



HAL
open science

Cylindrospermopsin accumulation and release by the benthic cyanobacterium *Oscillatoria* sp. PCC 6506 under different light conditions and growth phases.

Myriam Bormans, Marion Lengronne, Luc Brient, Charlotte Duval

► To cite this version:

Myriam Bormans, Marion Lengronne, Luc Brient, Charlotte Duval. Cylindrospermopsin accumulation and release by the benthic cyanobacterium *Oscillatoria* sp. PCC 6506 under different light conditions and growth phases.. *Bulletin of Environmental Contamination and Toxicology*, 2014, 92 (2), pp.243-247. 10.1007/s00128-013-1144-y . hal-00957621

HAL Id: hal-00957621

<https://hal-univ-rennes1.archives-ouvertes.fr/hal-00957621>

Submitted on 20 Mar 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Cylindrospermopsin accumulation and release by the benthic cyanobacterium
Oscillatoria sp PCC 6506 under different light conditions and growth phases**

Bormans Myriam^{1*}, Lengronne Marion¹, Brient Luc¹, Duval Charlotte²

¹UMR 6553 ECOBIO, CNRS Université de Rennes 1, 35042 Rennes, France

²UMR 7245 CNRS Museum National d'Histoire Naturelle, 75005 Paris, France

*corresponding author : myriam.bormans@univ-rennes1.fr, phone : +33 (0)2 2323 5979

ABSTRACT

We have studied the dynamics of cylindrospermopsin concentration [CYN] of a benthic cyanobacterium of the genus *Oscillatoria* under various light conditions over the different growth phases. The present study is the first one reporting on the effect of abiotic factors on the CYN accumulation and release by a benthic species. In particular we have measured the concentrations of both intracellular and extracellular [CYN]. We found that the total CYN content is highest during the exponential growth phase at intermediate light level ($10\mu\text{E m}^{-2} \text{s}^{-1}$) and during the stationary growth phase at more extreme lower and higher light levels. Our results also indicate that the amount of the extracellular form varied between 56 and 96% of the total CYN concentrations. We found no relationship between CYN content and growth rates. These results suggest many similarities with planktonic species but also highlight some differences.

KEYWORDS

Benthic cyanobacteria; cylindrospermopsin content; intracellular; extracellular; light intensity; growth phase

INTRODUCTION

Cyanobacteria are known to produce a variety of toxins, which have adverse effects on animals and humans (Codd et al. 2005). These toxins include microcystins, cylindrospermopsins, saxitoxins, nodularins, anatoxins and more recently beta-methylamino-L-alanine or BMAA. Although microcystin is the only toxin routinely monitored for health risks, there are an increasing number of studies reporting on the presence of all types of toxins in waterbodies. In particular, cylindrospermopsin (CYN) is rapidly being recognised as one of the globally problematic freshwater toxin due to its ever expanding distribution from tropical to temperate zones (Rücker et al. 2007 ; Brient et al. 2008 ; Sinha et al. 2012) and its ability to bioaccumulate in freshwater organisms (Kinnear 2010).

CYN is a tricyclic alkaloid cytotoxin produced by several cyanobacterial species : *Cylindrospermopsis raciborskii*, *Aphanizomenon ovalisporum*, *Aphanizomenon flos aquae*, *Anabaena bergii*, *Anabaena lapponica*, *Umezakia natans*, *Raphidiopsis curvata* and other unidentified species. Recently, several oscillatorioides have been identified as producing CYN as *Lyngbya wollei* (Seifert et al. 2007) and *Oscillatoria* sp. (Mazmouz et al. 2010). CYN is highly water soluble and is primarily present in extracellular form (Chiswell et al. 1999 ; Rücker et al. 2007). CYN degrades relatively slowly by photodegradation (Wörmer et al. 2010) and biodegradation (Wörmer et al. 2008) explaining relatively high extracellular concentrations in the water.

In the field, Wiedner et al. (2008) found that CYN was essentially produced as extracellular form during the decline of a bloom. Some laboratory experiments on *Cylindrospermopsis raciborskii* (Saker and Griffith 2000; Hawkins et al. 2001; Dyble et al. 2006) confirm this

tendency by demonstrating an increased accumulation of extracellular CYN during the stationary growth phase.

Abiotic factors that influence the accumulation and release of CYN have been studied both in the field and in laboratory experiments. The effect of temperature has been studied showing globally that an increase in temperature results in a decrease of CYN accumulation (Saker and Griffith 2000; Preußel et al. 2009; Cires et al. 2011). Nutrient concentrations also play a role in CYN accumulation. In studies with *Aphanizomenon ovalisporum*, an increase of CYN accumulation with limiting phosphorus conditions was shown (Bar Yosef et al. 2010) while higher CYN accumulation in cells grown in a culture medium with lower phosphorus concentration was also observed by Cires et al. (2011). Higher intracellular CYN concentration was also observed in controlled laboratory experiments with several strains of *Cylindrospermopsis raciborskii* in lower nitrogen medium (Saker and Neilan 2001). Light intensity has also been examined as a factor influencing CYN accumulation. Dyble et al. (2006) indicated an increase of total CYN concentration with an increase in light intensity between 18 and 75 $\mu\text{E m}^{-2} \text{s}^{-1}$ for the species *Cylindrospermopsis raciborskii* at 25°C. Similarly Cires et al. (2011) observed a significant increase of intracellular CYN between 15 and 60 $\mu\text{E m}^{-2} \text{s}^{-1}$ on the species *Aphanizomenon ovalisporum* at 28°C. However, according to Preußel et al. (2009), the impact of light intensity is temperature dependent, it increased at 20°C and decreased at 25°C for *Aphanizomenon flos aquae* for light ranging between 10 and 60 $\mu\text{E m}^{-2} \text{s}^{-1}$.

To our knowledge, all of the studies reported in the literature on the role of abiotic factors in CYN accumulation have involved planktonic species. As CYN concentration has recently been measured on benthic species, we examined how abiotic factors affect CYN accumulation and release by a benthic species. In particular, this study examines the role of light intensity and growth phase on both intracellular and extracellular CYN concentrations by the benthic *Oscillatoria sp.* PCC 6506 recently identified as producing CYN (Mazmouz et al. 2010) under unlimited nutrient conditions. We hypothesize that CYN content will vary as a function of stressed conditions, therefore we expect differences between growth phases and between light levels.

MATERIALS AND METHODS

Strain and cultures conditions

The strain PCC 6506 is a filamentous species with slightly flexed trichomes. The CYN producing strain *Oscillatoria sp.* PCC 6506 was provided by the Pasteur Culture Collection of Cyanobacteria (Pasteur Institute in Paris) as an axenic culture which was grown in a BG11 medium at 20°C in a 14h/10h light dark cycle. Three light intensities were tested (2, 10 and 20 $\mu\text{E m}^{-2} \text{s}^{-1}$) after a preliminary experiment identified that the culture was not growing at higher light intensity. Prior to the experiment each culture was pre-adapted for three weeks to its appropriate light intensity. All experiments were performed in triplicate batch cultures. An initial biomass of 10^5 cells mL^{-1} was used in each culture. Toxin concentrations were measured at the same time (day 2, day 23 and day 51) for all three light levels.

Biomass and growth kinetics

Oscillatoria sp. cell density was estimated by converting the OD at 750 nm into cell density (cells mL^{-1}) based on a linear relationship between these two parameters. The specific growth rates, μ (day^{-1}), were calculated during the exponential growth phase according to the following equation:

$$\mu = (\ln C_{t+\Delta t} - \ln C_t) / (\Delta t)$$

where C_t and $C_{t+\Delta t}$ represent the cell density (cells mL⁻¹) at times t and $t + \Delta t$, respectively.

Cell counts were obtained with a Nageotte cell on an optical microscope counting a minimum of 2000 cells. Optical density was measured twice a week to follow the evolution of the biomass, identify growth phases and calculate growth rates. The toxin concentration was tested prior to the experiment to determine appropriate volumes to sample. Samples of 1 mL were collected in each replicate during the different phases (latent, exponential and stationary) and analysed for intracellular and extracellular CYN concentrations.

Toxin analyses

Separation of intracellular and extracellular CYN was achieved by low vacuum filtration on a GF/F filter. The filtrate was stored at -18°C for extracellular analysis. For the intracellular form, the biomass recovered on the filter was dried for 3h at 35°C and resuspended in 5 mL of ultrapure water before being stored with the filter at -18°C. After thawing, extractions were performed according to Welker et al. (2002) and adapted for the specification of strain PCC 6506 as of Mazmouz et al. (2010). Samples (biomass and filter) were subjected to ultrasounds using a Sonics Vibracell probe for 60s at 130 W then placed on an orbital shaker at 300 rpm for 1h at room temperature. They were again subjected to 60s of ultrasounds and filtered on 0.45 µm nylon filter.

Concentrations were measured using a Cyindrospermopsin ELISA kit from Abraxis and analysed with a spectrophotometer ELX800UV from BioTEK instruments. The detection limit was estimated at 0.04 µg L⁻¹ and the range between 0.04 and 2.5 µg L⁻¹. Therefore, the samples had to be diluted with ultrapure water before analysis. Concentrations of both intracellular and extracellular forms are reported per cell.

Statistical analysis

Results are expressed as mean +/- standard deviations. One-way ANOVA for non equal variance followed by a post-hoc Tukey test was performed for all statistical tests during each of the growth phases with significance threshold set at 0.05.

RESULTS

A high correlation ($R^2 = 0.92$; $n = 45$ not shown) was found between optical density at 750 nm and cell numbers estimated by counting on an optical microscope. This relation, which was similar for the three light intensities, was used to infer the evolution of biomass. Growth rates and length of growth phases varied with light intensities. Growth rates were significantly different ($p < 0.05$, Tukey test) between each light intensity with the highest value of $0.136 \pm 0.003 \text{ day}^{-1}$ at $10 \mu\text{E m}^{-2} \text{ s}^{-1}$, an intermediate value of $0.118 \pm 0.005 \text{ day}^{-1}$ at $2 \mu\text{E m}^{-2} \text{ s}^{-1}$ and the lowest value of $0.100 \pm 0.003 \text{ day}^{-1}$ at $20 \mu\text{E m}^{-2} \text{ s}^{-1}$. Very low variance was observed between the triplicates for each modality. While the latent phase length (4 days) was similar for all light intensities, the exponential growth phase was longer at $10 \mu\text{E m}^{-2} \text{ s}^{-1}$ (40 days) and was equal to 30 days at the other two light intensities.

At the beginning of the experiment, during the latent phase, there was no significant difference in the CYN content (either forms) between the replicates cultured at different light intensities ($p > 0.05$, Tukey test). We observed an intracellular CYN concentration varying between 1.9 and 2.6 fg cell⁻¹ and an extracellular CYN concentration varying between 9.3 and 12.4 fg cell⁻¹ with over 80% as extracellular CYN (Fig.1).




Fig. 1 : Intracellular and extracellular CYN concentrations (fg cell^{-1}) as a function of light during the latent, exponential and stationary growth phase

During the exponential growth phase the highest proportion of the toxin was again in extracellular form at all light levels varying between 56 % at $2 \mu\text{E m}^{-2} \text{s}^{-1}$ to 84 % at $10 \mu\text{E m}^{-2} \text{s}^{-1}$. CYN concentrations were significantly lower ($p < 0,05$, Tukey test) at $2 \mu\text{E m}^{-2} \text{s}^{-1}$ for both forms with values of $1.40 \pm 0.04 \text{ fg cell}^{-1}$ as intracellular and $1.77 \pm 0.06 \text{ fg cell}^{-1}$ as extracellular form compared to the other light levels (Fig.1). During the exponential growth phase, we therefore observed a significantly lower ($p < 0,05$, Tukey test) total CYN content at $2 \mu\text{E m}^{-2} \text{s}^{-1}$ compared to the other two light levels.

During the stationary phase, again most of the toxin was in extracellular form with proportions varying with light levels between 89 % at $10 \mu\text{E m}^{-2} \text{s}^{-1}$ and 96 % at $2 \mu\text{E m}^{-2} \text{s}^{-1}$.

Both intracellular and extracellular CYN concentrations varied with light intensity and were significantly higher ($p < 0.05$, Tukey test) at $20 \mu\text{E m}^{-2} \text{s}^{-1}$ than at the other light levels. Total CYN content was maximal at $20 \mu\text{E m}^{-2} \text{s}^{-1}$ during the stationary phase with concentrations per cell of $19.6 \pm 2.4 \text{ fg cell}^{-1}$ (Fig.1). For a typical biomass observed in the field of $10^5 \text{ cells mL}^{-1}$, these values would be equivalent to $2 \mu\text{g L}^{-1}$, a value twice the recommended drinking water guideline of $1 \mu\text{g L}^{-1}$ (Humpage and Falconer 2003).

Although the maximum total CYN content during the exponential growth phase coincided with the maximum growth rate (at $10 \mu\text{E m}^{-2} \text{s}^{-1}$), the lowest CYN contents at $2 \mu\text{E m}^{-2} \text{s}^{-1}$ were not associated with the lowest growth rates. Therefore we did not observe a consistent relationship between growth rate and either intracellular or total CYN content (Fig. 2).




Fig. 2 : Total and intracellular CYN concentrations (fg cell⁻¹) versus growth rates (d⁻¹) during the exponential phase at different light intensities.

DISCUSSION

In this study, we showed a dominance of extracellular CYN content regardless of the growth phase and the light level for the benthic species *Oscillatoria sp.* Dominance of extracellular CYN (62%) for the same strain *Oscillatoria sp.* PCC 6506 was also observed by Mazmouz et al. (2010) after 11 days of growth at 20°C and in a nutrient rich medium and a constant light level of $15 \mu\text{E m}^{-2} \text{s}^{-1}$, which was likely to correspond to the exponential phase. The results presented here on the CYN content for a benthic species being mostly extracellular are in accordance with results for planktonic species, although very few studies have reported on

CYN release and presented extracellular concentrations values. Dominance of extracellular CYN was observed in the field in lakes dominated by the *Aphanizomenon* genus (Wiedner et al. 2008). In laboratory experiments, extracellular proportions varied with abiotic factors and species. Cires et al. (2011) reported over 60% of extracellular CYN at extreme temperature for *Aphanizomenon ovalisporum* and Preußel et al. (2009) reported extracellular CYN concentration up to 58% under stress conditions, suggesting that the release of CYN was stress induced. Increases in extracellular concentrations under stress conditions have also been reported for microcystin (MC) producers associated with high light (Wiedner et al. 2003) or grazing pressure (Zhang et al. 2010). Nevertheless, for MC producers, the majority of the toxin is in intracellular form except during cell lysis.

Although the CYN concentration was found to be mostly in extracellular form, we observed a growth phase effect. Our results suggest relatively higher intracellular CYN content during the exponential growth phase and higher extracellular CYN content during the stationary phase. Very few studies have examined the CYN accumulation and release during the stationary phase (Hawkins et al. 2001; Dyble et al. 2006). However, they all report an increase of extracellular form during the stationary phase in accord with an increase observed in the field at the end of the bloom in a seasonal dynamics (Wiedner et al. 2008). This characteristic seems therefore to be common to all CYN producers studied so far.

We observed also a light effect on CYN accumulation and release. A statistically significant lower concentration of extracellular CYN was observed at the intermediate light level during the stationary phase. Due to the higher biomass reached this result can be in part due to a dilution effect, which occurs when the growth rate is higher than the toxin production rate. Observed higher extracellular CYN concentration at lower and higher light levels are consistent with a more limiting environment associated with stressed conditions. Indeed, stress related release of extracellular CYN has been suggested under N limitation (Preußel et al. 2009). We also observed a light intensity effect with lower CYN content of both intracellular and extracellular form during the exponential phase at low light leading to a lower total CYN content. This result is similar to that of planktonic species, which tend to show higher accumulation of intracellular CYN concentration at higher light levels.

We did not observe a monotonic relationship between growth rate and either intracellular or total CYN concentration. This result is similar to published studies on other CYN producers, which do not show a consistent relationship between CYN concentration and μ . Indeed high CYN concentrations have been measured at high light intensities in *Cylindrospermopsis raciborskii* when growth rates were not at their maximum (Dyble et al. 2006). Similarly, Saker and Neilan (2001) found high intracellular CYN concentrations in low nitrogen *Cylindrospermopsis raciborskii* cultures while growth rates were suboptimal. There were no correlations between growth rates associated with variations of both temperature and light and either intracellular or total CYN contents in studies on *Aphanizomenon flos aquae* (Preußel et al. 2009; Cires et al. 2011). This lack of relationship between growth rate and CYN content contrasts with MC producers which often show a strong linear relationship between intracellular MC and μ (Orr and Jones 1998; Briand et al. 2012) suggesting different functional roles for these toxins.

In summary, we found that a benthic species produced predominantly extracellular CYN with an increase during the stationary phase in accord with planktonic CYN producers possibly indicating a stress response. No correlation was found between growth rate and CYN content again in a similar way to planktonic CYN producers' response. Hence the results presented in

this study, although preliminary, suggest many similarities with planktonic CYN producers. Further studies should include testing intra species variability amongst *Oscillatoria* as well as other benthic species before generalization can be attempted. As the CYN toxin is predominantly present in an extracellular form in the environment, its presence is particularly problematic in water bodies used for drinking water purposes or recreational activities. In view of major ecological implications from benthic CYN producers, which include grazing by benthic organisms bioaccumulating in the food chain, it is somewhat surprising that benthic cyanobacteria have received so little attention.

Acknowledgements

This work was financed by the National Agency for Security of Environment and Health (ANSES) through the TOXCYN project. The experiment was run in the phytotronic chambers of the experimental facilities ECOLEX of ECOBIO.

REFERENCES

- Bar-Yosef Y, Sukenik A, Hadas O, Viner-Mozzini Y, Kaplan A (2010) Enslavement in the water body by toxic *Aphanizomenon ovalisporum*, inducing alkaline phosphatase in phytoplanktons. *Curr Biol* 20:1557–1561
- Brient L, Lengronne M, Bormans M, Fastner J. 2008. First occurrence of cylindrospermopsin in freshwater in France. *Environ Toxicol* 24(4):415-420
- Briand E, Bormans M, Quiblier C, Salençon MJ, Humbert JF (2012) Evidence of the cost of the production of Microcystins by *Microcystis aeruginosa* under differing light and nitrate environmental conditions. *PlosOne* 7(1):e29981
- Chiswell RK, Shaw GR, Eaglesham G, Smith MJ, Norris RL, Seawright AA, Moore MR (1999) Stability of cylindrospermopsin, the toxin from the cyanobacterium, *Cylindrospermopsis raciborskii* : Effect of pH, temperature and sunlight on decomposition. *Environ Toxicol* 14:155–161
- Cires S, Wörmer L, Timon J, Wiedner C, Quesada A (2011) Cylindrospermopsin production and release by the potentially invasive cyanobacterium *Aphanizomenon ovalisporum* under temperature and light gradients. *Harmful Algae* 10(6):668–675
- Codd GA, Morrison LF, Metcalf JS (2005) Cyanobacterial toxins: risk management for health protection. *Toxicol Appl Pharmacol* 203:264–272
- Dyble J, Tester PA, Litaker RW (2006) Effects of light intensity on cylindrospermopsin production in the cyanobacterial HAB species *Cylindrospermopsis raciborskii*. *Afr J Mar Sci* 28:309-312
- Hawkins PR, Putt E, Falconer I, Humpage A (2001) Phenotypical variation in a toxic strain of the phytoplankter, *Cylindrospermopsis raciborskii* (Nostocales, Cyanophyceae) during batch culture. *Environ Toxicol* 16(6):480–487
- Humpage AR and Falconer IR (2003) Oral toxicity of the cyanobacterial toxin cylindrospermopsin in male Swiss albino mice: Determination of no observed adverse effect level for deriving a drinking water guideline value. *Environ Toxicol* 18(2):94–103
- Kinnear S (2010) Cylindrospermopsin : A decade of progress on bioaccumulation research. *Mar Drugs* 8:542–564

- Mazmouz R, Chapuis-Hugon F, Mann S, Pichon V, Méjean A, Ploux O (2010) Biosynthesis of cylindrospermopsin and 7-epicylindrospermopsin in *Oscillatoria* PCC 6506 ; Identification of the gene cluster and toxin analysis. *Appl Environ Microbiol* 76(15):4943–4949
- Orr P, Jones G (1998) Relationship between microcystin production and cell division rates in nitrogen-limited *Microcystis aeruginosa* cultures. *Limnol Oceanogr* 43(7):1604-1614
- Preußel K, Wessel G, Fastner J, Chorus I (2009) Response of cylindrospermopsin production and release in *Aphanizomenon flos-aquae* (cyanobacteria) to varying light and temperature conditions. *Harmful Algae* 8:645–650
- Rücker J, Stüken A, Nixdorf B, Fastner J, Chorus I, Wiedner C (2007) Concentrations of particulate and dissolved cylindrospermopsin (CYN) in 21 *Aphanizomenon* dominated lakes of North East Germany. *Toxicon* 50:800–809
- Saker ML, Griffith DJ (2000) Effects of temperature on growth and cylindrospermopsin content of seven isolates of *C. raciborskii* (Nostocales, Cyanophyceae) from water bodies in northern Australia. *Phycologia* 39:349-354
- Saker ML, Neilan BA (2001) Varied diazotrophies, morphologies, and toxicities of genetically similar isolates of *Cylindrospermopsis raciborskii* (Nostocales, Cyanophyceae) from Northern Australia. *Appl Environ Microbiol* 67(4):1839–1845
- Seifert M, McGregor G, Eaglesham G, Wickamasinghe W, Shaw G (2007) First evidence for the production of cylindrospermopsin and deoxy-cylindrospermopsin by the freshwater benthic cyanobacterium, *Lyngbia wollei* (Fallow ex Gomont) Spezialeand Dick. *Harmful Algae* 6:73–80
- Sinha R, Pearson LA, Davis TW, Burford MA, Orr PT, Neilan BA (2012) Increased incidence of *Cylindrospermopsis raciborskii* in temperate zones - Is climate change responsible? *Water Res* 46:1408–1419
- Welker M, Bickel H, Fastner J (2002) HPLC-PDA detection of cylindrospermopsin – opportunities and limits. *Water Res* 36:4659–4663
- Wiedner C, Visser PM, Fastner J, Metcalf JS, Codd GA, Mur LR. 2003. Effects of light on the microcystin content of *Microcystis* strain PCC 7806. *Applied and Environ Microb* 69:1475-1481
- Wiedner C, Rücker J, Fastner J, Chorus I, Nixdorf B (2008) Seasonal dynamics of cylindrospermopsin and cyanobacteria in two German lakes. *Toxicon* 52:677-686
- Wörmer L, Cires S, Carrasco D, Quesada A (2008) Cylindrospermopsin is not degraded by co-occurring natural bacterial communities during a 40-day study. *Harmful Algae* 7:206–213
- Wörmer L, Huerta-Fontela M, Cires S, Carrasco D, Quesada A (2010) Natural photodegradation of the cyanobacterial toxins microcystin and cylindrospermopsin. *Environ Sci Technol* 44(8):3002-3007
- Zhang X, Hu H, Men Y, Christoffersen KS. 2010. The effect of *Poteroiochromonas* abundance on production of intra and extracellular microcystin-LR concentration. *Hydrobiologia* 652:237-246