Pseudo-first-order reaction of chemically and biologically formed green rusts with HgII and C₁₅H₁₅N₃O₂: effects of pH and stabilizing agents (phosphate, silicate, polyacrylic acid, and bacterial cells)

P-Ph Remy, M. Etique, A. A. Hazotte, A-S Sergent, N. Estrade, C. Cloquet, K. Hanna, F. P. A. Jorand

► To cite this version:

P-Ph Remy, M. Etique, A. A. Hazotte, A-S Sergent, N. Estrade, et al.. Pseudo-first-order reaction of chemically and biologically formed green rusts with HgII and C₁₅H₁₅N₃O₂: effects of pH and stabilizing agents (phosphate, silicate, polyacrylic acid, and bacterial cells). Water Research, 2015, 70, pp.266-278. 10.1016/j.watres.2014.12.007 . hal-01153424

HAL Id: hal-01153424

<https://hal-univ-rennes1.archives-ouvertes.fr/hal-01153424>

Submitted on 22 Oct 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Pseudo-first-order reaction of chemically and biologically formed green
rusts with Hg^{II} and $\text{C}_{15}\text{H}_{15}\text{N}_3\text{O}_2$: effects of pH and stabilizing agents
(phosphate, silicate, polyacrylic acid, and bacterial cells)**

P.-Ph. Remy^{1, 2}, M. Etique^{1, 2}, A.A. Hazotte^{1, 2, §}, A.-S. Sargent^{1, 2}, N. Estrade^{3, 4#}, C. Cloquet^{3, 4},
K. Hanna^{5, 6}, and F.P.A. Jorand^{1, 2*}

¹Université de Lorraine, LCPME, UMR 7564, Institut Jean Barriol, Villers-lès-Nancy, F-54601, France

²CNRS, LCPME, UMR 7564, Institut Jean Barriol, Villers-lès-Nancy, F-54601, France

³CNRS, CRPG, UMR 7358, BP 20, Vandœuvre-lès-Nancy, F-54501, France

⁴Université de Lorraine, CRPG, UMR 7358, BP 20, Vandœuvre-lès-Nancy, F-54501, France

⁵ENSCR, CNRS, UMR 6226, CS 50837, Rennes Cedex 7, F-35708, France

⁶Université européenne de Bretagne, Rennes, F-35000, France

[§]Present address: LUNAM University, Subatech-LPGN, UMR 6457 & 6112, BP 92208, F-44322 Nantes Cedex 3, France

[#]Present address: Pacific Centre for Isotopic and Geochemical Research, EOAS, The University of British Columbia, 2207 Main Mall, Vancouver, British Columbia, V6T 1Z4, Canada.

*Corresponding author: frederic.jorand@univ-lorraine.fr; +33 (0)383 685 248

Keywords: Reductive transformation, layered double hydroxides, cationic mercury, methyl red, biogenic green rust, surface area concentration

Abstract

The kinetics of Hg^{II} and methyl red (MR) reduction by hydroxycarbonate green rust (GR1) and by hydroxysulfate green rust (GR2) were studied in the presence of naturally occurring organic and inorganic ligands (phosphate, polyacrylic acid, bacterial cells, silicate). The reducing ability of biogenic hydroxycarbonate green rust (GR1_{bio}), obtained after microbial reduction of lepidocrocite by *Shewanella putrefaciens*, was also investigated and compared to those of chemically synthesized GR1 and GR2 (GR1_{ab} and GR2_{ab}). Pseudo first-order rate constants (k_{obs}) of Hg^{II} reduction (at pH 7, 8.2, and 9.5) and MR reduction (at pH 7) were determined and were normalized to the structural Fe^{II} content of GRs (k_{FeII}) and to the estimated concentration of surface Fe^{II} sites (k_{s}). The k_{s} values ranged from $0.3 \text{ L mmol}^{-1} \text{ min}^{-1}$ to $43 \text{ L mmol}^{-1} \text{ min}^{-1}$ for the Hg reduction, and from $0.007 \text{ L mmol}^{-1} \text{ min}^{-1}$ to $3.4 \text{ L mmol}^{-1} \text{ min}^{-1}$ for the MR reduction. No significant discrepancy between GR_{ab} and GR_{bio} was observed in term of reactivity. However, the reduction kinetics of MR was generally slower than the Hg^{II} reduction kinetics for all tested GRs. While a slight difference in Hg^{II} reduction rate was noted whatever the pH values (7.0, 8.2, or 9.5), the reduction of MR was significantly affected in the presence of ligands. A decrease by a factor of 2 – 200, depending on the type of ligand used, was observed. These data give new insights into the reactivity of GRs in the presence of co-occurring organic and inorganic ligands, and have major implications in the characterization of contaminated systems as well as water treatment processes.

1. Introduction

Green rusts (GRs) are mixed ferrous–ferric hydroxides which have a layered structure characterized by alternating positively charged hydroxide layers $\{\text{Fe}^{\text{II}}_{(1-x)}\text{Fe}^{\text{III}}_x(\text{OH})_2\}^{x+}$ and hydrated anionic interlayers $\{(x/n)\text{A}^{n-} \cdot m\text{H}_2\text{O}\}^{x-}$. Two types of GRs are distinguished: green rust 1 (GR1) and green rust 2 (GR2) containing either planar or spherical anions (*e.g.* CO_3^{2-} , Cl^-), and non-planar anions (*e.g.* SeO_4^{2-} , SO_4^{2-}), respectively (Génin *et al.*, 2006). GRs are present in the environment as corrosion products of Fe-based materials (Refait *et al.*, 2003), and as minerals (*e.g.* fougèrite) in hydromorphic soils (Génin *et al.*, 1998), in ground water (Christiansen *et al.*, 2009), or in suspended matter of stratified lakes (Zegeye *et al.*, 2012). GRs are described as highly reactive compounds, especially in reducing several organic and inorganic contaminants (Myneni *et al.*, 1997; Erbs *et al.*, 1999; Loyaux-Lawniczak *et al.*, 2000; Hansen *et al.*, 2001; Williams and Scherer, 2001; Lee and Batchelor, 2002; O’Loughlin *et al.*, 2003; Elsner *et al.*, 2004; O’Loughlin *et al.*, 2004; Mitsunobu *et al.*, 2008; Kone *et al.*, 2009) and their roles in the biogeochemical redox cycling of iron and other elements are of growing interest to the scientific community (*e.g.* : Carlson *et al.*, 2012; Zegeye *et al.*, 2012). Although their reactivity has been widely investigated with chemically synthesized GR2 (SO_4^{2-}), the reactivity of GR1, microbially synthesized GR1/GR2, and GR2 in the presence of cells or polymers has been scarcely reported.

GRs can be synthesized at the laboratory scale by a partial oxidation of ferrous salts or by a coprecipitation of Fe^{II} with Fe^{III} salts in aqueous solution (Schwertmann and Fechter, 1994; Génin *et al.*, 1998; Bocher *et al.*, 2004). They can also be formed by bacterial activities via a bioreduction of ferrihydrite or lepidocrocite under anoxic conditions (Fredrickson *et al.*, 1998; Ona-Nguema *et al.*, 2002) or by an indirect Fe^{II} oxidation by biogenic nitrite (Etique *et al.*, 2014a). In the natural environment, the formation of GRs is sometimes incorrectly considered as a biological process, whereas the reduction of Fe^{III} species or the oxidation of

Fe^{II} species are only due to an enzymatic reaction. The resulting coprecipitation leading to the formation of GR in the extracellular medium is only based on a chemical process. Furthermore, no difference in structure and composition between chemically and biologically formed GRs have been evidenced, except for the size of crystals, which are larger for the biological GR than for its chemical counterpart (Zegeye *et al.*, 2005), and for the presence of extracellular-polymeric substances (EPS) and/or bacterial cells surrounding biological GR crystals (Zegeye *et al.*, 2010; Jorand *et al.*, 2013). These EPS and cells are perceived as affecting GR reactivity, which might explain the higher stability of the biogenic GR (for several years in aqueous phase) as compared to the chemical one. However, this phenomenon remains poorly studied and barely understood. Although recent work has demonstrated that the extracellular polymers (EPS from activated sludge in waste water treatment) significantly affect the reactivity of GR1 towards anionic inorganic pollutants (NO₃⁻), probably by “screen effect” on reactive sites (Zegeye *et al.*, 2014), knowledge on this reactivity with metallic or organic contaminants is essential to provide greater insight into applied or natural processes in which GRs are involved (*e.g.* Su and Puls, 2004; Bearcock *et al.*, 2011).

For that purpose, the reactivity of a biologically/chemically synthesized hydroxycarbonate GR1 (GR1_{bio}/GR1_{ab}), and chemically formed sulfate GR2 (GR2_{ab}) towards Hg^{II} and C₁₅H₁₅N₃O₂ (methyl red MR) was investigated in presence of various stabilizing agents. Previous works have already studied the reactivity of GR2_{ab} towards Hg^{II} and MR (O’Loughlin *et al.*, 2003; Kone *et al.*, 2009), but to the best of our knowledge, the reactivity of biogenic GRs towards these two contaminants has not yet been documented. In anoxic environments, green rust phases are represented by the fougèrite mineral, which is a hydroxycarbonate green rust (GR1) (Génin *et al.*, 1998; Mills *et al.*, 2012). Therefore, the most representative iron phase of this fougèrite mineral would be the GR1_{bio}. GR2_{ab} is known to be relatively stable as long as common anoxic conditions are preserved at pH ~7 (Ruby *et*

al., 2006), while GR1_{ab} is perceived as unstable iron hydroxides rapidly turning into a mixture of magnetite (Fe^{II}Fe^{III}₂O₄) and siderite (FeCO₃) (Taylor *et al.*, 1985; Benali *et al.*, 2001). However, under strict anoxic conditions and neutral to slightly alkaline pH (7 – 9) (this work), GR1_{ab} can be preserved and stored for several days (Etique *et al.*, 2014b). Nonetheless, many experiments were performed with GR1_{ab} supplemented with PO₄³⁻ that prevents any further undesirable “transformation” (Bocher *et al.*, 2004). Silicate, quartz, organic polymers, and bacterial cells – compounds which are commonly found in environmental settings or in water or soil remediation processes – were also described to inhibit/prevent GRs dissolution/transformation (Zegeye *et al.*, 2010; O’Loughlin *et al.*, 2010; Sergent *et al.*, 2011; Jorand *et al.*, 2013). Thus, to test the ability of these “stabilizers” to affect the reactivity of GR1 and GR2, the reduction of Hg^{II} and MR was carried out in the presence of phosphates, silicates, polyacrylic acid (paa), and bacterial cells, at pH values allowing a stability of GRs (pH > 7). Control experiments were also performed to evaluate the contribution of aqueous Fe^{II} and bacterial cells to the transformation of target compounds. The depletion of mercury was monitored over time using ICP-AAS. The MR reduction was monitored by UV-Visible spectrophotometry.

2. Chemical background

2.1. Mercury and GR

Mercury has two main stable oxidation states: Hg⁰ and Hg^{II}. Elemental Hg⁰ is gaseous under environmental conditions, and under environmentally relevant pH values Hg^{II} exists in pure water in aqueous form as Hg²⁺ and mercury dihydroxide (Hg(OH)₂) (1) with a pK_a value of 6.09 (Pourbaix, 1963):

Thus, the predominant mercury species with pH higher than or equal to 6.09 is $\text{Hg}(\text{OH})_2$, and the reduction half-reaction equation (2) is:

The reduction of Hg^{II} is effective in the presence of solid Fe^{II} but insignificant with aqueous Fe^{II} (Charlet *et al.*, 2002). The mixed Fe^{II} - Fe^{III} minerals, such as sulfate GR2, are oxidized by Hg^{II} to magnetite (O'Loughlin *et al.* 2003). The $\text{Fe}^{\text{II}}/\text{Fe}^{\text{III}}$ ratio of hydroxycarbonate GR1 can vary from 1/2 to 2/1 (trébeurdenite and fougèrite, respectively) (Mills *et al.*, 2012), while it is equal to 2/1 for sulfate GR2, as far as we know (Refait *et al.*, 2006). The oxidation of GR1 and GR2 to magnetite by $\text{Hg}(\text{OH})_2$ involves an exchange of 2 electrons and is given by equations (3) and (4):

Due to the presence of bicarbonate, the oxidation of GR1 leads to a lower production of H^+ than that of GR2.

2.2. Methyl red and GR

Methyl red (MR) or 2-(N,N-dimethyl-4-aminophenyl)azobenzenecarboxylic acid ($\text{C}_{15}\text{H}_{15}\text{N}_3\text{O}_2$) is an azo dye with one azo group ($-\text{N}=\text{N}-$) used as a common pH indicator with a $pK_a = 5.1$. As other azo dyes, MR is a recalcitrant pollutant causing a significant burden on the environment. The reduction of MR is known to be achieved with the cleavage of the azo bond (Khalid *et al.*, 2008; Kone *et al.*, 2009), leading to the formation of two aromatic amines (carboxyanilic acid: $\text{R}_1\text{-NH}_2$ and *p*-dimethylaminoaniline: $\text{R}_2\text{-NH}_2$) as:

Thus, 4 $\text{Fe}^{\text{II}}_{\text{GR}}$ are theoretically needed to reduce MR and, assuming that GR is oxidized into FeOOH , the reduction equations can be written as:

3. Materials and methods

3.1. Chemicals

All solutions were prepared using ACS grade chemicals (except when specified) and 18.2 MΩ cm N₂-purged pure water (Purelab Option-Q, Elga LabWater, Antony, France). The mercury solution (750 μM) was prepared from an HgCl₂ salt 99.5% (1166745, Merck) in HCl 0.5 M (30721-2.5L, Sigma-Aldrich) and the pH was adjusted to 7.0 ± 0.1 with NaOH 1M. Ferrous sulfate heptahydrate (FeSO₄•7H₂O, 99%), ferrous chloride tetrahydrate (FeCl₂•4H₂O, 99%), and sodium phosphate (Na₂HPO₄•7H₂O) were purchased from Sigma-Aldrich. Polyacrylic acid (C₃H₄O₂)_n was provided from a stock solution of 2 g L⁻¹ (Mw 2,000 g mol⁻¹, Aldrich, 323667). The methyl red (C₁₅H₁₅N₃O₂) was provided by Sigma-Aldrich (98%, 32654-25G).

The mercury concentration present in each reactant (0.3 – 1.9 ppb) was measured as blank (Table S1).

3.2. Synthesis of GRs

Abiotically formed GRs were synthesized under O₂-free conditions in an anaerobic chamber scavenging oxygen with palladium catalysts and H₂ (N₂/H₂, 95/5) (Coy Laboratory Products Inc.[®]). The GR1_{ab} was obtained using a coprecipitation method as described by Bocher *et al.* (2004) by adding a mixture of Fe^{II} and Fe^{III} salts to a mixture of sodium hydroxide and sodium carbonate. Ferrous sulfate heptahydrate FeSO₄•7H₂O and ferric sulfate pentahydrate Fe₂(SO₄)₃•5H₂O were dissolved in 40 mL of pure water. A [Fe^{II}]/[Fe^{III}] ratio of 2 was chosen with {[Fe^{II}]+[Fe^{III}]} = 0.5 M. Magnetic stirring (500 rpm) ensured fast and

complete dissolution. Di-sodium carbonate Na_2CO_3 ($[\text{CO}_3^{2-}] = 0.5 \text{ M}$) was dissolved in 40 mL of a 1 M NaOH solution in order to have $[\text{OH}^-]/\{[\text{Fe}^{\text{II}}]+[\text{Fe}^{\text{III}}]\} = 2$. The carbonate solution was added to the solution of iron under the same conditions of magnetic stirring, and a bluish-green precipitate appeared immediately. GR1_{ab} was used within a week of its formation to test its reactivity with Hg^{II} and MR, or was stabilized by addition of extra $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$ salt $\{[\text{PO}_4]_0/[\text{Fe}]\} = 0.8\%$ (*i.e.* 4 mM PO_4^{3-}) ($\text{GR1}_{\text{ab+p}}$) or silicic acid (H_4SiO_4 4 mM) ($\text{GR1}_{\text{ab+Si}}$), or paa (500 mg L^{-1}) ($\text{GR1}_{\text{ab+paa}}$). A $\{[\text{CO}_3^{2-}]/[\text{SO}_4^{2-}]\}$ ratio of 1 was chosen to prevent the formation of sulfate GR2_{ab} as described by Bocher *et al.* (2004).

The abiotic sulfate GR2 (GR2_{ab}) was synthesized using the coprecipitation method developed by Ruby *et al.* (2003). Briefly, 20 millimoles of an iron solution, containing 13.3 millimoles Fe^{II} (3.7 g of $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$) and 6.7 millimoles Fe^{III} (1.63 g of $\text{Fe}_2(\text{SO}_4)_3 \cdot 5\text{H}_2\text{O}$), were added to 400 mL of pure water. Iron salts were precipitated by addition of NaOH (50 mL, 40 mM) with a syringe and under magnetic stirring to precipitate all the iron salts ($n\text{NaOH} / n\text{Fe} = 2$).

The biotic carbonate GR1 (GR1_{bio}) was produced according to Zegeye *et al.* (2007) by incubation of *S. putrefaciens* CIP 8040^T (2.5×10^9 cells mL^{-1}) at 30°C in a basal medium containing 160 mM of sodium formate (HCOONa) and 300 mM of lepidocrocite ($\gamma\text{-FeOOH}$) as the sole electron donor and acceptor, respectively.

All GRs synthesized were washed in O_2 -free deionized water by centrifugation (5 min, $10,000 \times g$) and the concentration of GRs was determined by the spectrophotometric ferrozine method measuring the concentration of Fe^{II} and Fe^{total} (Viollier *et al.*, 2000) after HCl extraction (6 M) in a Tecan Infinite M200 PRO microplate reader. GRs were characterized by X-ray diffraction (XRD), transmission electron microscopy (TEM), and transmission Mössbauer spectroscopy (TMS).

3.3. Experimental set-up of Hg^{II} and methyl red reduction by GRs

The Hg^{II} solution (5 μM , initial starting concentration) was introduced into a 250 mL glass reactor (borosilicated glass, Schott) containing 150 mL of pure water with a GR suspension (500 μM , $\text{Hg}/\text{GR} = 1/100$). Initially, the glass reactor pH was adjusted to 7.0, 8.2, and 9.5 with NaOH 1N or HCl 0.5N. The glass reactor was continuously N_2 -purged and stirred (300 rpm) in order to outgas Hg^0 and prevent any oxidation from ambient air. One milliliter of the unfiltered suspension (*i.e.* with dissolved and adsorbed Hg) was sampled at regular time intervals and was immediately dissolved in an acidic solution of HNO_3/HCl (500 μL HNO_3 at 65% with 500 μL HCl at 30%, w/v), before being diluted ten times. This acidic digestion was performed in order 1) to stop the reaction and preserve the mercury speciation, and 2) to dissolve the mineral phase and to release the potential adsorbed Hg fraction in the aqueous phase. Thus, the potential adsorption of Hg on GR surfaces should not bias the determination of reduction kinetic. Moreover, the formation of Hg^0 was qualitatively checked by trapping it during the reaction between Hg^{II} species and green rust in a solution of KMnO_4 50 mM (in H_2SO_4 10%), where the elemental volatile mercury was re-oxidized into Hg^{II} under acidic conditions (data not shown).

The methyl red (MR) solution was prepared in pure water and the pH was adjusted to 7.0 ± 0.1 with NaOH 1N. The MR solution (final concentration $\sim 15 \mu\text{M}$) was introduced in a glass batch reactor containing 150 mL of pure water with a GR suspension (500 μM , $\text{MR}/\text{GR} \sim 1/33$). During the reaction, the medium was continuously stirred (300 rpm) and 1 mL samples were removed over time after adjusting the pH to 7.0 ± 0.1 in order to avoid shifting the maximum peak (at 430 nm).

Control experiments were also carried out for Hg^{II} and MR on the same terms as before, respectively. The $\text{Fe}^{\text{II}}_{\text{aq}}$ control was performed (0.2 mM, pH 7) to assess the reactivity of Hg^{II} or MR with aqueous Fe^{II} versus structural Fe^{II} . Such amounts of Fe^{II} were chosen assuming

that ~0.2 mM of $\text{Fe}^{\text{II}}_{\text{aq}}$ (*i.e.* passing through 0.2 μm) was measured in a suspension of GR (not shown). A bacterial control with 6×10^5 cells mL^{-1} of *S. putrefaciens* was performed to evaluate the effect of bacteria in the GR1_{bio} suspension. The cell density selected was similar to what is required for 500 μM GR1_{bio}.

3.4. Analyses

The Hg concentration was determined with a Milestone DMA-80 Direct Mercury Analyzer (Milestone GmbH, Germany). Its unique processing of samples by thermal decomposition, amalgamation, and atomic absorption spectrometry provides a direct analysis of the sample matrices.

The MR concentration was measured by a spectrophotometric analytical method (Cary 60 UV-Vis, Agilent Technologies). A broad band in the visible MR spectrum was characterized at 430 nm in pH 7 (Fig. S1) and was assigned to the azo band (Hou *et al.*, 2007). To avoid artifacts caused by suspended particles, short centrifugations were performed (2 min, $14,000 \times g$) before determining MR concentration. The sorption of MR by GR was not considered here, as we assumed it to be negligible as previously shown (Kone *et al.*, 2009).

XRD data were collected with a D8 Bruker diffractometer, equipped with a monochromator and a position-sensitive detector. The X-ray source was a Co anode ($k = 0.17902$ nm). The diffractogram was recorded in the 3-64 2θ range, with a 0.0359° step size and collecting time of 3 s per point.

TMS analyses were performed using a constant-acceleration spectrometer with a 50 mCi source of ^{57}Co in Rh. The spectrometer was calibrated with a 25 μm foil of α -Fe at room temperature. The cryostat consisted of a closed cycle helium Mössbauer cryogenic workstation with vibration isolation stand manufactured by Advance Research Systems®. Helium exchange gas was used to thermally couple the sample to the refrigerator, allowing

variable temperature operations from 7 to 300 K. Computer fittings were done using Lorentzian-shape lines.

Transmission electron microscopy was conducted using CM20/STEM Philips coupled with an energy dispersive X-ray system (EDX) using a voltage of 200 kV. One drop of the suspension was laid on an amorphous carbon-coated grid and loaded into the analysis holder of the microscope under 10^{-8} Torr vacuum.

The surface area of GRs was determined by multipoint N_2 -BET analysis using a Coulter (SA113) surface area analyzer. Prior to specific surface area determination, GR suspensions were dried by vacuum desiccation for 48 h.

4. Results and discussion

4.1. Characterization of GRs

The greenish solid phase formed by either the microbial reduction of lepidocrocite (γ -FeOOH) by *S. putrefaciens* or chemical synthesis was characterized by XRD (Fig. 1a) and the d-values obtained were summarized in Table S2. The carbonate GR1 was identified as a major secondary iron mineral produced by the bacterial activity (GR1_{bio}) and as a main product from the coprecipitation method in the presence of carbonate and stabilized with phosphate (GR1_{ab+P}) or not stabilized (GR1_{ab}), which is in close agreement with previous studies (Drissi *et al.*, 1995; Zegeye *et al.*, 2007). TEM images and electron diffraction patterns were also performed and hexagonal crystals of GR1 were observed in various sizes from 5 μ m to 10 μ m for GR1_{bio} and from 100 nm to 300 nm for GR1_{ab+P} (Fig. 1b, c and d). This is consistent with the values reported in previous investigations (Ona-Nguema *et al.*, 2002; Zegeye *et al.*, 2014).

The formation of hydroxysulfate GR2 (GR2_{ab}) by the coprecipitation method from Ruby *et al.* (2003) was confirmed both by the d-values from the XRD diffractogram and by

the electron diffraction pattern, which is in agreement with the literature (Simon *et al.*, 1997) (Fig. 1, Table S2). Hexagonal particles of GR2_{ab} were observed by TEM with sizes ranging from 200 to 400 nm, which is in line with what Zegeye *et al.* (2005) had reported.

All the GRs formed were quasi-free of crystallographic impurities at a discernible level of XRD (Fig. S2). The Fe^{II}/Fe^{III} ratio was determined by TMS or by ferrozine method on solids (Fig. S3, Table S3): $1.2 \text{ (GR1}_{ab+P}) \leq 1.6 \text{ (GR1}_{bio}) \leq 2.0 \text{ (GR2}_{ab}) \leq 2.2 \text{ GR1}_{ab}$, and the specific surface area (SSA) was also measured by N₂-BET analysis (Table 1): $13 \text{ m}^2 \text{ g}^{-1} \text{ (GR1}_{bio}) \leq 14 \pm 2 \text{ m}^2 \text{ g}^{-1} \text{ (GR2}_{ab}) \leq 31 \pm 6 \text{ m}^2 \text{ g}^{-1} \text{ (GR1}_{ab+P} \text{ and GR1}_{ab})$ and also expressed as surface area concentration (A) in Table 1. These SSA values of green rusts were slightly lower than those obtained by Williams and Scherer (2001): $47 \pm 7 \text{ m}^2 \text{ g}^{-1}$. This fluctuation of SSA of green rusts depends on the synthesis method, which influences the crystal size. Moreover, the delay between its synthesis and its use modifies the crystal size probably by an Ostwald ripening.

Having determined SSA for each GR, the concentration of the surface Fe^{II} sites available can be reasonably estimated since the reactivity of GRs is based on this value. Here, an example is given for GR1_{bio} with an initial concentration of 500 μM and with the A value of $4.2 \text{ m}^2 \text{ L}^{-1}$ (Table 1). Knowing that the average common value of the surface site density is 5 Fe^{II} sites per nm^2 for iron oxides (Williams and Scherer, 2001), the number of Fe^{II} sites per nm^2 and per L of GR1_{bio} was determined at $2.1 \times 10^{19} \text{ Fe}^{\text{II}} \text{ sites nm}^{-2} \text{ L}^{-1}$ (i.e. $5 \text{ Fe}^{\text{II}} \text{ sites nm}^{-2} \times 4.2 \times 10^{18} \text{ nm}^2 \text{ L}^{-1}$). The total concentration of Fe^{II} sites on GR1_{bio} surface was therefore estimated at 34.8 μM (Table 1).

4.2. Kinetics of Hg^{II} reduction by GRs

Regardless the GR species, the Hg^{II} concentrations decreased rapidly according to an exponential decay at neutral and alkaline pH values (7.0, 8.2, and 9.5) (Fig. 2). Since GRs were present in great excess, their concentrations are assumed to have been constant

throughout the reaction and the decrease in Hg concentration can be described by a pseudo first-order reaction (equations 8, 9):

$$d[\text{Hg}]_t/dt = -k_{\text{obs}} \times [\text{Hg}]_t \quad (8)$$

$$[\text{Hg}]_t = [\text{Hg}]_0 \times \exp(-k_{\text{obs}} \times t) \quad (9)$$

where k_{obs} is the pseudo first-order rate constant (min^{-1}), and $[\text{Hg}]_t$ and $[\text{Hg}]_0$, the total concentrations of cationic mercury at time t and 0 , respectively. The fitting by a pseudo first-order reaction is consistent with other studies with GR and various oxidants such as selenite, CCl_4 , nitrate, and Cr^{VI} (Erbs *et al.*, 1999; Hansen *et al.*, 2001; Williams and Scherer, 2001; Etique *et al.*, 2014b). The values of k_{obs} were determined from equation (9) by plotting $-\ln([\text{Hg}]_t/[\text{Hg}]_0)$ as a function of time t . To take into account the Fe^{II} content of GRs (Note S1), k_{obs} values were normalized to Fe^{II} content ($k_{\text{Fe}^{\text{II}}}$, $\text{L mmol}^{-1} \text{min}^{-1}$) (Note S2), but it does not significantly affect the hierarchy as regards the kinetic rate constants (Table 1). Similarly, a k_{obs} normalization to the concentration of Fe^{II} sites on the green rust surface (k_{S} , $\text{L mmol}^{-1} \text{min}^{-1}$) (Note S3) does not seem to influence significantly the decrease of the Hg reduction rate from pH 7.0 to pH 9.5 (Table 1). However, at pH 8.2 the k_{S} values clearly highlight the increase of Hg reduction kinetics when $\text{Hg}^{\text{II}}/\text{GR}$ ratio increases from 1/80 to 1/0.8 for GR1_{bio} and from 1/100 to 1/10 for GR1_{ab} (Table 1, Fig. S4). Thus, this approach with k_{S} appeared to be more relevant compared to the expressions of kinetics rate constants in k_{obs} and $k_{\text{Fe}^{\text{II}}}$.

Control tests were performed with aqueous Fe^{II} (0.2 mM, pH = 7.0) or with bacteria (pH = 7.0) and no significant decrease in Hg concentration was observed over time ($k_{\text{obs}} < 10^{-3} \text{ min}^{-1}$) (Fig. S5a). This indicates that the depletion of Hg is neither due to a reduction reaction with the aqueous Fe^{II} , nor to sorption on glass walls. Similarly, bacteria associated with GR1_{bio} do not appear to be responsible for the Hg^{II} decrease. Thus, *S. putrefaciens* is not able, in our experimental conditions, to reduce Hg^{II} . It is worth noting that other iron reducers such as *S. oneidensis* MR-1 or *Geobacter* spp are able to reduce Hg^{2+} especially with lower

amounts of Hg^{II} ($0.15 \mu\text{M}$) and under growth conditions (Wiatrowski *et al.* 2006). In the present work, *S. putrefaciens* was in stationary growth phase and no electron donor or carbon source were added (non-growth conditions).

These results demonstrate that GRs reduced effectively Hg^{II} to Hg^0 . Compared to other Fe^{II} solids, these reaction kinetics are of the same order of magnitude as those induced by magnetite (Wiatrowsky *et al.*, 2006) or by the Fe^{II} species sorbed onto phlogopite (Charlet *et al.*, 2002). O'Loughlin *et al.* (2003) have reported slightly higher rates of Hg^{II} reduction by GR2, as they reduced 98.7% of the initial Hg^{II} (6.4 mmol L^{-1} of GR2_{ab} with $400 \mu\text{mol L}^{-1} \text{Hg}^{\text{II}}$, $\text{GR}/\text{Hg}^{\text{II}} = 16$) after 30 min. Here, we report a thorough reduction of Hg^{II} in less than an hour (with a ratio 6 times higher $\text{GR}/\text{Hg}^{\text{II}} = 100$ as compared to O'Loughlin *et al.*, 2003). Nevertheless, it cannot be excluded that part of Hg^0 could still be trapped in the solid phase, which would underestimate the kinetics of Hg^{II} reduction (Pasakarnis *et al.*, 2013).

4.2.1. Evolution of the Fe^{II} concentration and determination of the stoichiometry of the reduction of Hg^{II} species by GRs

The green rust reactivity was also evaluated by measuring the evolution of the total Fe^{II} concentration of the suspension. Among all experiments, only 3 runs present a detectable decrease of the $\text{Fe}^{\text{II}}/\text{Fe}_{\text{tot}}$ ratio in the presence of Hg^{II} species for a ratio $\text{Hg}^{\text{II}}/\text{GR}$ of 1/10 (GR1_{ab}) (Fig. S4a), 1/8 and 1/0.8 (GR1_{bio}) (Fig. S4b). For all the other experiments, the Fe^{II} concentration did not vary significantly since GR was used in large excess in comparison to the initial amount of Hg^{II} species ($5 \mu\text{M}$). During the reduction of mercury, the $\text{Fe}^{\text{II}}/\text{Fe}_{\text{tot}}$ ratio lowered of 0.06 units for GR1_{ab} (1/10) and GR1_{bio} (1/8) (Fig. S4). However, a sharp decrease from 0.67 to 0.46 was noted for the 1/0.8 ratio of $\text{Hg}^{\text{II}}/\text{GR1}_{\text{bio}}$ (Fig. S4b). Thus, Fe^{II} species of GR were oxidized to Fe^{III} species during the reduction of Hg^{II} species, and this chemical observation was confirmed by the characterization of magnetite as end product (Fig. S6).

Moreover, the stoichiometry of the reaction between Fe^{II} species and Hg^{II} species was determined at 1/10, 1/8 and 1/0.8 ratios of $\text{Hg}^{\text{II}}/\text{GR}$ (Fig. 3). The expected ratio $\text{Hg}^{\text{II}}:\text{Fe}^{\text{II}}_{\text{GR}}$ corresponding to the amount of consumed Hg^{II} ions over the amount of oxidized Fe^{II} species during the reduction of mercury by green rust is 1:2 according to the equation (3). This theoretical ratio of $\text{Hg}^{\text{II}}:\text{Fe}^{\text{II}}_{\text{GR}}$ was confirmed by the determination of the 1:1.9 experimental ratio (Fig. 3). Thus, this is in agreement (95 %) with the stoichiometry of the reaction involving the oxidation of 2 moles of ferrous iron per 1 mole of Hg^{II} reduced species.

4.2.2. Effect of pH on GR reactivity

In aqueous systems, GR particles act as Lewis acid and coordinate water or hydroxyl groups. This is why the particles surface chemistry is highly dependent of pH value. Recently, the point of zero charge (pzc) of GRs (GR1 and GR2) has been accurately determined at 8.3 ± 0.1 (Guilbaud *et al.*, 2013). Therefore, at $\text{pH} < 8.3$, the surface of GR particles is protonated, leading to a positive surface charge (Fig. 4); at $\text{pH} > 8.3$, the GR surface is negatively charged (Fig. 4). The evolution of the net surface charge of GR with pH may affect the rate and the extent of the reduction of contaminants such as chromate or mercury. Williams and Scherer (2001) demonstrated that the reduction of anionic pollutants such as CrO_4^{2-} was enhanced by a sorption onto positively charged surface of GR. Thus, the most significant reduction of Hg^{II} should be expected for a negatively charged surface of GRs ($\text{pH} > 8.3$), where the electrostatic interactions are more favorable for Hg^{2+} sorption.

However, the predominant phase of mercury at pH of this study is $\text{Hg}(\text{OH})_2$ (Fig. 4). At all tested pH values and whatever GRs used (GR2_{ab}, GR1_{bio}, and GR1_{ab+p}), the highest value of k_{obs} ($\sim 4 \times 10^{-2} \text{ min}^{-1}$) was obtained at pH 7.0, when GRs have a positive surface charge (Fig. 2a, Table 1). This would suggest, as it was previously considered for the reaction of magnetite with Hg^{II} (Wiatrowsky *et al.*, 2009), that hydroxyl groups of $\text{Hg}(\text{OH})_2$ could react with protonated GR particles, thus enhancing the sorption of mercury at the surface of GR

followed by a reduction into Hg^0 . Indeed, when pH values rise from 7.0 to 8.2 (or 9.5), the values of k_{obs} for $\text{GR1}_{\text{ab+P}}$, GR1_{bio} , and GR2_{ab} decrease by a factor of ~20%, 30%-50%, and 60%-70%, respectively (Table 1). GR reactivity at alkaline pH can be ranked as: $\text{GR1}_{\text{ab+P}} > \text{GR1}_{\text{bio}} > \text{GR2}_{\text{ab}}$. No significant difference in k_{obs} values was noted for each type of GRs between pH 8.2 and 9.5 (Table 1). Thus, the reduction of $\text{Hg}(\text{OH})_2$ by GR would be favored at $\text{pH} < 8.3$ (Fig. 4). Therefore, the fluctuation of the surface charge of GR in the function of pH governs the adsorption of $\text{Hg}(\text{OH})_2$ and its reduction.

Coupled with the adsorption/reduction of $\text{Hg}(\text{OH})_2$, a decrease in pH was observed for the reaction with GR2_{ab} , whereas pH values remained almost constant with $\text{GR1}_{\text{bio}}/\text{GR1}_{\text{ab+P}}$ (Table S4). These observations are consistent with equations (5) and (6), since the reduction of Hg^{II} is associated with the production of two H^+ for GR2 instead of one H^+ for GR1 (due to the presence of bicarbonate at the pH range values investigated).

4.2.3. Minor impact of cells and phosphate on the Hg^{II} reduction

Phosphate and bacterial cells (as bacterial bodies with polymeric substances) are known to improve the stability of GRs (Bocher *et al.*, 2004; Jorand *et al.*, 2013) and to affect their reactivity towards nitrate (Etique *et al.*, 2014b; Zegeye *et al.*, 2014). To test whether these stabilizing agents would affect the reactivity of the studied GRs towards Hg^{II} , experiments were performed with and without bacterial cells or phosphate, and the rate and the extent of the Hg^{II} reduction were determined. When GR2_{ab} was supplemented with *S. putrefaciens* cells ($\text{GR2}_{\text{ab+cells}}$), k_{S} did not significantly change compared to GR2_{ab} (Fig. 2, Table 1). Similarly, no significant difference was noted between GR1_{ab} and $\text{GR1}_{\text{ab+P}}$ (*i.e.* GR1 supplemented with 0.8% phosphate) (Fig. 2b, Table 1). Therefore, bacterial cells or phosphates have a minor impact on the reactivity of GRs towards Hg^{II} , and the GR1_{bio} can reduce Hg^{II} as effectively as GR1_{ab} . This suggests that *S. putrefaciens* did not inhibit significantly the reduction of Hg^{II} by GRs as it has already been shown for other bacteria (Mishra *et al.*, 2011).

4.3. Kinetics of methyl red reduction by GRs

Here also, the reaction kinetics can be described by a pseudo first-order reaction (equations 10, 11):

$$d[\text{MR}]_t/dt = -k_{\text{obs}} \times [\text{MR}]_t \quad (10)$$

$$[\text{MR}]_t = [\text{MR}]_0 \times \exp(-k_{\text{obs}} \times t) \quad (11)$$

where k_{obs} is the pseudo first-order rate constant (min^{-1}), and $[\text{MR}]_t$ and $[\text{MR}]_0$ the total concentrations of MR at time t and 0 , respectively. However, it should be noted that a certain number of kinetic curves present a latency time (slight sigmoidal shape) ($\text{GR1}_{\text{ab+Si}}$ and $\text{GR1}_{\text{ab+paa}}$, Fig. 5), and a better description of this lag-time merits to be investigated in further studies.

The values of k_{obs} were determined from equation (11) by plotting $-\ln([\text{MR}]_t/[\text{MR}]_0)$ as a function of time t (Table 2). The Fe^{II} content of GRs and the concentration of Fe^{II} sites on the green rust surface were also taken into account to normalize k_{obs} values and to define $k_{\text{Fe}^{\text{II}}}$ ($\text{L min}^{-1} \text{mmol}^{-1}$) and k_s ($\text{L mmol}^{-1} \text{min}^{-1}$), but it does not significantly affect the hierarchy as regards the kinetic rate constants (Table 2).

MR was completely reduced by GR2_{ab} in less than 30 min with a k_s value of $3.4 \text{ L mmol}^{-1} \text{min}^{-1}$ ($t_{1/2} = 5 \text{ min}$) (Fig. 5). Generally, all the GRs used showed lower k_s values than those obtained for Hg^{II} reduction (except for the Hg^{II} /GR ratios of 1/10, 1/8 and 1/0.8). For example, MR was fully reduced by GR1_{ab} in 120 min (Fig. 5) instead of only 60 min for a total reduction of Hg^{II} (Hg^{II} /GR ratio of 1/100) (Fig. 2a). However, k_s obtained for a reaction of MR with GR1_{bio} had a significantly lower value ($0.057 \text{ L mmol}^{-1} \text{min}^{-1}$) (Table 2) when compared to the reaction between GR1_{bio} (1/80) and Hg^{II} ($k_{\text{obs}} = 1.18 \text{ mmol}^{-1} \text{min}^{-1}$) (Table 1). Thus, the time needed to reduce MR by GR1_{bio} (1200 min) (Fig. 5) is 20 times higher than for the reduction of Hg^{II} (60 min) (Fig. 2a).

4.3.1. Critical role of bacterial cells and phosphate on MR reduction

Since *Shewanella* spp. were shown to degrade MR (Khalid *et al.*, 2008), the degradation of MR by GR1_{bio} could be due to an active metabolism of cells involved in the GR1_{bio} formation and remaining in the mineral suspension. To assess the effect of *S. putrefaciens* on MR removal, a series of batch experiments were performed with a fresh cell suspension in the same physiological state (late stationary phase), and with a heat inactivated suspension. In both cases, no significant decrease in MR levels was obtained indicating that cells present with GR_{bio} do not contribute to MR removal (Fig. S5b).

To test whether bacteria could affect the reactivity of GRs, *Shewanella* cells were added to GR2_{ab}, the most reactive GR (high k_s). A significant decrease in the MR reduction rate by a factor of 200 was observed relative to GR2_{ab} (Table 2, Fig. 5), which indicates that bacterial cells inhibit considerably the MR reduction by GRs. Similarly, k_s of GR1_{ab+P} was ~200 times lower than that of GR2_{ab} (0.007 L mmol⁻¹ min⁻¹ and 3.4 L mmol⁻¹ min⁻¹, respectively, Table 2), thereby underscoring that phosphates impaired significantly the reactivity of GR. This is in line with a recent work dealing with the effect of phosphates on the reduction of nitrate by GR (Etique *et al.*, 2014b).

4.3.2. A moderate inhibition of GRs reactivity towards MR by silicate and polyacrylic acid

To investigate whether other environmentally relevant components could affect GR reactivity, experiments were performed with H₄SiO₄ and paa. Such compounds can represent (or be models of) aqueous Si from quartz associated with soil particles or extra-cellular polymeric substances from biofilms (Sergent *et al.*, 2011; Jorand *et al.*, 2013). The k_s of GR1_{ab+Si} and GR1_{ab+paa} (0.156 L mmol⁻¹ min⁻¹ and 0.193 L mmol⁻¹ min⁻¹, respectively) were ~20 times higher than that of GR1_{ab+P} and ~10 times lower than that of GR2_{ab} (Table 2, Fig. 5). This result is in line with a recent study pointing out that GR1 stabilized with EPS (GR1_{ab+paa}) is quasi-unreactive towards nitrate (Zegeye *et al.*, 2014). Thus, it implies that paa,

H₄SiO₄, and cells would have a less pronounced “screen effect” than phosphate, or be sorbed onto different reactive sites as compared to MR.

As stated before, the stoichiometric difference (Fe^{II}/Fe^{III}) between GRs could influence the rate of reduction, as well as the concentration of surface Fe^{II} site of GR. However, even considering k_s or k_{FeII} , the hierarchy of GR reactivity with MR remains unchanged (Table 2):

$$k_s(\text{GR1}_{ab+p}) < k_s(\text{GR2}_{ab+cells}) < k_s(\text{GR1}_{bio}) < k_s(\text{GR1}_{ab+Si}) < k_s(\text{GR1}_{ab+paa}) < k_s(\text{GR1}_{ab}) < k_s(\text{GR2}_{ab}).$$

To know whether MR was effectively reduced and not only sorbed onto the solid phase, the presence of degradation products from MR reduction was checked. At the end of the reduction, the main absorption peak at 430 nm had disappeared while a band in the ultraviolet region at 305 nm had increased (Fig. S1). This absorbance band in the UV region could be assigned to the presence of *p*-dimethylaminoaniline. This result is consistent with equation (5), as well as with the findings of Hou *et al.* (2007) and Kone *et al.* (2009) who reported that methyl orange and methyl red were reduced by zero-valent iron and GR2_{ab}, respectively, through cleavage of the azo bond into sulfanilic or carboxyanilic acid and *p*-dimethylaminoaniline. Control with aqueous Fe^{II} was performed and no reduction of MR was observed (Fig. S5b).

4.4. MR and Hg^{II} reduction mechanisms by GRs

4.4.1. Estimation of the number of moles of contaminant that could be reduced by GRs

The number of moles of Hg^{II} species or MR that could be reduced for a given surface area of GR at the given pH value can only be determined for a run highlighting an incomplete consumption of the contaminant. Thus, the initial concentration of GR used during the reduction of the pollutant has not to be sufficient to fully reduce the initial amount of Hg^{II} species or MR. This phenomenon is described by the term saturation.

Additional experiment was carried out, where Hg^{II} species were added successively to the GR1_{bio} suspension (400 μM) in quantities of 5 μM to 100 μM to reach a final Hg^{II}

concentration of 130 μM in the reaction medium (pH 8.2) (Fig. 6). One day after the beginning of the reaction, the quantity of Hg^{II} removed was 77 μM (Fig. 6), representing 59 % of the total addition of Hg^{II} . Between each addition of Hg^{II} , the kinetics of the reaction decreased (first addition: $k_{\text{obs},\#1} = 4.1 \pm 0.1 \times 10^{-2} \text{ min}^{-1}$; third addition: $k_{\text{obs},\#3} = 1.6 \pm 0.2 \times 10^{-2} \text{ min}^{-1}$; last addition: $k_{\text{obs},\#5} = 5.0 \times 10^{-4} \text{ min}^{-1}$) suggesting that the availability of Fe^{II} sites on the surface of GR1_{bio} lowered. However, all the Fe^{II} sites on the GR1_{bio} surface do not seem to be saturated with a Hg^{II} /GR ratio of 1/5.2 (i.e. 77/400). This ratio was increased in another additional run performed with a Hg^{II} /GR ratio of 1/0.8 at pH 8.2. A plateau was reached at 1.66 μM (Fig. S4b), evidencing a partial reduction of mercury with an initial concentration of GR1_{bio} of 4 μM . Here, a saturation was highlighted and the amount of Hg^{II} species reduced by surface area concentration of GR1_{bio} was estimated at $1.1 \times 10^{-4} \text{ moles of Hg}^{\text{II}} \text{ m}^{-2}$.

For the reduction of MR, none of our carried out experiments showed an incomplete consumption of $\text{C}_{15}\text{H}_{15}\text{N}_2\text{O}_3$ species at pH 7 (Fig. 5). Nevertheless, we can estimate the minimum amount of $\text{C}_{15}\text{H}_{15}\text{N}_2\text{O}_3$ species that can be reduced by GR1_{bio} at $4.3 \times 10^{-6} \text{ moles of MR m}^{-2}$ or more.

4.4.2. Parameters influencing the kinetics of the MR reduction vs the Hg^{II} reduction

The MR reaction kinetics were more affected in the presence of GR stabilizing agents, such as phosphate or bacterial cells, than the reduction of Hg^{II} (GR2_{ab+p}/ Hg^{II} and GR2_{ab+cells}/ Hg^{II} , Fig. 2a; GR2_{ab+p}/MR and GR2_{ab+cells}/MR, Fig. 5). These stabilizing compounds have already proved their ability to prevent the formation of magnetite from GR1_{ab} or GR1_{bio} dissolution (Bocher *et al.*, 2004; Jorand *et al.*, 2013). It was suggested that reactive ferrous sites involved in GR dissolution and located on the lateral faces would be hidden by phosphate (Bocher *et al.*, 2004). For example, the reduction of nitrate by GR1 implies these lateral Fe^{II} sites, and its kinetics is strongly affected by phosphate (Etique *et al.*, 2014b), by EPS (Zegeye *et al.*, 2014), or by silicate (data not shown). In the present

investigation, since GR stabilizing agents affect MR reduction, MR would preferentially react with the same Fe^{II} sites of GR as those involved by the sorbing ligands. As Hg^{II} reduction was slightly affected by phosphate or cells, we may suppose that Hg^{II} would preferentially react with other reactive sites, which might rather be located on the basal face of the GR crystal. Other GR stabilizers such as silicate and paa, are also known to prevent unstable GRs from transforming into magnetite (Sergent *et al.*, 2011; Jorand *et al.*, 2013), however, they affect to a lesser degree the MR reduction by GRs.

In addition, the presence of sorbing ligands could slow down the reactivity of GRs by electrostatic repulsion and/or steric hindrance. Indeed, the interactions of negatively charged MR with GR surfaces covered with a layer of anionic ligands such as phosphate anions, cells, or polymers might also explain the decrease in the reduction rate of MR as compared to that of Hg.

5. Conclusion

The difference in reactivity between GRs has more to do with the nature of the contaminant (Hg^{II} or MR) than with the type of GRs (GR1 or GR2). At pH 7, the reaction of GRs with MR is generally slower than with Hg^{II} , and moreover, the reduction of MR seems to be more affected by stabilizing agents than Hg^{II} reduction. More precisely, the reduction of MR is sharply limited by phosphate and bacterial cells, whereas in the presence of silicate and paa this reduction is moderate. This difference in GR reactivity towards mercury and methyl red would probably be based on the nature of Fe^{II} surface sites involved during the reduction: lateral Fe^{II} surface sites would be monopolized for MR reduction, whereas basal Fe^{II} surface sites would interact with Hg^{II} species.

Although the Hg^{II} reduction is slightly hindered in the presence of stabilizing agents, the variation of pH suspension from 7.0 to 9.5 leads to a decrease of the kinetics rate. This finding can be explained by the fluctuation of the surface charge of GR (pH 7.0: positive, pH

8.2: zero, pH 9.5: negative) that governs the adsorption and the reduction of $\text{Hg}(\text{OH})_2$, the predominant phase of Hg^{II} species at pH of this investigation.

In environmental conditions where the fougèrite mineral can be represented by biologically synthesized hydroxycarbonate green rust (GR1_{bio}), Hg^{II} species would be more reduced than MR. Indeed, the amount of Hg^{II} species reduced by surface area of GR1_{bio} (1.1×10^{-4} moles of $\text{Hg}^{\text{II}} \text{ m}^{-2}$) is significantly higher than those of MR (4.3×10^{-6} moles of MR m^{-2}). In engineering applications using GRs as a reactive sorbent and/or reducing agent, more attention should be paid to the co-occurring ligands as regards the type of target pollutants. These results have strong implications in attenuation and remediation studies aiming to predict the fate and transport of methyl red and mercury in natural or engineered systems.

Acknowledgements

The authors acknowledge financial support from the Agence Nationale de la Recherche (France) (“Arctic Metals” project), the Ministère de l’Enseignement Supérieur et de la Recherche (France), and the Zone Atelier Moselle (Lorraine, France). We thank J. Ghanbaja for TEM analysis, I. Bihannic for XRD analysis, B. Grégoire and A. Razafitianamaharavo for BET analysis, and M. Abdelmoula for TMS analysis.

List of symbols

GR1_{ab} :	abiotically synthesized hydroxycarbonate green rust
GR2_{ab} :	abiotically synthesized hydroxysulfate green rust
GR1_{bio} :	microbially synthesized hydroxycarbonate green rust
$\text{GR1}_{\text{ab+P}}$:	abiotically synthesized hydroxycarbonate green rust with phosphate
$\text{GR1}_{\text{ab+Si}}$:	abiotically synthesized hydroxycarbonate green rust with silicic acid
$\text{GR1}_{\text{ab+paa}}$:	abiotically synthesized hydroxycarbonate green rust with polyacrylic acid

543	GR _{2ab+cells} :	abiotically synthesized hydroxysulfate green rust with heat-killed cells
544	A:	surface area concentration ($\text{m}^2 \text{L}^{-1}$)
545	k_{obs} :	pseudo first-order rate constant (min^{-1})
546	k_{FeII} :	k_{obs} normalized to the Fe^{II} content of GRs ($\text{L mmol}^{-1} \text{min}^{-1}$)
547	k_{s} :	k_{obs} normalized to the estimated Fe^{II} surface sites of GRs ($\text{L mmol}^{-1} \text{min}^{-1}$)
548	paa:	polyacrylic acid

549

550 **Appendix A. Supplementary data**

551 Supplementary data associated with this article (Tables S1 – S4; Notes S1 – S3; and Fig. S1 –
552 S6) can be downloaded online.

553

554 **References**

- 555 Bearcock, J. M., Perkins, W. T., Pearce, N. J. G., 2011. Laboratory studies using naturally
556 occurring "green rust" to aid metal mine water remediation. *Journal of Hazardous*
557 *Materials* 190, 466–473.
- 558 Benali, O., Abdelmoula, M., Refait, P., Génin, J.-M. R. 2001. Effect of orthophosphate on the
559 oxidation products of Fe(II)–Fe(III) hydroxycarbonate: the transformation of green rust to
560 ferrihydrite. *Geochimica et Cosmochimica Acta* 65, 1715–1726.
- 561 Bocher, F., Géhin, A., Ruby, C., Ghanbaja, J., Abdelmoula, M., Génin J.-M. R. 2004.
562 Coprecipitation of Fe(II–III) hydroxycarbonate green rust stabilized by phosphate
563 adsorption. *Solid State Sciences* 6 (1), 117–124.
- 564 Carlson, H. K., Clark, I. C., Melnyk, R. A., Coates, J. D. 2012. Toward a mechanistic
565 understanding of anaerobic nitrate-dependent iron oxidation: balancing electron uptake and
566 detoxification. *Frontiers in Microbiology* 3 (57), 1–6.

- Charlet, L., Bosbach, D., Peretyashko, T. 2002. Natural attenuation of TCE, As, Hg linked to the heterogeneous oxidation of Fe(II): an AFM study. *Chemical Geology* 190, 303–319.
- Christiansen, B. C., Balic-Zunic, T., Dideriksen, K., Stipp, S. L. S. 2009. Identification of green rust in groundwater. *Environmental Science & Technology* 43, 3436–3441.
- Drissi, S. H., Refait, P., Abdelmoula, M., Génin, J.-M. R. 1995. The preparation and thermodynamic properties of iron(II)-iron(III) hydroxide-carbonate (green rust); Pourbaix diagram of iron in carbonate-containing aqueous media. *Corrosion Science* 37, 2025–2041.
- Elsner M., Schwarzenbach R. P., Haderlein, S. B. 2004. Reactivity of Fe(II)-bearing minerals toward reductive transformation of organic contaminants. *Environmental Science & Technology* 38, 799–807.
- Erbs, M., Hansen, H. S. B., Olsen, C. E. 1999. Reductive dechlorination of carbon tetrachloride using iron(II) iron(III) hydroxide sulfate (green rust). *Environmental Science & Technology* 33, 307–311.
- Etique, M., Jorand, F. P. A., Zegeye, A., Grégoire, B., Despas, C., Ruby, C. 2014a. Abiotic process for Fe(II) oxidation and green rust mineralization driven by a heterotrophic nitrate reducing bacteria (*Klebsiella mobilis*). *Environmental Science & Technology* 48, 3742–3751.
- Etique, M., Zegeye, A., Grégoire, B., Carteret, C., Ruby, C. 2014b. Nitrate reduction by mixed iron(II-III) hydroxycarbonate green rust in the presence of phosphate anions: the key parameters influencing the ammonium selectivity. *Water Research* 62, 29–39.
- Fredrickson, J. K., Zachara, J. M., Kennedy, D. W., Dong, H., Onstott, T. C., Hinman, N. W., Li, S.-M. 1998. Biogenic iron mineralization accompanying the dissimilatory reduction of hydrous ferric oxide by a groundwater bacterium. *Geochimica et Cosmochimica Acta* 62, 3239–3257.

- Génin, J.-M. R., Abdelmoula, M., Ruby, C., Upadhyay, C. 2006. Speciation of iron; characterisation and structure of green rusts and $\text{Fe}^{\text{II-III}}$ oxyhydroxycarbonate fougérite. *Comptes Rendus Geosciences* 338, 402-419.
- Génin, J.-M. R., Bourrié, G., Trolard, F., Abdelmoula, M., Jaffrezic, A., Refait, P., Maître, V., Humbert, B., Herbillon, A. 1998. Thermodynamic equilibria in aqueous suspensions of synthetic and natural Fe(II)–Fe(III) green rusts: occurrences of the mineral in hydromorphic soils. *Environmental Science & Technology* 32, 1058–1068.
- Guilbaud, R., White, M. L., Poulton, S. W. 2013. Surface charge and growth of sulphate and carbonate green rust in aqueous media. *Geochimica et Cosmochimica Acta* 108, 141-153.
- Hansen, H. C. B., Guldberg, S., Erbs, M., Koch, C.-B. 2001. Kinetics of nitrate reduction by green rusts effects of interlayer anion and Fe(II):Fe(III) ratio. *Applied Clay Sciences* 18, 81–91.
- Hou, X.-L., Yang, Z., Yeung, K.-S., Wong, H. N. C., 2007. Chapter 5.3 Five-membered ring systems: furans and benzofurans. *Progress in Heterocyclic Chemistry* 18, (C), 187–217.
- Jorand, F. P. A., Sergent, A.-S., Remy, P.-P., Bihannic, I., Ghanbaja, J., Lartiges, B., Hanna, K., Zegeye, A. 2013. Contribution of anionic vs neutral polymers to the formation of green rust 1 from $\gamma\text{-FeOOH}$ bioreduction. *Geomicrobiology Journal* 30, 600–615.
- Khalid, A., Arshad, M., Crowley, D. E. 2008. Decolorization of azo dyes by *Shewanella* sp. under saline conditions. *Applied Microbiology and Biotechnology* 79, 1053-1059.
- Kone, T., Hanna, K., Abdelmoula, M., Ruby, C., Carteret, C., 2009. Reductive transformation and mineralization of an azo dye by hydroxysulphate green rust preceding oxidation using H_2O_2 at neutral pH. *Chemosphere* 75, 212–219.
- Lee, W., Batchelor, B., 2002. Abiotic reductive dechlorination of chlorinated ethylenes by iron-bearing soil minerals. 2. Green rust. *Environmental Science & Technology* 36, 5348-5354.

- 616 Loyaux-Lawniczak, S., Refait, P., Ehrhardt, J.-J., Lecomte, P., Génin, J.-M. R., 2000.
- 617 Trapping of Cr by formation of ferrihydrite during the reduction of chromate ions by
- 618 Fe(II)-Fe(III) hydroxysalt green rusts. *Environmental Science & Technology* 34, 438-443.
- 619 Mills, S. J., Christy, A. G., Génin, J.-M. R., Kameda, T., Colombo, F., 2012. Nomenclature of
- 620 the hydrotalcite supergroup: natural layered double hydroxides. *Mineralogical Magazine*
- 621 76, 1289-1336.
- 622 Mishra, B., O'Loughlin, E. J., Boyanov, M. I., Kemner, K. M., 2011. Binding of Hg^{II} to high-
- 623 affinity sites on bacteria inhibits reduction to Hg^0 by Mixed $\text{Fe}^{\text{II/III}}$ phases. *Environmental*
- 624 *Science & Technology* 45, 9597–9603.
- 625 Mitsunobu, S., Takahashi, Y., Sakai, Y., 2008. Abiotic reduction of antimony (V) by green
- 626 rust ($\text{Fe}_4(\text{II})\text{Fe}_2(\text{III})(\text{OH})_{12}\text{SO}_4 \cdot 3\text{H}_2\text{O}$). *Chemosphere* 70, 942–947.
- 627 Myneni, S. C. B., Tokunaga, T. K., Brown Jr., G. E., 1997. Abiotic Se redox chemistry in the
- 628 presence of Fe(II, III)-oxides. *Science* 278, 1106 – 1109.
- 629 O'Loughlin, E. J., Burris, D. R., 2004. Reduction of halogenated ethanes by green rust.
- 630 *Environmental Toxicology and Chemistry* 23, 41-48.
- 631 O'Loughlin, E. J., Kelly, S. D., Kemner, K. M., Csencsits R., Cook, R. E. 2003. Reduction of
- 632 Ag^{I} , Au^{III} , Cu^{II} , and Hg^{II} by $\text{Fe}^{\text{II/III}}$ hydroxysulfate green rust. *Chemosphere* 53, 437-446.
- 633 O'Loughlin, E. J., Gorski C. A., Scherer M. M., Boyanov M. J., Kemner K. M. 2010. Effects
- 634 of oxyanions, natural organic matter, and bacterial cell numbers on the bioreduction of
- 635 lepidocrocite ($\gamma\text{-FeOOH}$) and the formation of secondary mineralization products.
- 636 *Environmental Science & Technology* 44, 4570-4576.
- 637 Ona-Nguema, G., Abdelmoula, M., Jorand, F., Benali, O., Géhin, A., Block, J.-C., Génin J.-M.
- 638 R., 2002. Iron (II, III) hydroxycarbonate green rust formation and stabilization from
- 639 lepidocrocite bioreduction. *Environmental Science & Technology* 36, 16-20.

- 640 Pasakarnis, T. S., Boyanov, M. I., Kemner, K. M., Mishra, B., O'Loughlin, E. J., Parkin, G.,
641 Scherer, M. M., 2013. Influence of chloride and Fe(II) content on the reduction of Hg(II)
642 by magnetite. *Environmental Science & Technology* 47, 6987-6994.
- 643 Pourbaix, M., 1963. *Atlas d'équilibres électrochimiques*. Gauthier-Villars et Cie, Paris, pp.
644 422-427.
- 645 Refait, P., Memet, J.-B., Bon, C., Sabot, R., Génin, J.-M. R., 2003. Formation of the Fe(II)–
646 Fe(III) hydroxysulphate green rust during marine corrosion of steel. *Corrosion Science* 45,
647 833-845.
- 648 Refait, P., Abdelmoula, M., Génin, J.-M. R., Sabot, R., 2006. Green rusts in electrochemical
649 and microbially influenced corrosion of steel. *Comptes Rendus Geosciences* 338, 476-487.
- 650 Ruby, C., Géhin, A., Abdelmoula, M., Génin, J.-M. R., Jolivet, J. P., 2003. Coprecipitation of
651 Fe(II) and Fe(III) cations in sulphated aqueous medium and formation of hydroxysulphate
652 green rust. *Solid State Sciences* 5, 1055-1062.
- 653 Ruby, C., Haissa, R., Géhin, A., Abdelmoula, M., Génin, J.-M. R., 2006. Chemical stability
654 of hydroxysulphate green rust synthesized in the presence of foreign anions: carbonate,
655 phosphate and silicate. *Hyperfine Interactions* 167, 803-807.
- 656 Schwertmann, U., Fechter, H., 1994. The formation of green rust and its transformation to
657 lepidocrocite. *Clay Minerals* 29, 87-92.
- 658 Sergent, A.-S., Jorand, F., Hanna, K., 2011. Effects of Si-bearing minerals on the nature of
659 secondary iron mineral products from lepidocrocite bioreduction. *Chemical Geology* 289,
660 86–97.
- 661 Simon L., Francois M., Refait P., Renaudin G., Lelaurain M., and Génin J.-M. R., 2003.
662 Structure of the Fe(II-III) layered double hydroxysulphate green rust two from Rietveld
663 analysis. *Solid State Sciences* 5, 327–334.

- 664 Su, C, Puls, R. W., 2004. Significance of iron(II,III) hydroxycarbonate green rust in arsenic
665 remediation using zerovalent iron in laboratory column tests. *Environmental Science &*
666 *Technology* 38, 5224-5231.
- 667 Taylor, R. M., Schwertmann, U., Fechter, H., 1985. A rapid method for the formation of
668 Fe(II)-Fe(III) hydroxycarbonate. *Clay Minerals* 20, 147–151.
- 669 Viollier, E., Inglett, P. W., Hunter, K., Roychoudhury, A. N., Van Cappellen, P. 2000. The
670 ferrozine method revisited: Fe(II)/Fe(III) determination in natural waters. *Applied*
671 *Geochemistry* 15, 785-790.
- 672 Wiatrowski, H. A., Das, S., Kukkadapu, R., Ilton, E. S., Barkay, T., Yee, N. 2009. Reduction
673 of Hg(II) to Hg(0) by magnetite. *Environmental Science & Technology* 43, 5307–5313.
- 674 Wiatrowski, H. A., Ward, P. M., Barkay, T., 2006. Novel reduction of mercury(II) by
675 mercury-sensitive dissimilatory metal reducing bacteria. *Environmental Science &*
676 *Technology* 40, 6690–6696.
- 677 Williams, A., Scherer, M. M., 2001. Kinetics of Cr(VI) reduction by carbonate green rust.
678 *Environmental Science & Technology* 35, 3488–3494.
- 679 Zegeye, A., Bonneville, S., Benning, L. G., Sturm A., Fowle D. A., Jones, C. A., Canfield, D.
680 E., Ruby C., MacLean, L. C., Nomosatryo, S., Crowe, S. A., Poulton, S. W., 2012. Green
681 rust formation controls nutrient availability in a ferruginous water column. *Geology* 40,
682 599-602.
- 683 Zegeye, A., Etique, M., Carteret, C., Ruby, C., Schaaf, P., Francius, G., 2014 Origin of the
684 differential nanoscale reactivity of biologically and chemically formed green rusts crystals
685 investigated by chemical force spectroscopy. *Journal of Physical Chemistry C* 118, 5978–
686 5987.
- 687 Zegeye, A., Mustin, C., Jorand, F. 2010. Bacterial and iron oxide aggregates mediate
688 secondary iron mineral formation: green-rust vs magnetite. *Geobiology* 8, 209-22.

- 689 Zegeye, A., Ona-Nguema, G., Carteret, C., Huguet, L., Abdelmoula, M., Jorand, F., 2005.
690 Formation of hydroxysulphate green rust 2 as a single iron(II–III) mineral in microbial
691 culture. *Geomicrobiology Journal* 22, 389–399.
- 692 Zegeye, A., Ruby, C., Jorand, F., 2007. Kinetic and thermodynamic analysis during
693 dissimilatory γ -FeOOH reduction: formation of green rust 1 and magnetite.
694 *Geomicrobiology Journal* 24, 51-64.

Table 1. Kinetics rate constants (k_{obs} , min^{-1}), k_{obs} normalized to surface Fe^{II} sites (k_{s} , $\text{L mmol}^{-1} \text{min}^{-1}$) and k_{obs} normalized to structural Fe^{II} (k_{FeII} , $\text{L mmol}^{-1} \text{min}^{-1}$) of the reduction of Hg^{II} ($5 \mu\text{M}$) by chemically synthesized carbonated green rust (GR1_{ab} or $\text{GR1}_{\text{ab+P}}$ supplemented with phosphate), biogenic carbonated green rust (GR1_{bio}), chemically synthesized sulfated green rust (GR2_{ab}), or GR2_{ab} with a cell suspension ($\text{GR2}_{\text{ab+cells}}$). Each batch was conducted with 0.5 mM ($\text{Hg}^{\text{II}}/\text{GR} = 1/100$) or 0.05 mM ($\text{Hg}^{\text{II}}/\text{GR} = 1/10$) of GR in duplicates (except when specified). A = surface area concentration ($\text{m}^2 \text{L}^{-1}$). ND = not determined.

	A ($\text{m}^2 \text{L}^{-1}$)	Structural Fe^{II} (mM)	Surface Fe^{II} sites (μM)	$\text{Hg}^{\text{II}}/\text{GR}$	Rate constants	pH 7.0 $\times 10^{-2}$	pH 8.2 $\times 10^{-2}$	pH 9.5 $\times 10^{-2}$
GR1_{ab}	10 ± 2^a	2.2	83	1/100	k_{obs} k_{s}^{II} $k_{\text{Fe}}^{\text{II}}$	ND ND ND	2.5 ± 0.3 30 ± 3 1.1 ± 0.1	ND ND ND
		0.22	8.3	1/10	k_{obs} k_{s}^{II} $k_{\text{Fe}}^{\text{II}}$	ND ND ND	8.6 ± 0.1 1036 ± 12 39.1 ± 0.5	ND ND ND
$\text{GR1}_{\text{ab+P}}$	10 ± 2	1.2	83	1/100	k_{obs} k_{s}^{II} $k_{\text{Fe}}^{\text{II}}$	3.8 ± 0.2 45 ± 2 3.2 ± 0.2	3.0 ± 0.6 36 ± 7 2.5 ± 0.5	3.1 ± 0.2 37 ± 2 2.6 ± 0.2
GR1_{bio}	4.2	1.3	34.8	1/80	k_{obs} k_{s}^{II} $k_{\text{Fe}}^{\text{II}}$	4.1 ± 0.2 118 ± 6 3.1 ± 0.1	2.2 ± 0.3 63 ± 8 1.7 ± 0.2	2.7 ± 0.5 78 ± 14 2.1 ± 0.4
		0.13	3.48	1/8	k_{obs} k_{s}^{II} $k_{\text{Fe}}^{\text{II}}$	ND ND ND	3.5 ± 0.4 1006 ± 114 27 ± 3	ND ND ND
		0.013	0.348	1/0.8	k_{obs} k_{s}^{II} $k_{\text{Fe}}^{\text{II}}$	ND ND ND	1.5^b 4310 115	ND ND ND
GR2_{ab}	4.6 ± 0.7	2.0	38.2	1/100	k_{obs} k_{s}^{II} $k_{\text{Fe}}^{\text{II}}$	4.1 ± 0.2 107 ± 5 2.0 ± 0.1	1.6 ± 0.1 42 ± 2 0.80 ± 0.05	1.1^b 29 0.6
$\text{GR2}_{\text{ab+cells}}$	4.6 ± 0.7	2.0	38.2	1/100	k_{obs} k_{s}^{II} $k_{\text{Fe}}^{\text{II}}$	1.8 ± 0.4^c 47 ± 11 0.9 ± 0.2	2.1 ± 0.5 55 ± 13 1.05 ± 0.25	2.8 ± 0.9 73 ± 24 1.4 ± 0.5

^a Assuming the surface area was the same than $\text{GR1}_{\text{ab+P}}$

^b $n = 1$

^c $n = 3$ independent experiments

Table 2: Kinetics rate constants (k_{obs}), k_{obs} normalized to surface Fe^{II} (k_s , $\text{L mmol}^{-1} \text{ min}^{-1}$) and k_{obs} normalized to structural Fe^{II} ($k_{\text{Fe}}^{\text{II}}$, $\text{L mmol}^{-1} \text{ min}^{-1}$) of the reduction of methyl red (MR) by chemically synthesized carbonated green rust (GR_{1ab}), GR_{1ab} supplemented with 4 mM phosphate (GR_{1ab+p}) or with 4 mM H_4SiO_4 (GR_{1ab+Si}) or with 0.5 g L^{-1} of polyacrylic acid (GR_{1ab+paa}) or by biogenic carbonated green rust (GR_{1bio}), chemically synthesized sulfated green rust (GR_{2ab}), or GR_{2ab} supplemented with a cell suspension (GR_{2ab+cells}). All experiments were done at pH 7.0 $\pm$ 0.1 and in duplicates (except for GR_{1ab+Si}). A = surface area concentration.

A	Structural Fe^{II} (mM)	Surface Fe^{II} sites (μM)	MR/GR	$k_{\text{obs}} \times 10^{-3}$ (min^{-1})	$k_s \times 10^{-3}$ ($\text{L mmol}^{-1} \text{ min}^{-1}$)	$k_{\text{Fe}}^{\text{II}} \times 10^{-3}$ ($\text{L mmol}^{-1} \text{ min}^{-1}$)
GR _{1ab}	2.2	83	1/33	28 $\pm$ 4	337 $\pm$ 48	13 $\pm$ 2
GR _{1ab+p}	1.2	83	1/33	0.6 $\pm$ 0.1	7 $\pm$ 1	0.5 $\pm$ 0.1
GR _{1ab+Si}	1.2	83	1/33	13 ^b	156	11
GR _{1ab+paa}	1.2	83	1/33	16 ^b	193	13
GR _{1bio}	1.3	34.8	1/26	2.0 $\pm$ 0.2	57 $\pm$ 6	1.5 $\pm$ 0.1
GR _{2ab}	2.0	38.2	1/33	130 $\pm$ 50	3403 $\pm$ 1308	65 $\pm$ 25
GR _{2ab+cells}	2.0	38.2	1/33	0.6 $\pm$ 0.2	16 $\pm$ 5	0.3 $\pm$ 0.1

^a Assuming the surface area was the same than GR_{1ab+p}

^b n = 1

Fig. 1. X-ray diffractograms (Co, $\lambda = 0.178897$ nm) of the biogenic carbonated green rust 1 after microbial reduction of lepidocrocite ($\gamma\text{-FeOOH}$) by *Shewanella putrefaciens* (GR1_{bio}); chemically synthesized carbonated green rust 1 (GR1_{ab}), chemically synthesized carbonated green rust 1 stabilized by phosphate (GR1_{ab+P}) and sulfated green rust 2 (GR2_{ab}) (a); images of GR1_{bio} (b), GR2_{ab} (c), and GR1_{ab+P} (d) crystals. Inserts show the electron diffraction pattern of each GR. With GR1_{ab+P}, a few minor peaks ($d = 0.186$ nm; $d = 0.223$ nm; $d = 0.258$ nm) were detected on XRD diffractogram (a), corresponding probably to chukanovite ($\text{Fe}_2(\text{OH})_2\text{CO}_3$), but this phase was not confirmed by Mössbauer analysis.

a)

Fig 2a

b)

Fig 2b

Fig. 2. Removal of mercury over time in the presence of green rusts: 400 μM of biologically synthesized hydroxycarbonate green rust (GR1_{bio}), and 500 μM of chemically synthesized hydroxycarbonate green rust (GR1_{ab}), GR1_{ab} supplemented with phosphate (GR1_{ab+P}), hydroxysulfate green rust (GR2_{ab}) and GR2_{ab} supplemented with cells (GR2_{ab+cells}). The amount of Hg remaining in solution (Hg_t/Hg_0) is plotted as a function of time at a pH of 7.0 ± 0.1 (a), 8.2 ± 0.1 (b), and 9.5 ± 0.1 (c). The changes in pH after a 60 h incubation time period are reported in Table S7. Bars are the errors of two independent experiments.

Fig. 3. Evolution of the mercury consumption as a function of the Fe^{II} consumption for time between 0 and 60 min with Hg^{II}/GR ratios of 1/10 for GR1_{ab} (circle), 1/8 (triangle) and 1/0.8 (square) for GR1_{bio}. The slope of 0.53 fits well with the expected ratio Hg^{II}:Fe^{II}_{GR} of 1:2. Error bars were drawn for two independent assays.

Fig. 4. – Green rust particles behavior at low/high pH and preferential adsorption/reduction of Hg(OH)₂.

Fig. 5. Removal of methyl red (MR) in the presence of green rusts: 400 μM of biologically synthesized hydroxycarbonate green rust (GR1_{bio}), and 500 μM of chemically synthesized hydroxycarbonate green rust (GR1_{ab}), GR1_{ab} supplemented with phosphate ($\text{GR1}_{\text{ab+P}}$), silicate ($\text{GR1}_{\text{ab+Si}}$), polyacrylic acid ($\text{GR1}_{\text{ab+paa}}$), hydroxysulfate green rust (GR2_{ab}) and GR2_{ab} supplemented with cells ($\text{GR2}_{\text{ab+cells}}$). The MR remaining in solution (MR_t/MR_0) is plotted as a function of time. The pH was manually held at a constant value of 7.0 before each sampling by addition of HCl 1M or NaOH 1M. Bars are the errors of two independent experiments.

Fig. 6. Kinetics of the Hg^{II} reduction by biologically synthesized hydroxycarbonate green rust (GR1_{bio}) with five successive additions of Hg^{II} with an interval of one hour between each addition in the reaction medium containing an initial GR1_{bio} concentration of $400 \mu\text{M}$. The quantity of Hg^{II} added each hour is indicated by the value above the red arrow. The total amount of Hg^{II} added in the GR1_{bio} suspension is $130 \mu\text{M}$ and after 24 h, $77 \mu\text{M}$ of $\text{Hg}(\text{II})$ was reduced into Hg^0 .

1. The reactivity of green rusts (GR) towards Hg^{II} or methyl red (MR) was tested
2. Bacteria and phosphate did not affect GR reactivity with Hg^{II}
3. Bacteria and ligands (phosphate, silicate and polymers) affect GR reactivity with MR
4. Environmental substances may hinder GR reactivity as regards to the target pollutant

Supplementary information of:

**Pseudo-first-order reaction of chemically and biologically formed green
rusts with Hg^{II} and $\text{C}_{15}\text{H}_{15}\text{N}_3\text{O}_2$: effects of pH and stabilizing agents
(phosphate, silicate, polyacrylic acid, and bacterial cells)**

P.-Ph. Remy^{1,2}, M. Etique^{1,2}, A.A. Hazotte^{1,2,§}, A.-S. Sergent^{1,2}, N. Estrade^{3,4#}, C. Cloquet^{3,4},
K. Hanna^{5,6}, and F.P.A. Jorand^{1,2*}

¹Université de Lorraine, LCPME, UMR 7564, Villers-lès-Nancy, F-54601, France

²CNRS, LCPME, UMR 7564, Villers-lès-Nancy, F-54601, France

³CNRS, CRPG, UMR 7358, BP 20, Vandœuvre-lès-Nancy, F-54501, France

⁴Université de Lorraine, CRPG, UMR 7358, BP 20, Vandœuvre-lès-Nancy, F-54501, France

⁵ENSCR, CNRS, UMR 6226, CS 50837, Rennes Cedex 7, F-35708, France

⁶Université européenne de Bretagne, Rennes, F-35000, France

[§]Present address: LUNAM University, Subatech-LPGN, UMR 6457 & 6112, BP 92208, F-44322 Nantes Cedex 3, France

[#]Present address: Pacific Centre for Isotopic and Geochemical Research, EOAS, The University of British Columbia, 2207 Main Mall, Vancouver, British Columbia, V6T 1Z4, Canada.

*Corresponding author: frederic.jorand@univ-lorraine.fr; +33 (0)383 685 248

22

23 Table S1: Total mercury concentration in each compound used for experimentation with mercury test.

	Pure water	SnCl ₂	GR1 _{bio}	GR1 _{ab+P}	GR2 _{ab}	Sp	Fe ^{II} _{aq.}	Hg ^{II}	GR1 _{ab}
Hg (ppb)	0.482	0.760	0.370	1.872	0.315	0.574	0.519	104.514	ND

24

25 Table S2a: d_{hkl} (nm) parameters from XRD analysis for chemically synthesized and biogenic carbonated green rust from the present study
 26 (GR1_{ab+P}, GR1_{ab}, and GR1_{bio}) and from the literature ^a(Drissi *et al.*, 1995), ^b(Zegeye *et al.*, 2007).

27 n.d. : no data

Relative intensity	hkl	d_{hkl}	GR1 _{ab+P}	d_{hkl}	GR1 _{ab}	d_{hkl}	GR1 _{bio}	Ref ^a abiotic	Ref ^b biotic
100	003	0.759		0.758		0.759		0.751	0.751
32	006	0.377		0.379		0.378		0.373	0.375
1	101	0.274		-		0.274		0.273	n.d.
15	012	0.268		-		0.268		0.265	0.266
3	104	0.247		-		0.248		0.247	n.d.
12	015	0.235		0.235		0.235		0.233	0.233
1	107	0.209		-		0.209		0.208	n.d.
9	018	0.198		0.197		0.197		0.196	0.196
1	0012	0.189		0.189		0.188		0.188	n.d.
2	1010	0.175		-		0.174		0.173	n.d.

28

29

Table S2b: d_{hkl} (nm) parameters from XRD analysis for chemically synthesized hydroxysulfate green rust from the present study (GR2_{ab}) and from the literature ^a(Simon *et al.*, 1997), ^b(Hansen *et al.*, 1994).

n.d. : no data

Relative intensity	hkl	d_{hkl} GR2 _{ab}	Ref ^a	Ref ^b
100	001	1.091	1.09	1.10
80	002	0.547	0.545	0.549
60	003	0.365	0.362	0.366
20	004/100	0.276	0.275	0.275
30	101	0.67	0.266	0.226
30	102	0.246	0.245	n.d.
30	005/103	0.220	0.219	0.2195

38
39
40
41
42
43
44
45
46

47
48
49
50

47
48
49
50

Table S4. Values of pH for batch experiments (n=2) before (initial pH) and after 60 min of incubation (final pH) with Hg^{II} .

Batch experiment	Initial pH	final pH
SnCl_2	2.4	2.4
GR2_{ab}	7.0 ± 0.2	6.6 ± 0.1
	8.0 ± 0.1	7.2 ± 0.1
	9.5 ± 0.1	8.9 ± 0.3
$\text{GR2}_{\text{ab+cells}}$	7.0 ± 0.1	6.5 ± 0.1
GR1_{bio}	7.0 ± 0.1	7.2 ± 0.3
	8.1	8.0
	9.5 ± 0.1	9.0 ± 0.6
GR1_{ab}	8.2 ± 0.1	9.0 ± 0.1
$\text{GR1}_{\text{ab+P}}$	7.0 ± 0.1	7.6 ± 0.3
	8.0 ± 0.1	8.0 ± 0.2
	9.5 ± 0.1	8.8 ± 0.3
Fe^{II} (2mM)	3.3 ± 0.1	3.5 ± 0.1

56 Note S1. Calculation of structural Fe^{II}

57 $[\text{Fe}^{\text{total}}] = [\text{Fe}^{\text{II}}] + [\text{Fe}^{\text{III}}] = [\text{GR}] \times 6$

58 $[\text{Fe}^{\text{II}}]/[\text{Fe}^{\text{III}}] = \text{Mössbauer ratio}$

59 *e.g.* $\text{GR1}_{\text{ab+P}}$

60 $[\text{Fe}^{\text{II}}] = 1.2 \times [\text{Fe}^{\text{III}}]$

61 $1.2 \times [\text{Fe}^{\text{III}}] + [\text{Fe}^{\text{III}}] = [\text{Fe}^{\text{total}}]$

62 $[\text{Fe}^{\text{III}}] = [\text{Fe}^{\text{total}}]/2.2$

63 $[\text{Fe}^{\text{II}}] = [\text{Fe}^{\text{total}}] - [\text{Fe}^{\text{total}}]/2.2 = 1.63 \text{ mM}$

64 $[\text{Fe}^{\text{II}}]\text{GR1}_{\text{ab+P}} = 1.63 \times 10^{-3} \text{ M}$

65 $[\text{Fe}^{\text{II}}]\text{GR1}_{\text{bio}} = 1.5 \times 10^{-3} \text{ M}$

66 $[\text{Fe}^{\text{II}}]\text{GR2}_{\text{ab}} = 2 \times 10^{-3} \text{ M}$

67 $[\text{Fe}^{\text{II}}]\text{GR1}_{\text{ab}} = 2 \times 10^{-3} \text{ M}$

68

69 Note S2. Calculation of $k_{\text{FeII}} (\text{L mol}^{-1} \text{ min}^{-1}) = k_{\text{obs}}$ normalized to the structural Fe^{II}
70 concentration.

71 $k_{\text{FeII}} = k_{\text{obs}} / [\text{Fe}^{\text{II}}]\text{GR}$

72 Where

73 $[\text{Fe}^{\text{II}}]\text{GR}$ is the structural Fe^{II} concentration (mol L^{-1})

74

75 Note S3. Calculation of $k_s (\text{L mmol}^{-1} \text{ min}^{-1}) = k_{\text{obs}}$ normalized to the surface Fe^{II} sites (μM)
76 (see manuscript)

77 $k_s = k_{\text{obs}}/(\text{surface } \text{Fe}^{\text{II}} \text{ sites})$

78

79

80

Figure S1: UV-Vis spectra of initial MR solution (+) and the same solution after 300 min of incubation with 500 μM GR2_{ab} (x).

Figure S2. X-ray diffractograms of biologically synthesized hydroxycarbonate green rust (GR1_{bio}) before addition of Hg^{II} at the concentration of 5 μ M (green line, $t = 0$ h), and 24 h after (black line, $t = 24$ h). The initial concentration of GR1_{bio} was 400 μ M. The lattice planes of green rust (GR) are written in brackets. The intensity is expressed in arbitrary unit (a. u.).

Figure S3: Mössbauer spectra of a) GR1_{bio}, b) GR1_{ab+P}, and c) GR2_{ab}.

b)

Fig. S3 b) GR1_{ab+P}

98 c)

100 Figure S3c GR2_{ab}

Figure S4. Removal of mercury and oxidation of ferrous iron content of chemically synthesized green rust (GR1_{ab}, a) and biologically formed green rust (GR1_{bio}, b) over the time at pH 8.2. The initial concentration of Hg^{II} species was 5 μ M and the initial concentrations of green rusts used were 4 μ M, 40 μ M, 400 μ M for GR1_{bio}, and 50 μ M, 500 μ M for GR1_{ab}. The Hg^{II}/GR ratios were 1/0.8, 1/8, 1/80 and 1/10, 1/100 respectively. Error bars were drawn for two independent assays.

112

a)

b)

Figure S5. (a): Disappearance of mercury over time in the presence of 50 μM Sn^{II} (pH 3.0 ± 0.5), and 0.2 mM aqueous Fe^{II} (pH 7.0 ± 0.1) or 6×10^5 cells mL^{-1} of *Shewanella putrefaciens* (pH 7.0 ± 0.1) (cells), a cell density that is equivalent to that used for the synthesis of GR_{bio} and that is expected to remain with $GR1_{bio}$. The changes in pH after a 60 h incubation time period are reported in Table S7. Bars are the errors of two independent experiments. The stannic solution ($SnCl_2$, 50 μM pH = 3) was prepared from a stock solution (22.5 mM in HCl 1.5 M) ($SnCl_2$ salt, 98% anhydrous, 196981000, Acros). It was used as an indicator for Hg^{II} reduction, since Sn^{II} is known to be a powerful reducer (Zheng and Hintelmann, 2010). In order to avoid Sn^{II} precipitation, which can occur along with increasing pH, Hg^{II} reduction by aqueous Sn^{II} were only performed at pH 3.

(b): Disappearance of methyl red (MR) over time in the presence of 0.2 mM aqueous Fe^{II} (pH 7.0), 6×10^5 cells mL^{-1} of a fresh *S. putrefaciens* suspension (cells) or heat inactivated cells (dead Sp cells), or a solution of MR in pure water.

Figure S6. X-ray diffractograms of biologically synthesized hydroxycarbonate green rust (GR1_{bio}) before addition of Hg^{II} at the concentration of 5 μ M (green line, $t = 0$ h), and 24 h after (black line, $t = 24$ h). The initial concentration of GR1_{bio} was 4 μ M. The green rust (GR) was oxidized into magnetite (M) during the reduction of mercury. The lattice planes are written in brackets. The intensity is expressed in arbitrary unit (a. u.).

135

136 References for supplementary data:

- 137 Bocher, F., Géhin, A., Ruby, C., Ghanbaja, J., Abdelmoula, M., Génin J.-M.R. 2004.
138 Coprecipitation of Fe(II–III) hydroxycarbonate green rust stabilized by phosphate
139 adsorption. *Solid State Sciences*, 6 (1), 117–124.
- 140 Drissi, SH, Refait, P, Abdelmoula, M, Génin, J.-M.R. 1995. The preparation and
141 thermodynamic properties of iron(II)-iron(III) hydroxide-carbonate (green rust); Pourbaix
142 diagram of iron in carbonate-containing aqueous media. *Corrosion Science* 37, 2025–
143 2041.
- 144 Hansen H. C. B., Borgaard O. K., Sorensen J. 1994. Evaluation of the free energy of
145 formation of Fe(II)-Fe(III) hydroxide-sulphate (green rust) and its reduction of nitrite.
146 *Geochimica et Cosmochimica Acta* 58, 2599-2068.
- 147 Ruby, C., Haissa, R., Géhin, A., Abdelmoula, M., Génin, J.-M.R., 2006. Chemical stability of
148 hydroxysulphate green rust synthesized in the presence of foreign anions: carbonate,
149 phosphate and silicate. *Hyperfine Interactions*, 167: 803-807.
- 150 Simon L., Francois M., Refait P., Renaudin G., Lelaurain M., and Génin J.-M. R., 2003.
151 Structure of the Fe(II-III) layered double hydroxysulphate green rust two from Rietveld
152 analysis. *Solid State Sciences* 5, 327–334.
- 153 Zegeye, A., Ruby, C., Jorand, F., 2007. Kinetic and thermodynamic analysis during
154 dissimilatory γ -FeOOH reduction: formation of green rust 1 and magnetite.
155 *Geomicrobiology Journal* 24, 51-64.
- 156 Zheng, W., Hintelmann, H., 2010. Nuclear field shift effect in isotope fractionation of
157 mercury during abiotic reduction in the absence of light. *The Journal of Physical*
158 *Chemistry A* 114 (12), 4238-4245.