A year (2014-2015) of plants in Proteomics journal. Progress in wet and dry methodologies, moving from protein catalogs, and the view of classic plant biochemists

Rosa Sanchez-Lucas, Angela Mehta, Luis Valledor, Francisco Cabello Hurtado, M. Cristina Romero-Rodriguez, Lyudmila Simova Stoilova, Sekvan Demir, Luis E. Rodriguez De Francisco, Ana M. Maldonado Alconada, Ana L. Jorin-Prieto, et al.

► To cite this version:

Rosa Sanchez-Lucas, Angela Mehta, Luis Valledor, Francisco Cabello Hurtado, M. Cristina Romero-Rodriguez, et al.. A year (2014-2015) of plants in Proteomics journal. Progress in wet and dry methodologies, moving from protein catalogs, and the view of classic plant biochemists. Proteomics, 2016, 16 (5), pp.866-876. 10.1002/pmic.201500351 . hal-01239816

HAL Id: hal-01239816

<https://hal-univ-rennes1.archives-ouvertes.fr/hal-01239816>

Submitted on 24 Mar 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A year (2014-2015) of plants in *Proteomics* journal. Progress in wet and dry methodologies, moving from protein catalogs, and the view of classic plant biochemists

Rosa Sanchez-Lucas¹, Angela Mehta², Luis Valledor³, Francisco Cabello Hurtado⁴, M. Cristina Romero-Rodriguez⁵, Lyudmila Simova Stoilova⁶, Sekvan Demir¹, Luis E. Rodriguez de Francisco^{1,7}, Ana M Maldonado Alconada¹, Ana L Jorrin-Prieto¹, Jesus V. Jorrín-Novo^{1}*

¹ Agroforestry and Plant Biochemistry and Proteomics Research Group, Dpt. Biochemistry and Molecular Biology, University of Cordoba-CeiA3, Cordoba, Spain.

² Embrapa Recursos Genéticos e Biotecnologia (CENARGEN), Brasília, DF, Brazil.

³ Dpt. Biology of Organisms and Systems (BOS), University of Oviedo, Oviedo, Spain.

⁴ Mechanisms and Origin of Biodiversity Team, UMR 6553-Ecobio, University of Rennes, Rennes, France.

⁵ Centro Multidisciplinario de Investigaciones Tecnológicas, and Dpto. de Fitoquímica, Facultad de Ciencias Químicas, Universidad Nacional de Asunción, San Lorenzo-Paraguay.

⁶ Plant Molecular Biology Department, Institute of Plant Physiology and Genetics, Bulgarian Academy of Sciences, Sofia, Bulgaria.

⁷ INTEC-Sto. Domingo, República Dominicana.

(*) Corresponding author

Author for correspondence. Jesus V. Jorrín Novo.

Bioquímica y Proteómica Vegetal y Agroforestal; Dpto. Bioquímica y Biología Molecular, Universidad de Córdoba.

Campus de Rabanales, Ed. Severo Ochoa, 14071 Córdoba, España

E-mail: bfljonoj@uco.es; Phone +34 957 218574; FAX +34 957 218439

Key words: plant proteomics, proteogenomics, systems biology, signaling, signal transduction

Total number of words (including references as well as figure and table legends): 8164

Abstract

The present review is an update of the previous one published in *Proteomics 2015 Reviews* special issue [1] covering the 2014-July 2015 period. It has been written on the bases of the publications that appeared in *Proteomics* journal during that period and the most relevant ones that have been published in other high-impact journals. Methodological advances and the contribution of the field to the knowledge of plant biology processes and its translation to agroforestry and environmental sectors will be discussed. This review has been organized in four blocks, with a starting general introduction (literature survey) followed by sections focusing on the methodology (*in vitro*, *in vivo*, wet and dry), proteomics integration with other approaches (systems biology and proteogenomics), biological information, and knowledge (cell communication, receptors and signaling), ending with a brief mention of some other biological and translational topics to which proteomics has made some contribution.

INTRODUCTION

The topic of plant proteomics was reviewed in 2015 on the basis of the articles published in *Proteomics* journal (from now just *Proteomics*, written in italics) between 2001, when the first issue appeared, and May 2014 [1]. The present review is an update covering the January 2014-July 2015 period in which methodological advances and the contribution of the field to the plant biology knowledge and its translation to agroforestry and environmental sectors will be discussed. It is not intended to be repetitive considering recent reviews but aiming to get to the reader for a deeper discussion of key topics. It was written to be included in the 2016 *Proteomics Review Special Issue*, since it is the last one of the series edited by Prof. Dunn, the editor in chief of *Proteomics* since it appeared and who recently retired. It is a matter of justice to recognize the enormous effort of Prof. Dunn in establishing and consolidating *Proteomics* as a leader journal as well as his scientific contribution to the field. The corresponding author, in particular, and the plant community, in general, should acknowledge the personal consideration and the support he always gave to us reflected, just as an example, in his input to the International Plant Proteomics Organization (INPPO; <http://www.inppo.com/>) society [2], even though plant studies are not within his field of investigation. He, better than nobody else, has been an actor, director, and observer of the evolution of Proteomic Science [3].

According to PubMed, during the period of 2014-July 2015 *Proteomics* published 600 papers distributed in 40 issues, either regular or special including *Reviews 2014 and 2015 Tissue Proteomics and Imaging Mass Spectrometry*, *Top-Down Proteomics*, *Focus on Biomedical Research Trends*, *Proteomics in Chromatin Biology and Epigenetics*, *Proteogenomics*, *Signal Transduction*, *Proteomics Data Visualization*, *Plant-based Foods: Seed, Nutrition and Human Health*, *Virology meets*

79 *Proteomics, Views on Protein N-terminal Biology, and Protein Structure*. These thematic issues
80 perfectly represented the most recent technical advances, the frontier of knowledge in biology, and the
81 interest in translational biomedicine. Of 592 papers published in *Proteomics*, 133 were selected for
82 preparing this review as they dealt with plants (33 original papers, 16 reviews, and 1 meeting report,
83 representing around 10 % of the total papers released by the journal) or presented the state of the art
84 of the field from a biological, methodological or translational point of view.

85 After a PubMed search the terms proteomics or proteome in the title/abstract returned 8553
86 hits. Compared with the other -omics approaches, this figure was slightly or much higher than
87 transcriptomics/transcriptome (7728) or metabolomics/metabolome (3064) searches, respectively.
88 While looking for the combined use of the different omics approaches, PubMed provided 52
89 proteomics, transcriptomics, and metabolomics reports, as well as 113 proteomics and
90 transcriptomics, and 56 proteomics and metabolomics papers. The integration of the different -omics
91 approaches will be discussed in a further section.

92 In the surveyed period, plants continued to be underrepresented in the current proteomics
93 literature, but significantly increasing, with 683 references in PubMed (almost 15% of the total) while
94 for humans, just as a comparative example, the figure was 2795, with quite a few related to the study
95 of cancer, as well as vascular and other diseases [4]. Despite the claimed general worry about food
96 supply, environmental concerns, and climate change, these issues have not been properly reflected in
97 research funding policy in most of the western countries and even less in developing nations. This
98 issue is very well discussed by Donald Kennedy (President Emeritus at Stanford University, and a
99 former editor-in-chief of Science) in Science's editorial "Building Agricultural Research" [5]. Human
100 welfare depends on well-connected health and food supply [6-8]. Famine is the major cause of
101 mortality on a worldwide basis, and increasing nutritional value of primary crops would drastically
102 reduce health problems in the less developed countries [9]. The situation seems to be changing, but
103 very slowly; at HUPO (<https://www.hupo.org/>) or general proteomics meetings such as European
104 EuPA (<http://www.eupa.org/>) or Spanish SEProt (<http://www2.cbm.uam.es/~seprot/>), room for non-
105 human proteomics and groups interested in different type of organisms (microorganisms, plant, and

animals) has been recently created, or is under discussion. The vision or sensations at the plant specific organization INPPO is that we are surviving within the field as a scientific ghetto. With no doubt, plant and human research can and should benefit from each other. Plants provide an almost unlimited experimental system and despite being far in evolution they share an important number of genes [10], with similar mechanisms of, for example, immunity [11], and host-pathogen interactions [12]. In addition, plants may be sources of drugs for disease treatment [13] and also of allergens [14, 15]. Some more examples of common phenomena between plant and animals are totipotency, the property of the cells to be developmentally reprogrammed, the conceptual basis of the stem cell projects [16], and the involvement of telomeres and telomerases in senescence and aging [17].

The topic of plant proteomics continues to be extensively reviewed, reflecting its expansion, relevance, and expectations; although we should assume by now that there is a long distance between expectations and realities, and that the full potential of proteomics is far, once again if compared with humans, from being fully exploited in plant biology research and crop science [18-20]. Here, in *Proteomics*, and also in other journals, much attention has been given to proteomics of plant stresses responses, the major topic in plant biology because of its implication in crop protection and biotechnological improvement [21-23]. Up to 15 plant proteomics reviews have been recently published in *Proteomics*, most of them in the Vol. 15, Iss. 5-6 (Reviews 2015) and 10 (Plant-based Foods: Seed, Nutrition and Human Health). Strikingly, the thematic issues covering the most recent technical novelties or the frontier in proteomics and biology knowledge have none or at most two plant papers per issue; it was the case of *Tissue Proteomics and Imaging Mass Spectrometry* (Vol.14, Iss. 7-8, one plant reference by Maciel et al., 2014 [24]), and *OMICS Views on Protein N-terminal Biology* (Vol. 15, Iss. 4; two plant original papers by Zhang et al., 2015 [25] and Venne et al., 2015 [26]). The review by Walton et al. (2015) [27] on signaling, a topic later discussed, deserves special attention, and is one of the plant reviews we should not miss. Regarding plant reviews in other journals, at least 15 have appeared, with some of them covering topics that will be mentioned later, including software, database and repositories [28] PTMs [29, 30], and signaling [31].

Accepted Article

132 Recently, Jorin-Novo (2015) [32] published an opinion article in which scientific standards
133 and MIAPE were discussed in the context of proteomics work and publications. From his experience,
134 the failure to meet proper scientific standards and those requested for a proteomics paper is the most
135 obvious and immediate reason for rejection. As just a simple example, the general terminology
136 employed in the literature included terms such as protein expression, when proteomics approaches
137 actually determine abundance [1, 32]. Abundance is the result not only of gene expression
138 (transcription and translation) but also of protein degradation and PTMs.

139 The list of referenced papers did not pretend to be exhaustive, but limited, according to the
140 author's considerations, to those that presented some methodological, biological or translational
141 novelties. Some of the revised papers fit in the general group of comparative analysis based on 2-DE
142 coupled to MS analysis and they were not included. On the other hand, special attention was given to
143 those using second to fourth generation techniques (labelling, gel-free, SRM/MRM [1]), those
144 integrating classic and -omics approaches, or focused on PTMs and interactomics. Regarding the
145 previous period reviewed, there have not been real novelties in the experimental (plant) system used,
146 the biological process studied or the translational purpose pursued [1]. There is still a general need to
147 go beyond the construction of a protein catalog accompanied by more speculation than real
148 discussion, although better annotations and deeper knowledge is required in some cases.

150 WET AND DRY METHODOLOGIES

151 *In vitro* and *in situ* proteomics analysis

152 Proteomics is a discipline in expansion with continuous novelties and improvements in
153 techniques, protocols, softwares and equipments [33, 34]. This is also true for quite standard and well
154 established protocols in the early proteomics workflow stages prior to MS analysis, as it is important to
155 optimize them to each experimental system (plant, organ, tissue, cell or subcellular fraction). Thus,
156 improved protocols dealing with protein extraction [35], depletion [36, 37], fractionation [38], gel

staining [39], multiplexing [40], trypsin protein digestion [41] and sample preparation for shotgun analysis [42, 43] have recently been published.

The use of different MS platforms in plant proteomics analysis was previously discussed [1], however it is important to mention that new ion fragmentation procedures as alternative to the current collision induced dissociation, CID, and electron transfer dissociation, ETD, such as photodissociation, have been claimed to produce enhanced sequence coverage and higher confidence in sequence assignment [44, 45].

One of the limitations of proteomics, when compared to genomic and transcriptomic analyses, is that proteins cannot be amplified as can DNA and therefore, optimized and efficient extraction methods must be developed. Thus, the very laborious and time consuming used protocols and the number of replicates (the three biological replicates performed as a rule are not enough to perform solid statistical analysis and get confident biological conclusions [46], it is recommended to perform at least 5 biological replicates to allow a reliable estimation of residual variance when more than two treatments are compared) are the main bottleneck that makes proteomics unpopular among plant biologists. The use of an appropriate extraction method is even more relevant for PTMs or interactomics studies that request fractionation steps. These issues have been addressed by Krahmer et al (2015) [47], who compared different alternatives for sample preparation compatible with large numbers of samples in phosphoproteomics analysis.

Protein extraction is still challenging and constitutes a big difficulty in the analysis of specific proteins such as integral membrane proteins [48]. Integral membrane proteomics analyses involve complicated protocols that include membrane purification and protein extraction. In order to analyze the symbiosome integral membrane proteins, Clarke et al. (2015) [49] reported a protocol that included isolation of symbiosomes with three-step Percoll gradients and preparation of the membrane system by ultracentrifugation of the broken symbiosomes supernatant, followed by bicarbonate stripping and chloroform-methanol protein extraction prior to MS analysis. Although the excessive manipulation of the sample may interfere in the integral membrane protein analysis it has a great

value from a descriptive point of view. Once the protein is identified, the expression of the corresponding genes and the associated phenotype can be assessed by using complementary approaches. As an example, in order to understand intracellular protein trafficking (a hot topic in plant biochemistry), Heard et al. (2014) [50] enriched partially overlapping subsamples of endosomal proteomes associated with endomembrane from *A. thaliana* seedlings and used affinity purification of fluorescent fusion constructs for seven subcellular marker proteins in order to predict its precise or shared location, a strategy that resembles LOPIT [51]

From the current literature it is clear that plant proteomics is moving from 2-DE/MS (still the bases of an important number of works published) to shotgun (bottom up gel free label free). Shotgun will be the dominant platform in the near future if accompanied by a powerful and potent MS analysis and equipment. It is simple and does not require much manipulation of the sample. The iTRAQ, quite popular among the Chinese groups, appeared as the third alternative, even though plant proteomicists did not pay much attention in the past to labelling techniques [52-55]. By using this technique, Pan et al. (2015) [56] have reported up to 13,706 and 12,124 proteins identified with a LTQ-Orbitrap hybrid mass spectrometer in *Arabidopsis thaliana* roots and leaves. Changes in abundance between plants supplied with different iron concentrations were observed for 886 proteins with only a low percentage of them showing good correlation with transcript levels.

Whatever the employed platform, it is very important that it is validated, as a preliminary step, for the experimental system being analyzed, as for any analytical tool, in order to allow the determination of the number of proteins confidently identified and quantified. Serial dilution of the sample could be enough for validation purposes for those approaches based on label-free [57].

Fourth generation *in vitro* proteomics, SRM, MRM, or mass-western, sustain hypothesis driven protein analysis. In the surveyed period just one paper, by Rogniaux et al. (2015) [58] was published in *Proteomics* reporting the use of targeted approach for determining the relative abundance of allergens in several wheat varieties, and showing the potential of the technique for translational, and food traceability, purposes. Although the use of these techniques in plants is still limited, targeted

proteomics is the way to go for validation of the candidate proteins identified in 2-DE or shotgun analyses. It is certainly a more appropriate follow-up study when compared to qRT-PCR analysis frequently reported in proteomics publications.

Imaging has opened new possibilities for *in vivo* or *in situ* proteomics analysis and integration with cellular biology techniques [59], giving, at the same time, the possibility of proteomics analysis at the cellular level [60]. It is a more or less well established technique for metabolomics analysis [61] but under development for proteins [62] and plant tissue samples. The technique has been the focus of the *Proteomics Tissue Proteomics and Imaging Mass Spectrometry* special issue (Vol. 14, Iss. 7-8), in which, two plant studies were reported. Maciel et al. (2014) [24] used the laser ablation imaging technique to evaluate the distribution of selenium and sulfur in leaves of transgenic and non-transgenic *A. thaliana*. One of the limitations of imaging studies is the processing of high amounts of datasets, and therefore attention has been given to data processing methods, as discussed by Wijetunge et al. [63] and Franceschi and Wehrens (2014) [64].

***In silico* analysis and data interpretation: software, databases and repositories**

Proteomics, as any -omics approach, generates from hundreds to thousands of identified products, being impossible to analyze and interpret all the data in a classic, mind based, manner [65]. The high amount of data generated and the costs of the analysis are the main reasons for the low number of samples and biological replicates analyzed. Protein profiling requires software and algorithms for protein identification, quantitation, functional and structural characterization, location, interaction, networks, integration, visualization and deposition in databases and repositories. These softwares are continuously created, improved and updated [66-69]. Recently, Sakata and Komatsu (2014) [28] have surveyed and reviewed plant proteomics databases and websites as well as recent technologies related to data collection and annotation.

The use of most algorithms and softwares is far from the current capacity of most plant biochemists working with proteomics. This is the reason why the dry part of the work is in the hands

of mass spectrometrists or bioinformaticians usually with no expertise in plant biology. This results in, more often than would be advisable, a blind acceptance of the computer provided data without a minimum rationale and critical evaluation of the data confidence, making it difficult to conclude from a biological point of view. The quite dynamic character of the proteomes and the fixed film provided by our experiments makes most interpretations speculative. The situation for a standard plant biochemist is even worse considering the number of algorithms that is continuously appearing [70]. Which one, what for, how to use them and translate to a biological discussion could be hell for most of us [71, 72]. In this regard, *Proteomics* a special issue, *Proteomics Data Visualization* (17, 10). In this and other regular issues a number of reviews discussing these questions have appeared [73-78]

The integration of disseminated data through databases is a pursued objective that will benefit the scientific community. In this direction, The BioMart Community Portal (www.biomart.org) is a good example the plant community should follow [79]. It includes over 800 different biological datasets spanning genomics, proteomics, model organisms, cancer data, ontology information and more.

PROTEOMICS INTEGRATION WITH OTHER -OMICS

Within the -omics approaches, the advantage, if so, of proteomics, and also metabolomics, is that the information provided is closer to the phenotype [80]. On the other hand, it is much more complicate at the methodological level (i.e. genomics and transcriptomics are favored by PCR and well established microarray techniques). At the molecular level, the phenotype is the result of gene expression through transcriptional and translational events and gene products and metabolites interaction, so proteins are just part of the story. As previously discussed [1], proteomics has limitations, the more relevant being proteome coverage, quantitation, identification in orphan organisms, PTM, and interactomics. Protein identification and quantification cannot always be as confident as we wish because of the organism, experimental design, and the employed techniques and

261 protocols, making our data interpretation conservative or speculative. In comparative proteomics, the
262 high variability in the protein profile among biological samples and the low number of biological
263 replicates employed in most studies hinder the biological interpretation. Thus, it is not easy to establish
264 protein abundance ranges in normal (control) samples, as it has been shown by Higdon et al. (2015)
265 [81]. These authors used data from two human proteome studies featured in Nature, and found that
266 protein abundance among biological replicates varied by ± 4 - to 10-fold for most proteins, with
267 coefficient of variation (CV) in between 62 and 117 %. To control or reduce this CV, it is
268 recommended to compare proteins identified with at least 3 or more unique peptides and normalize
269 abundances against housekeeping proteins with low variability.

270 Just by itself, and more than generating a list or catalog of protein species, proteomics can
271 provide real biological knowledge; however it would require, first, the use of a model systems, in
272 plants mostly *A. thaliana*, with a good collection of mutants for reverse and functional genetics
273 studies and, second, an appropriate experimental system, as for example cell suspensions. A good
274 example of this is the paper published by Smith et al. (2015) [82] in which signal transduction
275 pathways mediating plant programmed cell death was studied. By using isobaric-tagged relative and
276 absolute quantitation of proteins present in the extracellular matrix, 33 proteins were identified as
277 putative cell-death regulatory proteins. One of these proteins, the CYCLASE 1, which had previously
278 no known function, was a negative regulator of cell death in *Arabidopsis*.

279 The integration of the different -omics, in the direction of systems biology, is far from being a
280 reality or was only possible for a reduced number of research groups or through large international
281 consortia [83]. Systems biology, wide genomic approaches, and the recently coined proteogenomics,
282 in combination with image phenotyping, classical molecular, physiological and cell biology
283 approaches [84, 85] will get us to a deeper and better rather than to a partial and biased knowledge of
284 the biological systems. This knowledge is highly relevant for crop improvement aiming at
285 productivity yield increases and survival under extreme environmental conditions through classical,
286 molecular marker assisted, and biotechnological breeding [11, 86-88]

Combined proteomics and transcriptomics analysis has been used in the study of the maize response to waterlogging and resulted, in general, in good correlations between both data [54]. But this is not necessarily the rule and it depends on the gene or gene groups, being the consequence of transcriptional, posttranscriptional, translational and posttranslational events. Abundance of both molecules depends not only on their synthesis but also on their stability and degradation. It is also known that different mRNAs possess different translational efficacy [89].

At least one paper, by Gupta et al. (2015) [90], has been published in which integrated proteomics and metabolomics have been employed as experimental approaches. They analyzed differences between soybean lines with different color in the seed coat. Differences in flavonoids, which provide the molecular basis of the color, were correlated with levels in pathway-related enzymes. Finally, there are not many papers in which the combined use of the three -omics approaches are reported. In the surveyed period, two papers dealt with the detection of unintended effects in transgenic maize [91, 92].

In any case, single and easy multi-omics approach experiments can be always performed and the preliminary steps of the workflow should be evaluated as one of the limitations is just preparative. In order to make confident data comparison or validation, these preparations should come from the same sample. Independent experiments or extractions can originate artefacts or biases, mostly coming from the inherent biological variability. In this direction, recent workflows aimed to extract biomolecules (DNA, RNAs, proteins, and metabolites) have been developed [93]. Despite dedicated workflows showed their efficiency, combined protocols demonstrated their performance in isolating biomolecules suitable to be processed by high-throughput pipelines (gel free label free proteomics, high-throughput metabolomics analyses and next generation sequencing transcriptomic analyses of messenger or regulatory RNAs) while avoiding biases related to the use of not exactly the same sample for the different -omics level analyses. The full overlapping of the datasets makes it easier to interpret and analyze the data when performing a Systems Biology-based integrative approach.

Proteogenomics is a relatively new and emerging field at the intersection of genomics and proteomics, being part of the wide genomics approaches. Although it may have multiple derivations (RNA and protein correlation studies, proteomics and data validation, gene sequence analysis, among others), in the simplest way it can be understood as the use of proteomics data and protein MS derived sequences to improve gene sequence annotations [94-96]. From the proteomics perspective it has an important feedback as correct sequence annotation will increase the number and confidence of the protein identification [97], favoring, at the same time the studies of non-model organisms [98]. It will also go deeper into the knowledge of posttranscriptional and translational events giving rise to the different gene products, protein species or proteoforms [99].

The topic has been deeply reviewed in *Proteomics* (*Proteogenomics* special issue; Vol. 14, Iss. 23-24). Proteogenomic experiments and prediction tools will differ between genome-sequenced and –unsequenced, as well as eukaryotic or prokaryotic organisms [95, 100].

FROM PROTEOMICS TO BIOLOGY

Cell communication, receptors, and signaling pathways, are good examples of how proteomics can contribute to biological knowledge and be used for translational purposes. This topic has been well presented in a recent, excellent review, by Walton et al. (2015) [27]. Signaling is a hot topic of great relevance in biology mainly approached by using functional and reverse genetics. It is quite challenging as it deals with minor and recalcitrant, membrane bound, proteins, PTMs and interactions, whose investigation requires integration with classic, transcriptomics and metabolomics approaches. In the recent literature surveyed in the reviewed period, this topic is well covered with both review papers and original contributions. Some other topics are also of interest but are somehow related to this or are less represented in the most recent literature. This is the case of plant developmental processes, plant-pathogen interactions and responses to other biotic stresses [101-104].

Plants chemically communicate and interact with their biotic and abiotic surroundings, responding and adapting its biology to it. They also complete their developmental program (from seed

germination to seed production through vegetative and reproductive phases) in coordination with seasonal changes noticed mainly as variations in light, temperature and water availability. There are also daily changes associated to a circadian clock timed by, among other factors, light regimes. The integration of environmental responses with development is crucial for plant survival, growth, and from an agronomic point of view, productivity. Plants, as sessile organisms, are very versatile, and their genome expression and metabolism is continuously reprogrammed in response to physico-chemical and biological stimuli. Despite the high number of specialized organs, tissues and cell types within a plant, all of them respond coordinately, implicating in good intercellular connexion systems with chemical messages between receptor and effector cells. At the individual cell level, gene expression changes are mediated by receptors, intracellular and interorganule signals, and signal transduction pathways resulting in changes in gene expression through transcriptional, translational and posttranslational events. Quite a number of molecules are known that act as external or internal chemical signals, including phytohormones, and those from microorganisms (symbionts, parasites or pathogens) [101, 103-106]. On the other side, receptors and signal transduction pathways and their components are starting to be characterized [31, 107].

Proteomics is contributing to the above processes through descriptive, comparative, PTMs, and interaction studies. Walton et al. (2015) [26] updated the contribution in these areas and the different strategies employed in dissecting plant hormone signaling. Some following papers dealing with PTMs and interactomics related to cell signaling are highlighted. Others, using descriptive or comparative proteomics approaches have been skipped from this review.

Protein phosphorylation is the most common mechanism of signal and because of that phosphoproteomics is the immediate strategy to be employed. Silva-Sanchez et al., (2015) [108], and Li et al., (2015) [29] have summarized the current methodologies used in extracting, enriching, and fractionating phosphoproteins, and in their MS analyses for identification, quantitation, and phosphorylation site mapping, conclude on cross-talk between different PTMs (phosphorylation, glycosylation, and redox modifications) and data validation. They provided a list of phosphoproteins,

softwares and databases. As an alternative to the current protocols, Han et al. (2015) [109] have proposed a faster workflow analysis in the case of sample amount limitation that increases coverage; it included integrated multistep enzymatic digestion, phosphopeptide enrichment and database search (iMEED). In an attempt to elucidate post-translational processes mediating circadian events, Choudhary et al. (2015) [110] performed a phosphoprotein analysis employing TiO₂ phosphopeptide and MS analysis. They identified 1586 phosphopeptides on 1080 protein groups, out of which 102 changed in abundance, including clock proteins (ELF4, PRR3), transcription factors and kinases. By using isotope-assisted quantitative phosphoproteomics, Minkoff et al. (2015) [111] carried out a comparative phosphoproteomics study between wild and mutant ABA-insensitive *A. thaliana* plants. Multiple changes occurred as early as 5 minutes after ABA treatment, with a reduction of the phosphorylation events in the mutant.

Reactive oxygen species overproduction is a typical plant response to oxidative stress, these species acting as signals and causing protein redox state changes [30, 112, 113]. The reversible PTM methionine oxidation was studied in *A. thaliana* plants by Jacques et al. (2015) [114] using wild and catalase-2 knock-out mutants. Over 500 sites of oxidation in 400 proteins were reported. They showed that the activity of two specific glutathione-S-transferases was significantly reduced upon oxidation. Konert et al. (2015) [115] have gone deeper into the mechanism of ROS signaling and metabolic adjustment in *A. thaliana*. By using genetic, biochemical, and a combination of MS data-dependent acquisition and SRM, they showed how protein phosphatase 2A, through its regulatory B' γ subunit, interact with a cytosolic aconitase and control mitochondrial alternative oxidases.

Interactomics is the most challenging and less developed proteomics approach, but the most concluding one in order to probe *in vitro* and *in situ* better than *in silico* interactions between proteins and proteins and DNA, RNA or metabolites. It will help in developing models from networks mediating receptor capture of signals and gene expression changes. The standard proteomics analysis involves the capture of protein partners (affinity purification or immunoprecipitation) coupled to MS analysis. Other published approaches included *in vivo* cross-linking with para-formaldehyde and

immunoaffinity purification of cross-linked protein. It has been used by Obermeyer et al., (2014) [116] in the search for H(+) ATPase complexes in lily pollen. These techniques require high amount of sample and involve excessive manipulation, being subjected to biases or false positives. Microarray technology will be, with no doubt, the future for more feasible experiments in which dynamic interactions could be observed through time course experiments [117]. As an alternative, the parallel analysis of genomic, transcriptomics, and proteomics experiments could be performed [118].

Some missed relevant papers

Last but not least in the surveyed period are the publications covering topics that deserve to be mentioned in a general plant proteomics review, either because they report on technical innovations or novel applications to the plant biology research. As we did not intend to be exhaustive, and due to space limitation, we will just mention them.

Such is the case of the top-down proteomics [45, 119-121], N-acetylome [122], degradation of proteins with N-termini containing basic amino acids (the Arg/N-end rule) [25], and single cell type proteomics analysis. Plant organs are complex in terms of the different types of tissues and cells that are formed, with each one having its own proteome signature. Thus, for comparative proteomics purposes it is possible that only a discrete number of cells respond to the experimental variable under study and if so, by extracting the whole organ we might be diluting the response. In order to decipher single proteome analysis new methodologies should be developed [86]. Up to now the laser microdissection [123] has been applied, however it has been of very limited use with plants. What is close to be a technical reality for transcriptomics looks impossible for proteomics [124].

Acknowledgements

Thanks to the University of Córdoba (Programas Propio y de Fortalecimiento) and CNPq (AM, and JVJN) for financial support. L.V. was generously funded by the Juan de la Cierva Programme (JCI-2012-12444, Spanish Ministry of Economy and Competitiveness). Rosa Sánchez Lucas is a recipient

of a Formación de Profesorado Universitario fellowship (FPU14/00186, Spanish Ministry of Education, Culture and Sport). The funders had no role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

- [1] Jorin-Novo, J.V., Pascual, J., Sanchez-Lucas, R., Romero-Rodriguez, M.C., et al., Fourteen years of plant proteomics reflected in Proteomics: Moving from model species and 2DE-based approaches to orphan species and gel-free platforms. *Proteomics* 2015, 5-6, 1089-1112.
- [2] Deswal, R., Abat, J.K., Sehrawat, A., Gupta, R., et al., First systematic plant proteomics workshop in Botany Department, University of Delhi: Transferring proteomics knowledge to next-generation researchers and students. *Proteomics* 2014, 14, 1581-1586.
- [3] Dunn, M.J., Proteomics into its third decade. *Proteomics* 2015, 15, 1-2.
- [4] Thongboonkerd, V., LaBaer, J., Domont, G.B., Recent advances of proteomics applied to human diseases. *J. Proteome Res.*, 2014, 13, 4493-4496.
- [5] Kennedy, D., Building agricultural research. *Science* 2014, 346, 13.
- [6] Lühje, S., Deswal, R., Agrawal, G.K., Editorial. Plant-based foods: seed, nutrition and human health. *Proteomics* 2015, 15, 1638.
- [7] Gregorich, Z.R., Ge, Y., Top-down proteomics in health and disease: challenges and opportunities. *Proteomics* 2014, 14, 1195-1210.
- [8] Ebhardt, H.A., Root, A., Sander, C., Aebersold, R., Applications of targeted proteomics in systems biology and translational medicine. *Proteomics* 2015 15, 3193-3208.
- [9] FAO., Globally almost 870 million chronically undernourished - new hunger report. *Food and Agriculture Organization of the United Nations*, Rome 2012. Report at: <http://www.fao.org/news/story/en/item/161819/icode/>.
- [10] Gutiérrez, R.A., Green, P.J., Keegstra, K., Ohlrogge, J.B., Phylogenetic profiling of the *Arabidopsis thaliana* proteome: what proteins distinguish plants from other organisms? *Genome Biol.* 2004, R53.

- [11] Rodríguez, R.M., López-Vázquez, A., López-Larrea, C., Immune systems evolution. *Adv. Exp. Med. Biol.* 2012, 739, 237-251.
- [12] Suh, L.S., Suh, S.J., Sokol, P.A., Ohman, D.E., A simple alfalfa seedling infection model for *Pseudomonas aeruginosa* strains associated with cystic fibrosis shows AlgT (sigma-22) and RhlR contribute to pathogenesis. *Proc. Natl. Acad. Sci.* 2002, 99, 15699-15704.
- [13] Weidle, U.H., Tiefenthaler, G., Schiller, C., Weiss, E.H., et al., Prospects of bacterial and plant protein-based immunotoxins for treatment of cancer. *Cancer Genom. Proteom.* 2014, 11, 25-38.
- [14] Leung, P.S., Shu, S.A., Chang, C., The changing geoepidemiology of food allergies. *Clin. Rev. Allergy Immunol.* 2014, 46, 169-179.
- [15] Halim, A., Carlsson, M.C., Madsen, C.B., Brand, S., et al., Glycoproteomic analysis of seven major allergenic proteins reveals novel post-translational modifications. *Mol. Cell Proteomics* 2015, 14, 191-204.
- [16] Heidstra, R., Sabatini, S., Plant and animal stem cells: similar yet different. *Nat. Rev. Mol. Cell Biol.* 2014, 15, 301-312.
- [17] McKnight, T.D., Riha, K., Shippen, D.E., Telomeres, telomerase, and stability of the plant genome. *Plant Mol. Biol.* 2002, 48, 331-337.
- [18] Uhlén, M., Fagerberg, Hallström, B.M., Lindskog, C., et al., Proteomics. Tissue-based map of the human proteome. *Science* 2015, 347, 1260419.
- [19] Wilhelm, M., Schlegl, J., Hahne, H., Moghaddas Gholami, A., et al., Mass-spectrometry-based draft of the human proteome. *Nature* 2014, 509, 582-587.
- [20] Sharma, T.K., Ramanathan, R., Rakwal, R., Agrawal, G.K, Bansal, V., Moving forward in plant food safety and security through NanoBioSensors: Adopt or adapt biomedical technologies? *Proteomics* 2015, 15, 1680-1692.
- [21] Fang, X.P., Chen, J.P., Dai, L.Y., Ma, H.S., et al., Proteomic dissection of plant responses to various pathogens. *Proteomics* 2015, 15, 1525-1543.

- [22] Ngara, R., Ndimba, B., Understanding the complex nature of salinity and drought-stress response in cereals using proteomics technologies. *Proteomics* 2014, *14*, 611-621.
- [23] Kim, S.T., Kim, S.G., Agrawal, G.K., Kikuchi, S., Rakwal, R., Rice proteomics: A model system for crop improvement and food security. *Proteomics* 2014, *14*, 593-610.
- [24] Maciel, B.C.M., Barbosa, H.S., Pessôa, G.S., Salazar, M.M., *et al.*, Comparative proteomics and metallomics studies in *Arabidopsis thaliana* leaf tissues: Evaluation of the selenium addition in transgenic and nontransgenic plants using two-dimensional difference gel electrophoresis and laser ablation imaging. *Proteomics* 2014, *14*, 904-912.
- [25] Zhang, H., Deery, M.J., Gannon, L., Powers, S.J. *e al.*, Quantitative proteomics analysis of the Arg/N-end rule pathway of targeted degradation in *Arabidopsis* roots. *Proteomics* 2015, *15*, 2447-2457.
- [26] Venne, A.S., Solari, F.A., Faden, F., Paretti, T., *et al.*, An improved workflow for quantitative N-terminal charge-based fractional diagonal chromatography (ChaFRADIC) to study proteolytic events in *Arabidopsis thaliana*. *Proteomics* 2015, *15*, 2458-2469.
- [27] Walton, A., Stes, E., De Smet, I., Goormachtig, S., Gevaert, K., Plant hormone signaling through the eye of the mass spectrometer. *Proteomics* 2015, *15*, 1113-1126.
- [28] Sakata, K., Komatsu, S., Plant proteomics: from genome sequencing to proteome databases and repositories. *Methods Mol Biol.* 2014, *1072*, 29-42.
- [29] Li, J., Silva-Sanchez, C., Zhang, T., Chen, S., Li, H., Phosphoproteomics technologies and applications in plant biology research. *Front Plant Sci.* 2015, *6*, 430.
- [30] Akter, S., Huang, J., Waszczak, C., Jacques, S., *et al.*, Cysteines under ROS attack in plants: a proteomics view. *J Exp Bot.* 2015, *66*, 2935-2944.

- [31] Tripathi, P., Rabara, R.C., Rushton, P.J.N., A systems biology perspective on the role of WRKY transcription factors in drought responses in plants. *Planta*. 2014. 239, 255-266.
- [32] Jorrin-Novo, J., 2015. Scientific standards and MIAPs in Plant Proteomics research and publications: do we follow them? *Front. Plant Sci.* 2015; 6, 473.
- [33] Lesur, A., Domon, B., Advances in high-resolution accurate mass spectrometry application to targeted proteomics. *Proteomics* 2015, 15, 880-890.
- [34] Lee, D.C.H., Jones, A.R., Hubbard, S.J., Computational phosphoproteomics: From identification to localization. *Proteomics* 2015, 15, 950-963.
- [35] Wu, X.L., Gong, F.P., Wang, W., Protein extraction from plant tissues for 2DE and its application in proteomic analysis. *Proteomics* 2014, 14, 645-658.
- [36] Lee, H.M., Gupta, R., Kim, S.H., Wang, Y., *et al.*, Abundant storage protein depletion from tuber proteins using ethanol precipitation method: Suitability to proteomics study. *Proteomics* 2015, 15, 1765-1769.
- [37] Kim, Y.J., Wang, Y., Gupta, R., Kim, S.W., *et al.*, Protamine sulfate precipitation method depletes abundant plant seed-storage proteins: A case study on legume plants. *Proteomics* 2015, 15, 1760-1764.
- [38] Wildgruber, R., Weber, G., Wise, P., Grimm, D., Bauer, J., Free-flow electrophoresis in proteome sample preparation. *Proteomics* 2014, 14, 629-636.
- [39] Moritz, C.P., Marz, S.X., Reiss, R., Schulenburg, T., Friauf, E., Epicocconone staining: A powerful loading control for Western blots. *Proteomics* 2014, 14, 162-168.
- [40] Lollo, B., Steele, F., Gold, L., Beyond antibodies: New affinity reagents to unlock the proteome. *Proteomics* 2014, 14, 638-644.

- [41] Hildonen, S., Halvorsen, T.G., Reubsaet, L., Why less is more when generating tryptic peptides in bottom-up proteomics. *Proteomics* 2014, *14*, 2031-2041.
- [42] George, I.S., Fennell, A.Y., Haynes, P.A., Protein identification and quantification from riverbank grape, *Vitis riparia*: Comparing SDS-PAGE and FASP-GPF techniques for shotgun proteomic analysis. *Proteomics* 2015, *15*, 3061-3065.
- [43] Valledor, L., Weckwerth, W. In: Jorin-Novo, J., Weckwerth, W., Wienkoop, S., Komatsu, S., (Eds) *Plant Proteomics: Methods and Protocols*. Springer. Germany. pp 347-358.
- [44] Cannon, J.R., Kluwe, C., Ellington, A., Brodbelt, J.S., Characterization of green fluorescent proteins by 193 nm ultraviolet photodissociation mass spectrometry. *Proteomics* 2014, *14*, 1165-1173.
- [45] Dang, X.B., Young, N.L., Ultraviolet photodissociation enhances top-down mass spectrometry as demonstrated on green fluorescent protein variants. *Proteomics* 2014, *14*, 1128-1129.
- [46] Valledor, L., Jorrín, J., Back to the basics: Maximizing the information obtained by quantitative two dimensional gel electrophoresis analyses by an appropriate experimental design and statistical analyses. *J. Proteomics* 2011, *74*, 1-18.
- [47] Krahmer, J., Hindle, M.M., Martin, S.F., Le Bihan, T., Millar, A.J., Sample preparation for phosphoproteomic analysis of circadian time series in *Arabidopsis thaliana*. *Methods Enzymol.* 2015, *551*, 405-431.
- [48] Trotschel, C., Poetsch, A., Current approaches and challenges in targeted absolute quantification of membrane proteins. *Proteomics* 2015, *15*, 915-929.
- [49] Clarke, V.C., Loughlin, P.C., Gavrin, A., Chen, C., *et al.*, Proteomic analysis of the soybean symbiosome identifies new symbiotic proteins. *Mol Cell Proteomics* 2015, *14*, 1301-1322.

- [50] Heard, W., Sklenář, J., Tomé, D.F., Robatzek, S., Jones, A.M., Identification of Regulatory and Cargo Proteins of Endosomal and Secretory Pathways in *Arabidopsis thaliana* by Proteomic Dissection. *Mol. Cell Proteomics* 2015, 14, 1796-1813.
- [51] Dunkley, T.P.J., Watson, R., Griffin, J.L., Dupree, P., Lilley, K.S., Localization of Organelle Proteins by Isotope Tagging (LOPIT). *Mol. Cell Proteomics* 2004, 3, 1128-1134.
- [52] Glibert, P., Van Steendam, K., Dhaenens, M., Deforce, D., iTRAQ as a method for optimization: Enhancing peptide recovery after gel fractionation. *Proteomics* 2014, 14, 680-684.
- [53] Ma, S., Sun, Y., Zhao, X., Xu, P., Recent advance in high accuracy iTRAQ for quantitative proteomics [Article in Chinese]. *Sheng Wu Gong Cheng Xue Bao* 2015, 30, 1073-1082.
- [54] Yu, F., Han, X.S., Geng, C.J., Zhao, Y.X., *et al.*, Comparative proteomic analysis revealing the complex network associated with waterlogging stress in maize (*Zea mays* L.) seedling root cells. *Proteomics* 2015, 15, 135-147.
- [55] Zargar, S.M., Kurata, R., Inaba, S., Oikawa, A., *et al.*, Quantitative proteomics of *Arabidopsis* shoot microsomal proteins reveals a cross-talk between excess zinc and iron deficiency, *Proteomics* 2015, 15, 1196-1201.
- [56] Pan, I.C., Tsai, H.H., Cheng, Y.T., Wen, T.N., *et al.*, Post-Transcriptional Coordination of the *Arabidopsis* Iron Deficiency Response is Partially Dependent on the E3 Ligases RGLG1 and RGLG2. *Mol Cell Proteomics* 2015, 14, 2733-2752.
- [57] González-Fernández, R., Aloria, K., Arizmendi, J.M., Jorrin-Novo, J.V., Application of Label-Free shotgun nUPLC-MS^E and 2-DE approaches in the study of *Botrytis cinerea* mycelium. *J. Proteome Res* 2013, 12, 3042-3056.

- [58] Rogniaux, H., Pavlovic, M., Lupi, R., Lollier, V., *et al.*, Allergen relative abundance in several wheat varieties as revealed via a targeted quantitative approach using MS. *Proteomics* 2015, *15*, 1736-1745.
- [59] Caprioli, R.M., Imaging mass spectrometry: Molecular microscopy for enabling a new age of discovery. *Proteomics* 2015, *15*, 807-809.
- [60] Li, H., Smith, B.K., Shrestha, B., Márk, L., Vertes, A., Automated cell-by-cell tissue imaging and single-cell analysis for targeted morphologies by laser ablation electrospray ionization mass spectrometry. *Methods Mol. Biol.* 2015, *1203*, 117-127.
- [61] Bartels, B., Svatoš, A., Spatially resolved in vivo plant metabolomics by laser ablation-based mass spectrometry imaging (MSI) techniques: LDI-MSI and LAESI. *Front Plant Sci.* 2015, *6*, 471.
- [62] Kiss, A., Smith, D.F., Reschke, B.R., Powell, M.J., Heeren, R.M.A., Top-down mass spectrometry imaging of intact proteins by laser ablation ESIFT-ICR MS. *Proteomics* 2014, *14*, 1283-1289.
- [63] Wijetunge, C.D., Saeed, I., Boughton, B.A., Spraggins, J.M., *et al.*, EXIMS: an improved data analysis pipeline based on a new peak picking method for EXploring Imaging Mass Spectrometry data. *Bioinformatics* 2015 *31*, 3198-3206.
- [64] Franceschi, P., Wehrens, R., Self-organizing maps: A versatile tool for the automatic analysis of untargeted imaging datasets. *Proteomics* 2014, *14*, 853-861.
- [65] Rabilloud, T., Lescuyer, P., The proteomic to biology inference, a frequently overlooked concern in the interpretation of proteomic data: A plea for functional validation. *Proteomics* 2014, *14*, 157-161.
- [66] Fellers, R.T., Greer, J.B., Early, B.P., Yu, X., *et al.*, ProSight Lite: graphical software to analyze top-down mass spectrometry data. *Proteomics* 2015, *15*, 1235-1238.

- [67] Qi, D., Lawless, C., Teleman, J., Levander, F., *et al.*, Representation of selected-reaction monitoring data in the mzQuantML data standard. *Proteomics* 2015, *15*, 2592-2596.
- [68] San Clemente, H., Jamet, E., WallProtDB, a database resource for plant cell wall proteomics. *Plant Methods* 2015, *11*, 2.
- [69] Wen, B., Du, C., Li, G., Ghali, F., *et al.*, IPeak: An open source tool to combine results from multiple MS/MS search engines. *Proteomics* 2015, *15*, 2916-2920.
- [70] Gallardo, O., Ovelleiro, D., Gay, M., Carrascal, M., Abian, J., A collection of open source applications for mass spectrometry data mining. *Proteomics* 2014, *14*, 2275-2279.
- [71] Oveland, E., Muth, T., Rapp, E., Martens, L., *et al.*, Viewing the proteome: How to visualize proteomics data? *Proteomics* 2015, *15*, 1341-1355.
- [72] Bilbao, A., Varesio, E., Luban, J., Strambio-De-Castillia, C., *et al.*, Processing strategies and software solutions for data-independent acquisition in mass spectrometry. *Proteomics* 2015, *15*, 964-980.
- [73] Laukens, K., Naulaerts, S., Vanden Berghe, W., Bioinformatics approaches for the functional interpretation of protein lists: From ontology term enrichment to network analysis. *Proteomics* 2015, *15*, 981-996.
- [74] Mikalsen, S.O., Proteomics made more accessible. *Proteomics* 2014, *14*, 980-990.
- [75] Vaudel, M., Venne, A.S., Berven, F.S., Zahedi, R.P., *et al.*, Shedding light on black boxes in protein identification. *Proteomics* 2014, *14*, 1001-1005.
- [76] Aiche, S., Sachsenberg, T., Kenar, E., Walzer, M., *et al.*, Workflows for automated downstream data analysis and visualization in large-scale computational mass spectrometry. *Proteomics* 2015, *15*, 1443-1447.

- [77] Perez-Riverol, Y., Alpi, E., Wang, R., Hermjakob, H., Vizcaino, J.A., Making proteomics data accessible and reusable: Current state of proteomics databases and repositories. *Proteomics* 2015, *15*, 930-949.
- [78] Tyanova, S., Temu, T., Carlson, A., Sinitcyn, P., et al., Visualization of LC-MS/MS proteomics data in MaxQuant. *Proteomics* 2015, *15*, 1453-1456.
- [79] Smedley D, Haider S, Durinck S, Pandini L, et al., The BioMart community portal: an innovative alternative to large, centralized data repositories. *Nucleic Acids Res* 2015, *43*(W1), W589-598.
- [80] Feussner, I., Polle, A., What the transcriptome does not tell - proteomics and metabolomics are closer to the plants' patho-phenotype, *Curr Opin Plant Biol.* 2015, *26*, 26-31.
- [81] Higdon, R., Kolker, E., Can "normal" protein expression ranges be estimated with high-throughput proteomics? *J Proteome Res.*, 2015, *14*, 2398-2407.
- [82] Smith, S.J., Kroon, J.T., Simon, W.J., Slabas, A.R., et al., A Novel Function for Arabidopsis CYCLASE1 in Programmed Cell Death Revealed by Isobaric Tags for Relative and Absolute Quantitation (iTRAQ) Analysis of Extracellular Matrix Proteins, *Mol Cell Proteomics* 2015, *14*, 1556-1568.
- [83] Hussein, S.M., Puri, M.C., Tonge, P.D., Benevento, M., et al., Genome-wide characterization of the routes to pluripotency. *Nature* 2014, *516*, 198-206.
- [84] Fraas, S., Lüthen, H., Novel imaging-based phenotyping strategies for dissecting crosstalk in plant development, *J. Exp. Bot.* 2015, *66*, 4947-2955.
- [85] Stitt, M., Gibon, Y., Why measure enzyme activities in the era of systems biology? *Trends Plant Sci.*, 2014, *19*, 256-265.
- [86] Misra, B.B., Assmann, S.M., Chen S., Plant single-cell and single-cell-type metabolomics. *Trends Plant Sci.* 2014, *19*, 637-646.

- [87] Kage, U., Kumar, A., Dhokane, D., Karre, S., Kushalappa, A.C., Functional molecular markers for crop improvement. *Crit Rev Biotechnol.* 2015, 14, 1-14.
- [88] Zivy, M., Wienkoop, S., Renaut, J., Pinheiro, C., *et al.*, The quest for tolerant varieties: the importance of integrating "omics" techniques to phenotyping. *Front Plant Sci.* 2015, 6, 448.
- [89] Browning, K.S., Bailey-Serres, J., Mechanism of cytoplasmic mRNA translation. *Arabidopsis Book* 2015, 13, e0176.
- [90] Gupta, R., Min, C.W., Kim, S.W., Wang, Y., *et al.*, Comparative investigation of seed coats of brown- versus yellow-colored soybean seeds using an integrated proteomics and metabolomics approach. *Proteomics* 2015, 15, 1706-1716.
- [91] Barros, E., Lezar, S., Anttonen, M.J., Van Dijk, J.P., Comparison of two GM maize varieties with a near-isogenic non-GM variety using transcriptomics, proteomics and metabolomics. *Plant Biotechnol J.* 2010, 8, 436-451.
- [92] Decourcelle, M., Perez-Fons, L., Baulande, S., *et al.*, Combined transcript, proteome, and metabolite analysis of transgenic maize seeds engineered for enhanced carotenoid synthesis reveals pleiotropic effects in core metabolism. *J. Exp. Bot.* 2015, 66, 3141-3150.
- [93] Valledor, L., Escandon, M., Meijon, M., Nukarinen, E., *et al.*, A universal protocol for the combined isolation of metabolites, DNA, long RNAs, small RNAs, and proteins from plants and microorganisms. *Plant J.* 2014; 79, 173-180.
- [94] Gupta, N., Tanner, S., Jaitly, N., Adkins, J.N., *et al.* Whole proteome analysis of post-translational modifications: Applications of mass-spectrometry for proteogenomic annotation. *Genome Res.* 2007, 17, 1362-1377.
- [95] Kucharova, V., Wiker, H.G., Proteogenomics in microbiology: Taking the right turn at the junction of genomics and proteomics. *Proteomics* 2015, 14, 2660-2675.

- [96] Wang, X.J., Liu, Q., Zhang, B., Leveraging the complementary nature of RNA-Seq and shotgun proteomics data. *Proteomics* 2014, *14*, 2676-2687.
- [97] Romero-Rodríguez, C., Pascual, J., Valledor, L., Jorrín-Novó, J.V., Improving the quality of protein identification in non-model species. Characterization of *Quercus ilex* seed and *Pinus radiata* needle proteomes by using SEQUEST and custom databases. *J. Proteomics* 2014, *105*, 85-91.
- [98] Armengaud, J., Trapp, J., Pible, O., Geffard, O., *et al.*, Non-model organisms, a species endangered by proteogenomics. *J. Proteomics* 2014, *105*, 5-18.
- [99] Gawron, D., Gevaert, K., Van Damme, P., The proteome under translational control. *Proteomics* 2015, *14*, 2647-2659.
- [100] Chapman, B., Bellgard, M., High-throughput parallel proteogenomics: A bacterial case study. *Proteomics* 2014, *14*, 2780-2789.
- [101] De-la-Peña, C., Loyola-Vargas, V.M., Biotic interactions in the rhizosphere: a diverse cooperative enterprise for plant productivity. *Plant Physiol.* 2014, *166*, 701-719.
- [102] Gupta, R., Lee, S.E., Agrawal, G.K., Rakwal, R., *et al.*, Understanding the plant-pathogen interactions in the context of proteomics-generated apoplastic proteins inventory. *Front Plant Sci.* 2015, *6*, 352. [103] Hurley, B., Subramaniam, R., Guttman, D.S., Desveaux, D., Proteomics of effector-triggered immunity (ETI) in plants. *Virulence* 2014, *5*, 752-760.
- [104] Zdarska, M., Dobisová, T., Gelová, Z., Pernisová, M., *et al.*, Illuminating light, cytokinin, and ethylene signalling crosstalk in plant development. *J Exp Bot.* 2015, *66*, 4913-4931.
- [105] Heuberger, A.L., Robison, F.M., Lyons, S.M., Broeckling, C.D., Prenni, J.E., Evaluating plant immunity using mass spectrometry-based metabolomics workflows. *Front Plant Sci.* 2014, *5*, 291.
- [106] Walley, J., Xiao, Y., Wang, J.Z., Baidoo, E.E., Plastid-produced interorganelle stress signal MEcPP potentiates induction of the unfolded protein response in endoplasmic reticulum. *Proc Natl Acad Sci USA.* 2015, *112*, 6212-6217.

- [107] Grones, P., Griml, J., Auxin transporters and binding proteins at a glance. *J Cell Sci.* 2015, 128, 1-7.
- [108] Silva-Sanchez, C., Li, H.Y., Chen, S.X., Recent advances and challenges in plant phosphoproteomics. *Proteomics* 2015, 15, 1127-1141.
- [109] Han, D., Jin, J., Yu, J., Kim, K., Kim, Y., Integrated approach using multistep enzyme digestion, TiO₂ enrichment, and database search for in-depth phosphoproteomic profiling. *Proteomics* 2015, 15, 618-623.
- [110] Choudhary, M.K., Nomura, Y., Wang, L., Nakagami, H., Somers, D.E. Quantitative Circadian Phosphoproteomic Analysis of *Arabidopsis* Reveals Extensive Clock Control of Key Components in Physiological, Metabolic, and Signaling Pathways. *Mol Cell Proteomics* 2015, 14, 2243-2260.
- [111] Minkoff, B.B., Stecker, K.E., Sussman, M.R., Rapid Phosphoproteomic Effects of Abscissic Acid (ABA) on Wild-Type and ABA Receptor-Deficient *A. thaliana* Mutants. *Mol Cell Proteomics* 2015, 14, 1169-1182.
- [112] Liu, P., Zhang, H.M., Wang, H., Xia, Y.J., Identification of redox-sensitive cysteines in the *Arabidopsis* proteome using OxiTRAQ, a quantitative redox proteomics method. *Proteomics* 2014, 14, 750-762.
- [113] Dietz, K.J., Redox regulation of transcription factors in plant stress acclimation and development. *Antioxid Redox Signal.* 2014, 21, 1356-1372.
- [114] Jacques, S., Ghesquière, B., De Bock, P.J., Demol, H., *et al.*, Protein Methionine Sulfoxide Dynamics in *Arabidopsis thaliana* under Oxidative Stress. *Mol Cell Proteomics* 2015, 14, 1217-1229.
- [115] Konert, G., Trotta, A., Kouvonen, P., Rahikainen, M., *et al.*, Protein phosphatase 2A (PP2A) regulatory subunit B'γ interacts with cytoplasmic ACONITASE 3 and modulates the abundance of AOX1A and AOX1D in *Arabidopsis thaliana*. *New Phytol.* 2015, 205, 1250-1263.

- [116] Pertl-Obermeyer, H., Schulze, W.X., Obermeyer, G., In vivo cross-linking combined with mass spectrometry analysis reveals receptor-like kinases and Ca(2+) signalling proteins as putative interaction partners of pollen plasma membrane H(+) ATPases. *J Proteomics* 2014, 108, 17-29.
- [117] Wang, H., Li, H., Zhang, W., Wei, L.M., *et al.*, Multiplex profiling of glycoproteins using a novel bead-based lectin array. *Proteomics* 2014, 14, 78-86.
- [118] Wang, J.X., Peng, X.Q., Peng, W., Wu, F.X., Dynamic protein interaction network construction and applications. *Proteomics* 2014, 14, 338-352.
- [119] Dang, X.B., Scotcher, J., Wu, S., Chu, R.K., *et al.*, The first pilot project of the consortium for top-down proteomics: A status report. *Proteomics* 2014, 14, 1130-1140.
- [120] Erba, E.B., Investigating macromolecular complexes using top-down mass spectrometry. *Proteomics* 2014, 14, 1259-1270.
- [121] Gault, J., Malosse, C., Machata, S., Millien, C., *et al.*, Complete posttranslational modification mapping of pathogenic *Neisseria meningitidis* pilins requires top-down mass spectrometry. *Proteomics* 20214, 14, 1141-1151.
- [122] Dinh, T.V., Bienvenut, W.V., Linster, E., Feldman-Salit, A., *et al.*, Molecular identification and functional characterization of the first N α -acetyltransferase in plastids by global acetylome profiling. *Proteomics* 2015, 15, 2426-2435.
- [123] Gautam, V., Sarkar, A.K., Laser assisted microdissection, an efficient technique to understand tissue specific gene expression patterns and functional genomics in plants. *Mol Biotechnol.* 2015, 57, 299-308.
- [124] Klein, A.M., Mazutis, L., Akartuna, I., Tallapragada, N., *et al.*, Droplet Barcoding for Single-Cell Transcriptomics Applied to Embryonic Stem Cells. *Cell* 2015, 161, 1187-1201.