Graphical Abstract

Highlights

- 1st comprehensive review of major limitations of PAH removal by Fenton oxidation.
- Limitations under review are acidic pH, PAH unavailability and soil deterioration.
- Critical review of various strategies to address these limitations is presented.
- Gaps and future research needs are highlighted for application at field scale.

**Fenton oxidation to remediate PAHs in contaminated soils: A critical review of major
limitations and counter-strategies**

M. Usman^{a, b*}, K. Hanna^c, S. Haderlein^a

^a Department of Geosciences, Center for Applied Geosciences, University of Tübingen,
72074 Tübingen, Germany

^b Institute of Soil and Environmental Sciences, University of Agriculture, Faisalabad 38040,
Pakistan

^c Ecole Nationale Supérieure de Chimie de Rennes, UMR CNRS 6226, 11 Allée de Beaulieu,
35708 Rennes Cedex 7, France

* Corresponding author:

Dr. M. Usman

Research Fellow (Alexander von Humboldt Foundation)

Department of Geosciences, Center for Applied Geosciences, University of Tübingen

Hölderlinstr. 12, 72074 Tübingen, Germany

Email: muhammad.usman@uaf.edu.pk or: musmanch@yahoo.com

Phone No.: +49 70 71 29 78 924

Fax: +49 70 71 29 50 59

Abstract

Fenton oxidation constitutes a viable remediation strategy to remove polycyclic aromatic hydrocarbons (PAHs) in contaminated soils. This review is intended to illustrate major limitations associated with this process like acidification, PAH unavailability, and deterioration of soil quality along with associated factors, followed by a critical description of various developments to overcome these constraints. Considering the limitation that its optimal pH is around 3, traditional Fenton treatment could be costly, impractical in soil due to the high buffering capacity of soils and associated hazardous effects. Use of various chelating agents (organic or inorganic) allowed oxidation at circumneutral pH but factors like higher oxidant demand, cost and toxicity should be considered. Another alternative is the use of iron minerals that can catalyze Fenton-like oxidation over a wide range of pH, but mobility of these particles in soils (i.e. saturated and unsaturated zones) should be investigated prior to *in-situ* applications. The PAH-unavailability is the crucial limitation hindering their effective degradation. Research data is compiled describing various strategies to address this issue like the use of availability enhancement agents, extraction or thermal pretreatment. Last section of this review is devoted to describe the effects of various developments in Fenton treatment onto soil quality and native microbiota. Finally, research gaps are discussed to suggest future directions in context of applying Fenton oxidation to remediate contaminated soils.

Keywords

PAH Remediation; Fenton oxidation; pH; PAH unavailability; Soil quality

1. Introduction

Polycyclic aromatic hydrocarbons (PAHs) are organic compounds composed of two or more fused benzene rings. Their contamination in environment is of great concern due their high toxicity, carcinogenic effects and environmental persistence. There are sixteen PAHs that are considered as priority pollutants by United States Environmental Protection Agency (US EPA) and European community. For being recalcitrant in nature, PAHs tend to persist in the environment resulting in a ubiquitous distribution. The European Environment Agency (EEA) located about 1.5 millions of contaminated sites in Europe including 200,000 sites contaminated PAHs (E.E.A, 2012). Soil often acts as the ultimate depository of PAHs particularly due to their hydrophobic nature (Haritash and Kaushik, 2009). Owing to the strong persistence and ubiquitous occurrence, substantial research has been devoted to develop techniques to remove persistent PAHs from complex matrix like soil or sediments. Chemical oxidation has emerged as a viable remediation strategy to degrade PAHs in contaminated soils (Lee et al., 1998; Watts et al., 2002; Rivas, 2006; Ferrarese et al., 2008; Yap et al., 2011). The most common oxidants used in this technique are permanganate, ozone, persulfate and Fenton's reagent (H_2O_2 combining with iron) (ITRC, 2005). Among these oxidants, Fenton oxidation is showing promising potential to rapidly degrade PAHs in contaminated soils. Fenton treatments utilize the high reactivity of hydroxyl radical ($\cdot\text{OH}$), a very reactive chemical species, which is generated by the decomposition of hydrogen peroxide (H_2O_2) by iron through following reaction (Fenton, 1894):

Over the last two decades, use of Fenton oxidation has been well documented to treat PAHs in contaminated soils by numerous research studies. Apart from research data, few reviews focusing on chemical oxidation/Fenton-based treatments specifically for PAHs alone or along with other persistent organic pollutants in contaminated soils have been published (Rivas,

2006; Gan et al., 2009; Yap et al., 2011; Venny et al., 2012a). First review about chemical oxidation of PAHs was completed by F.J. Rivas (Rivas, 2006) where he focused mainly on the Fenton oxidation, ozone and high temperature pressurized water after a detailed discussion about sorption and extraction of PAHs from contaminated soils. Various techniques available for the remediation of PAH contaminated soils were reviewed by Gan and co-workers (2009) including bioremediation, solvent extraction, bioremediation, phytoremediation, chemical oxidation, photocatalytic degradation, electrokinetic remediation, thermal treatment and their integration. Yap and co-workers (2011) presented the first review specifically for the application of Fenton-based treatments (conventional, modified, Fenton-like along with integrated approaches) for PAHs-contaminated soils. Review by Venny and co-workers (2012a) described the fundamental principles and governing factors of Fenton oxidation to remediate soils contaminated with PAHs, polychlorinated biphenyls (PCBs), polychlorinated dibenzo-p-dioxins and furans and coupling of Fenton oxidation with other strategies like bioremediation and eletrokinetic remediation.

To the best of our knowledge, a systematic review illustrating the major limitations of Fenton treatment along with proposed solutions to overcome these constraints is still missing in literature. Thus, this review article is intended to collect research data related to potential constraints of Fenton oxidation in context of PAH contaminated soils. In addition, recent developments to overcome these limitations are critically evaluated along with associated factors in this review. These drawbacks would depend on the remedial strategy and scale, and the targeted zone in soils (e.g. aquifer or vadose zone). Despite the large amount of published studies related to Fenton treatments for PAH-contaminated soils at lab-scale (Yap et al., 2011), very limited data is available on the remediation performance at pilot/field scale (Liu et al., 1993; Brown, 1997; Lingle et al., 2001; ITRC, 2005; Taylor, 2012). Prior to the field or *in-situ* applications, feasibility studies are generally conducted at batch scale to determine the

95 extent and rate of PAH removal. However, few reports have investigated the Fenton
96 oxidation under column dynamic conditions, which more closely mimic field conditions
97 (Palmroth et al., 2006a; Laurent et al., 2012; Venny et al., 2012c; Lemaire et al., 2013b;
98 Venny et al., 2014). Figure 1 shows the different experimental setups used to investigate
99 Fenton oxidation in contaminated soils at laboratory scale. Column studies, which are better
100 representative of real field conditions, can be carried out under water-saturated (that
101 represents aquifer) and water-unsaturated conditions (that represent vadose zone).
102 We are convinced that this review article will be of significant interest for researchers
103 working on chemical oxidation of PAHs-contaminated soils. Finally, the literature lack of
104 information and data that is crucial prior to *in-situ* field applications is also highlighted.

106 **2. Major limitations of Fenton oxidation in PAH-contaminated soil**

107 Fenton oxidation has validated its potential as viable remediation technique for PAH-
108 contaminated soils but there are various limitations associated with this process. This review
109 article focuses on the three major limitations of this technique in context of PAH-
110 contaminated soil including: i) acidification of medium, ii) PAH unavailability and iii)
111 deterioration of soil quality, fertility and microbial community. These limitations are
112 discussed in the following sections along with a critical discussion of their solutions proposed
113 in literature.

115 **2.1. Acidification of reaction medium**

116 Traditional Fenton's reagent ($\text{Fe}^{\text{II}}\text{-H}_2\text{O}_2$) displays its maximum oxidation activity at a pH
117 about 2.8 – 3 (Safarzadeh-Amiri et al., 1996; Pignatello et al., 2006). Thus, most of the
118 remediation studies to remove PAHs in contaminated soils employed traditional Fenton
119 reagent at acidic pH (Yap et al., 2011) which is probably the most important limitation of this

process. This initial acidification (optimum pH is 3) is required to keep Fe^{II} in its soluble form (Tamura et al., 1980). The iron precipitation occurring after pH 4.0 led to the obstruction of activation processes (Usman et al., 2012), through several parasite reactions including the decomposition of H_2O_2 to oxygen. In addition, sorption of Fe^{II} or Fe^{III} ions onto mineral or organic constituents in soil increases with increase in pH (Hanna et al., 2013; Catrouillet et al., 2014) and thus, rendering the Fe species less available for Fenton activation.

This acidification step is operated not only to keep Fe in dissolved form but also because the $\bullet\text{OH}$ generation in reaction (1) is highest at pH 3 that usually decreases at higher or lower pH values (Gallard et al., 1998). Furthermore, reaction (1) actually involves the formation of a highly oxidised Fe adduct, often indicated as ferryl (e.g. ferryl ion FeO^{2+}), which does not necessarily evolve into $\bullet\text{OH}$ (Pignatello et al., 1999).

Initial acidification required for optimum efficiency of Fenton's reagent is costly and impractical in soil system due to buffering capacity of soils (Ershadi et al., 2011). This could also create a hazardous situation for soil quality and microbes in soil buffered systems, where a circumneutral pH prevails (Nam et al., 2001; Sahl and Munakata-Marr, 2006; Sirguey et al., 2008; Valderrama et al., 2009). Therefore, traditional Fenton oxidation was found incompatible with subsequent biological treatment to remediate PAH contaminated soil (Nam et al., 2001). It may also increase mobilization of heavy metals that would devastate the soil ecosystem. Acidic pH along with high amount of Fe^{II} was also linked to a remarkable increase in available Fe and Al in treated soil, which reached the toxicity level (Gan et al., 2013). This decrease in pH during Fenton oxidation was also found as the major limiting factor for subsequent vegetation in treated soil (Sirguey et al., 2008). It was reported by Valderrama and co-workers (2009) that use of more aggressive traditional Fenton oxidation does not favor the subsequent biological treatment which was explained by the residual

oxidant that may remain in the soil and inhibit the biodegradation after the application of higher oxidant dose.

Therefore, various strategies were considered in literature that allows chemical oxidation at circumneutral pH but without compromising the oxidation efficiency for efficient PAH degradation like the use of chelating agents or iron minerals. These strategies will be detailed in the following paragraphs.

2.1.1 Use of Chelating agents

Use of chelating agents along with traditional Fenton treatment allows chemical oxidation at circumneutral pH and the reaction is termed as modified Fenton (MF) oxidation (Nam et al., 2001). Chelating agents (CAs) are mostly organic compounds that bind iron and therefore maintain in its dissolved form. By binding with Fe^{II} or Fe^{III} , CAs can maintain their ability to catalyze H_2O_2 at circumneutral pH. These chelating agents are used to extend the applicability of Fenton oxidation to native soil pH that has more practical implications for *in-situ* remediation. Different kinds of chelating agents have been explored in MF (Table 1) including cyclodextrines (CD), ethylenediaminetetraacetic acid (EDTA), catechol (CC), gallic acid (GA), citric acid (CA), oxalic acid (OA), humic acid (HA), malic acid (MA), sodium citrate (SC) and other amino-poly-acetic acids (Nam et al., 2001; Gryzenia et al., 2009; Veignie et al., 2009; Venny et al., 2012b).

For example, Nam et al. (Nam et al., 2001) showed that the addition of organic CAs (such as CC and GA) along with ferric ion during MF oxidation (pH 6.0–6.5) could have distinct advantages due to its compatibility with subsequent bioremediation as compared to the traditional Fenton oxidation at pH 2–3. Their results revealed that >80% of low molecular weight (LMW) PAHs and only 20-40% of high molecular weight (HMW) PAHs were degraded during traditional Fenton oxidation. In their attempt to overcome the limitations of

conventional Fenton oxidation, combined MF and biodegradation resulted in > 98% removal of LMW-PAHs and > 70% removal of HMW-PAHs. In their attempt to examine HA as iron chelator in MF oxidation, Georgi and co-workers (2007) reported that presence of HA (50–100 mg L⁻¹) in MF system greatly enhanced the degradation of organic compounds in the range of pH 5–7. They also highlighted that similar degradation extent can be obtained by MF oxidation at pH 5 in the presence of HA and traditional Fenton process at pH 3 in the absence of HA (Georgi et al., 2007).

In addition to binding Fe^{II} ion for enhanced solubility at circumneutral pH, CD (Veignie et al., 2009) or HA (Georgi et al., 2007) also resulted in improving availability of hydrophobic organic pollutants in contaminated soils by forming complexes with pollutants. Formation of pollutant-CD-iron or pollutant-HA-iron complex in microenvironment allows hydroxyl radicals to specifically target on pollutants by eliminating the detrimental soil matrix effect for improved oxidation efficiency (Lindsey et al., 2003; Georgi et al., 2007; Veignie et al., 2009). Owing to the ability of CD to improve pollutant availability, it was also employed in soil washing solution (Viglianti et al., 2006), bioremediation (Bardi et al., 2000) and chemical oxidation (Usman et al., 2012) that improved the removal efficiency many times. A detailed discussion about the use of CD to improve PAH availability is presented below in Section 2.2.1 (Chemical Pretreatment).

Despite the promising efficiency of chelating agents to promote pollutant oxidation, their use is limited by cost, toxicity and potential negative effects on oxidation efficiency due to the non-productive consumption of the CAs (Xue et al., 2009b; Pardo et al., 2014). For being organic in nature, CAs may compete with target pollutants for oxidant and thus act as radical scavenger by increasing oxidant demand (Georgi et al., 2007; Xue et al., 2009b; Pardo et al., 2014). Oxidation of CAs themselves will thereby lead to the loss of their chelating ability which is required to enhance pollutant degradation (Georgi et al., 2007). Moreover, chelating

agents like EDTA can also pose environmental risks because of their persistence in the environment and their contribution to improve heavy metal mobility/bioavailability (Sillanpaa and Pirkanniemi, 2001; Yang et al., 2001; Rastogi et al., 2009).

Recently inorganic CAs like sodium pyrophosphate (SP) were proposed as an alternative in MF oxidation of PAHs in contaminated soils at circumneutral pH (Venny et al., 2012b; Venny et al., 2012c; Jorfi et al., 2013). Comparison of various organic and inorganic CAs (Fig. 2a) to assist MF oxidation revealed the following efficiency order for PAH degradation: SP > EDTA > OA > MA > SC (Venny et al., 2012b). Other studies by this research group have established the higher efficiency of SP during Fenton oxidation and its compatibility with biological treatment to promote PAH degradation (Venny et al., 2012c; Venny et al., 2014). Similarly, Jorfi et al. (Jorfi et al., 2013) performed Fenton oxidation by employing iron nano-particles along with various CAs to remove pyrene in contaminated soil. They reported that removal efficiency was found to be maximum for SP (93%) compared to 86%, 75%, 72% and 71% for EDTA, SC, HA and FA, respectively at $\text{H}_2\text{O}_2/\text{Fe}^{\text{III}}$ molar ratio of 10, pH 7 and the contact time of 6 h. Use of inorganic CAs could be advantageous since they are less likely to act as radical scavenger and do not increase the total organic carbon of system (Rastogi et al., 2009) although prolonged treatment could make SP to act as radical scavenger (Venny et al., 2012c). In addition to less oxidant scavenging, decomposition of H_2O_2 with SP was lower than with organic CAs (Fig. 2b) underscoring higher stability of oxidant when mixed with inorganic SP (Venny et al., 2012b).

2.1.2. Use of Fe-bearing minerals

In order to overcome the drawback of acidification employed in traditional Fenton oxidation, iron mineral can be used instead of soluble Fe^{II} at circumneutral pH, and the reaction is defined as a Fenton-like (FL) reaction. Iron minerals are able to catalyze Fenton-like

reactions over a wide range of pH values (Watts et al., 2002; Hanna et al., 2008; Jung et al., 2009). The water solubility of Fe-oxyhydroxides as well as the possible leaching of iron from their surface is low. The Fe-minerals catalyze oxidation mainly through heterogeneous reaction, i.e. activation of oxidant by Fe surface sites located on oxide surfaces. As a result, the ability of Fe to catalytically decompose H_2O_2 and generate hydroxyl radicals can be maintained with respect to traditional Fenton, but the overall oxidation kinetics is generally lower (Garrido-Ramírez et al., 2010). Various mechanisms are proposed for oxidative degradation of organic compounds on the surface of iron oxide catalysts during FL oxidation and are detailed elsewhere (Xue et al., 2009a; Garrido-Ramírez et al., 2010; Yap et al., 2011). There are various kinds of iron minerals including ferric- Fe^{III} (ferrihydrite, goethite, lepidocrocite, hematite etc.) and mixed Fe^{II} - Fe^{III} oxides (magnetite and green rust) (Cornell and Schwertmann, 2006). Heterogeneous catalysts including native or added iron oxides or certain transition metals can catalyze the FL oxidation in soils at circumneutral pH (Flotron et al., 2005; Usman et al., 2012), which may be especially advantageous for *in situ* remediation of contaminated soils where pH cannot be adjusted.

It was also reported by Jung and co-workers (2009) that H_2O_2 was more stable (as it has a ten times longer life time) under FL reaction (catalyzed by magnetite or hematite) than that with soluble Fe^{II} . Moreover, mineral-catalyzed oxidation system (3 ml of 30% H_2O_2 for 5 g contaminated sand) without pH adjustment showed better degradation (75.3%) of phenanthrene sorbed on sand (200 mg.kg^{-1}) than by Fenton activated by Fe^{II} (52.2%) (Jung et al., 2009).

Remediation of PAHs in contaminated soils was reported by Fenton-like oxidation catalyzed by native iron (Watts et al., 2002; Flotron et al., 2005) or iron minerals added into the system like goethite (Kanel et al., 2003; Kanel et al., 2004), hematite (Jung et al., 2009), iron oxyhydroxide/sodium dodecyl sulfate composites (Park and Kim, 2013) and magnetite

(Usman et al., 2012; Usman et al., 2013; Biache et al., 2015). Magnetite was found more effective than goethite and soluble Fe^{II} at acidic pH to remediate petroleum hydrocarbons in diesel and kerosene contaminated soil (Kong et al., 1998). Higher catalytic efficiency of magnetite, a mixed Fe^{II} - Fe^{III} oxide, as compared to the ferric minerals was correlated to its structural Fe^{II} (Matta et al., 2007; Hanna et al., 2008).

It was shown by Watts and co-workers (2002) that iron mineral present in solid matrix can allow the production of reactive oxidant species for strong degradation of PAHs. They reported that 85% of benzo[a]pyrene was removed when H_2O_2 (15 M) was catalyzed by naturally occurring iron minerals (33.52 mg.g^{-1} of soil) while a degradation of 70% was achieved when H_2O_2 (15 M) was catalyzed by soluble Fe^{II} (6.6 mM) in sand. Role of endogenous iron minerals was highlighted when application of sole H_2O_2 resulted in PAH degradation almost equivalent to that obtained with H_2O_2 + soluble Fe^{II} without pH adjustment in sludge (pH 8.8) and sediment (pH 4.4) bearing 94 mg.g^{-1} , 30 mg.g^{-1} and not determined total iron, respectively (Flotron et al., 2005). They reported similar findings in contaminated soil (pH 7.1) where Fe contents were not determined. Owing to the presence of native iron (16.4 mg.g^{-1}), application of Fenton oxidation (0.4 g $\text{H}_2\text{O}_2.\text{g}^{-1}$ soil) under saturated conditions resulted in significant degradation of PAHs (upto 52%) without any ferrous iron supplementation or initial acidification of soil (Palmroth et al., 2006a). Pardo and co-workers (2016) observed similar degradation of PAHs by Fenton oxidation (882 mmol.L^{-1} without pH adjustment) obtained by adding 1 mmol L^{-1} Fe^{III} or without external addition of Fe highlighting the role of endogenous Fe. They noted a similar concentration of iron in solution after external addition of 1 mmol L^{-1} Fe^{III} or iron extracted from soil (Pardo et al. 2016)

Kanel and co-workers (2004) reported that goethite can successfully catalyze FL oxidation (33.5 g goethite. kg^{-1} of soil and 5 M H_2O_2) with degradation extent of 73, 60, and 55% for

phenanthrene, anthracene, and pyrene, respectively in spiked sandy soil obeyed pseudo-first order kinetics at natural pH. Use of magnetite (Fig. 3) to catalyze FL oxidation (H_2O_2 :Fe of 20, 10% w/w magnetite) at circumneutral pH (7 – 8) resulted in PAH degradation extent of 90% and 60% in spiked sandy soil and in historically contaminated soils (sandy loamy type) sampled from former coking plant site, respectively (Usman et al., 2012). Under the similar experimental conditions, soluble Fe^{II} was found unable to catalyze H_2O_2 as similar oxidation efficiency (20% in spiked sandy soil and 5% in real soil) was observed when H_2O_2 was used alone without iron activation and that was correlated to the precipitation of Fe^{II} at circumneutral pH in tested soils. Lower degradation in real soil samples as compared to the spiked sandy soil was related to the soil matrix effect. Magnetite-catalyzed FL oxidation (oxidant/Fe molar ratio equal to 10:1) also resulted in strong PAH removal (60–70 %) under flow through conditions (flow rate 0.1 mL min^{-1} under water-saturated columns) in spiked sandy soil having 10% of magnetite w/w (Usman et al., 2013). Biache and co-workers (2015) also highlighted the strong efficiency of magnetite-catalyzed FL oxidation (14.5 mL of H_2O_2 (30%wt), 500 mg magnetite for 5 g soil) to degrade PAHs in three contaminated soils sampled from former coking plant site (20% removal), former gas plant site soil (92%) and active wood treatment facility (50%). Recently, pyrite (FeS_2) showed strong reactivity to catalyze FL oxidation of PAHs in soil washing wastewater at circumneutral pH (Choi et al., 2014). They reported a sustainable reactivity of this process that was linked to slow but continuous Fe^{II} dissolution from the pyrite surface leads to the continuous formation of reactive radicals during the oxidation reaction (Choi et al., 2014).

Pardo and co-workers (2016) recently conducted a study to optimize the application of the Fenton oxidation to remediate PAH contaminated soils. They evaluated the suitability of nZVI particles and Fe^{III} salt at different concentrations of initial iron ($1\text{--}5 \text{ mmol L}^{-1}$) and oxidant ($88\text{--}1765 \text{ mmol L}^{-1}$) without pH adjustment. Almost similar degradation of PAHs

was achieved for the majority of the experiments. But, they reported that aqueous concentration of iron was lower in case of nZVI and they linked to its lower solubility as compared to Fe^{III} salt at circumneutral pH. Owing to this behavior, Pardo and co-workers (2016) suggested that nZVI could act as a continuous Fe source that can be advantageous for real field application.

Regardless of the investigated Fe-bearing minerals or nZVI particles to promote the *in-situ* application of the oxidation treatment, profound investigations are required to evaluate injection mode of these particles in soils. It will depend on the targeted area in soil whether saturated aquifer (pump and treat) or unsaturated zone such as vadose zone (e.g. lysimeter system). It is known that the efficiency of treatment significantly depends on the contact “pollutant/catalyst/oxidant” (Lemaire et al., 2013a), and therefore particles with a high mobility must easily reach the contaminated targeted zones (Johnson and Amy, 1995). This mobility is strongly related to the composition of particles including functionalized or not, particle size (micro- or nano-sized), concentration of suspension, etc. (Jiang et al., 2010). Despite the mobility and colloidal transport of natural particles and some manufactured nanoparticles have been already studied (Hannah and Thompson, 2008; Lowry et al., 2012), there is limited information on the transport and fate of the catalyst particles used for remediation purposes.

Apart from these strategies, few studies reported the use of Fenton’s reagent (H_2O_2 + soluble Fe^{II}) to remove PAHs without adjusting the soil’s pH to avoid this initial acidification but that leads to the partial degradation of PAHs (Kulik et al., 2006; Silva et al., 2009; Laurent et al., 2012).

2.2. PAH unavailability

Another major constraint associated with Fenton oxidation in aged contaminated soils is the poor availability of PAHs for remediation (Flotron et al., 2005; Usman et al., 2012; Lemaire et al., 2013a; Choi et al., 2014) due to their strong hydrophobicity and partitioning into soil constituents especially organic matter (Riding et al., 2013). This is caused by the deep diffusion of the contaminants into the organic material that renders them unavailable for the hydroxyl radicals, which are generated in the aqueous phase (Bogan et al., 2003). A strong connection was reported between PAH availability and remediation efficiency of Fenton oxidation (Usman et al., 2013; Biache et al., 2015; Usman et al., 2016). As a matter of fact, magnetite-catalyzed FL oxidation was not effective unless PAH availability was enhanced in two aged coking plant soils (Usman et al., 2012). Due to this strong correlation, Biache and co-workers (2015) proposed that efficiency of Fenton based oxidation could serve as a valid method to estimate PAH availability in contaminated soils. In order to suppress this limitation and thus effectively degrade PAHs in contaminated soils, various strategies were considered and are discussed below.

2.2.1 Chemical pretreatment

Chemical pretreatment involves the use of various chemical reagents to enhance the PAH availability in contaminated soils as a pretreatment for or in combination with Fenton oxidation. Four types of enhancement-solubility agents have been used including co-solvent, surfactant, cyclodextrin and vegetable oil to solubilize PAHs. Co-solvents like ethanol were used to increase solubility of PAH in soils that ultimately reduces the surface tension between the compounds in soil and aqueous phase, and increase in PAH availability (Bonten et al., 1999).

344 Ethanol-Fenton (ETF) treatment resulted in significant increase in PAH removal of 11–28%
345 (Lundstedt et al., 2006), 97% (Lee et al., 1998) and 63–93% (Lee et al., 2002) than by Fenton
346 alone in contaminated soils (Fig. 4). Application of ETF treatment in soil spiked with
347 benz(a)anthracene (500 mg/Kg of soil) using optimized experimental conditions (1.0 ml of
348 ethanol, 0.2 ml of 0.5 M Fe^{II} , and 0.3 ml of 30% H_2O_2 per g of soil) removed 97% of target
349 PAH and that neither water nor sodium dodecyl sulfate (SDS as surfactant) show a
350 substantial effect on removal (Lee et al., 1998). Ethanol was widely used as a solvent because
351 of its safety, low cost, and excellent extraction ability (Lee and Hosomi, 2001a).

352 Recently, Yap and co-workers (2012) reported the use of a novel system, ethyl lactate-Fenton
353 (ELF) treatment for soils artificially contaminated with four PAHs (500 mg/kg each of
354 phenanthrene, anthracene, fluoranthene and benzo[a]pyrene). Their results (Fig. 4) indicate
355 that ELF ($[\text{H}_2\text{O}_2] = 666 \text{ mM}$, $\text{Fe}^{\text{II}} = 66.6 \text{ mM}$ with ethyl lactate, $\text{pH} = 3$) resulted in higher
356 PAH degradation (up to 97%) than ETF (70%) while conventional Fenton oxidation (without
357 ethanol or ethyl lactate) resulted in 31–57% of untreated PAHs. A recent study by the same
358 research group revealed the feasibility of using ELF to treat PAHs in aged soils (pH 4.4 and
359 6.2) having various quantities of soil organic carbon (Yap et al., 2015). Similar trend of
360 higher PAH degradation by ELF ($\text{H}_2\text{O}_2 = 667.6 \text{ mM}$, $\text{Fe}^{\text{II}} = 16.7 \text{ \& } 66.7 \text{ mM}$ with varying
361 amounts of ethyl lactate) as compared to ETF or traditional Fenton (Fig. 4) was also indicated
362 elsewhere (Gan et al., 2013). However, addition of co-solvents could change the general non-
363 selective behavior of $\cdot\text{OH}$ radicals towards selectivity as preferential degradation of
364 anthracene > benzo[a]pyrene > phenanthrene > fluoranthene was observed by ELF and ETF
365 treatments (Yap et al., 2012). They also suggested that degree of selectivity is co-solvent
366 dependent as selectivity was more pronounced in the ETF treatment than in the ELF
367 treatment. Moreover, presence of co-solvents during Fenton oxidation could also inhibit PAH
368 degradation by acting as radical scavenger (Qiang et al., 2000; Lundstedt et al., 2006; Yap et

al., 2012). But scavenging effect was more pronounced for EL than ET and it was correlated to the higher bond ratio of C–O to C–H and higher oxidation state of EL (Yap et al., 2012). This scavenging effect associated with ethanol was avoided by its use as a pretreatment step (5 mL of ethanol/g soil for 24 h) where it was allowed to evaporate before soil samples were subjected to Fenton-like oxidation (Usman et al., 2012). Treating ethanol pre-treated soil by Fenton-like (FL) oxidation (magnetite + H₂O₂) caused 20% removal of PAHs (Fig. 5). It is worthy to note that no PAH degradation was observed when FL oxidation was used directly without any chemical pre-treatment (Usman et al., 2012). On the other hand, FL oxidation after pretreatment using cyclodextrin (CD, 10 mM, overnight mixing) resulted in slightly lower PAH degradation (15%) as compared to ethanol pretreatment (Usman et al., 2012) probably due to scavenging effect of CD (Hanna et al., 2005). This scavenging effect concerns only CD in study by Usman and co-workers (2012), as ethanol was allowed to evaporate from the reactor before applying oxidants.

Use of surfactants was also found effective to improve the PAH availability decreasing the interfacial tension and increasing their partitioning to the hydrophobic cores of surfactant micelles. First use of surfactants in combination with Fenton's reagent was reported by Martens and Frankenberger (Martens and Frankenberger Jr, 1995) who found that use of sodium dodecyl sulfate (SDS) before Fenton oxidation resulted in higher PAH degradation (30-45%) compared to without SDS. Application of cationic surfactant cetylpyridinium chloride (CPC) in the pyrite Fenton system enhanced the PAH oxidation efficiency from 91% (without CPC) to 97% at circumneutral pH (Choi et al., 2014). Recently, use of SDS as surfactant with H₂O₂ + Fe^{III} resulted in higher PAH removal than the systems without SDS including H₂O₂ + Fe^{III} or H₂O₂ + nanoparticles of zerovalent iron (nZVI) (Pardo et al., 2016). It should be noted that surfactants can also be formed as a result of partial oxidation of hydrocarbons and/or native organic matter (Ndjou'ou and Cassidy, 2006; Gryzenia et al.,

2009). These authors demonstrated the potential of *in-situ* surfactant production during Fenton oxidation of hydrocarbon-contaminated soils. These surfactants accompanying the Fenton oxidation of hydrocarbons in soil are eventually removed by further chemical oxidation and/or biodegradation (Ndjou'ou and Cassidy, 2006; Gryzenia et al., 2009). Moreover, formation of reductants (e.g. superoxide) during Fenton oxidation could act as surfactant and contribute in enhanced desorption of PAHs (Watts et al., 1999). Non-selective degradation of PAHs was also linked to this probable production of surfactants during Fenton-like oxidation (Usman et al., 2012) that would have rendered HMW PAHs available for remediation (Gryzenia et al., 2009). Pretreatment of contaminated soils with vegetable oil prior to Fenton oxidation also improved the PAH degradation in contaminated soils (Bogan et al., 2003). They incorporated two different oils (corn oil and palm kernel oil) in Fenton treatment for PAH removal from highly contaminated aged soils. A significant increase in the PAH degradation (15–45%) was reported with 5% (per dry wt. of soil) oil addition, with a pronounced effect for HMW PAHs. On the other hand, Fenton treatment (without vegetable oil) could only degrade a few LMW PAHs.

2.2.2. Extraction pretreatment

It is well documented that artificially spiked PAHs (fresh contamination) are easier to oxidize than native PAHs because of their binding/sequestration onto soil constituents for years in later case (Nam et al., 2001; Flottron et al., 2005; Usman et al., 2012; Jorfi et al., 2013; Lemaire et al., 2013b). For example, Nam and co-workers (2001) described that Fenton treatment caused higher PAH removal in model spiked soil as compared to that in soil from former manufactured gas plant (MGP). They reported a highly efficient degradation of a mixture of PAHs in the model soil (noticeably, 84.5% and 96.7% of initial pyrene and Benzo(a)pyrene, respectively). In the MGP soil, the same treatment destroyed approximately

40% and 20% of pyrene and Benzo(a)pyrene, respectively. Flotron and co-workers (2005) also reported more efficient degradation of spiked PAHs (onto soil, sediment and sludge) than native PAHs in soil after Fenton oxidation.

Fenton oxidation ($\text{H}_2\text{O}_2:\text{Fe}^{\text{II}}$ of 20:1) in column experiments under water-unsaturated conditions caused significantly higher PAH degradation for spiked soils (55%) as compared to aged soils (30%) (Lemaire et al., 2013b). In another work, removal of pyrene in freshly contaminated soil was greater (91%) than that of the soil sample with aged pollution (43%) when MF oxidation (300 mM H_2O_2 , 5mM Fe^{III}) using iron nano-particles was applied (Jorfi et al., 2013). The pyrene degradation extent decreased from 88.9% in freshly contaminated soil to 74.7% in soil aged for 30 days although further increase in aging time did not significantly decrease the removal efficiency (Sun and Yan, 2008). As a matter of fact, incubation of a freshly contaminated coking plant soil by a ^{13}C -phenanthrene indicated that bacteria rapidly degraded the freshly-added PAH whereas the native contamination of the coking plant soil (including phenanthrene) is not affected (Cébron et al., 2011). But all these studies using spiked soils were based upon the PAHs whose solutions were freshly prepared in laboratory and were not extracted from the field soil.

This was done in a recent study where two ancient coking plant soils contaminated with 16 PAHs were extracted and solvent extractable organic matter (EOM) separated from these soils was spiked on sand (Usman et al., 2012). Then, application of Fenton-like oxidation ($\text{H}_2\text{O}_2:\text{Fe}$ of 20:1, 10% magnetite w/w, neutral pH) (FL) resulted in significant degradation of PAHs (90%) in sand spiked with EOM (Fig. 5). It should be noted that FL oxidation was unable to degrade PAHs when applied directly in both soils. It might be correlated to the long term aging of both soils as coking activities and steel industry continued on these soils from the very beginning of the 20th century till 1983 (Ouvrard et al., 2011). As a next step of study, Usman and co-workers (2012) applied extraction pretreatment to re-incorporate the PAHs

into the soil as fresh contaminants. It was done by following the same protocol as was used to separate EOM (for spiking sand) but without separating it from the soil, rather it was allowed to evaporate in soil. This procedure allowed PAH removal of 50-60% in re-contaminated soils by successive Fenton-like oxidation (oxidation conditions similar to spiked sand) as indicated in Fig. 5 (Usman et al., 2012). This drop in degradation efficiency between real soil (50-60%) and spiked sand (90%) was correlated to soil matrix effect because of the insoluble organic matter and mineral fraction (Flotron et al., 2005; Jonsson et al., 2007; Usman et al., 2012). But, application of similar pretreatment allowed a six-times higher PAH availability than raw soil measured by Tenax extraction that is closer to bioavailable fraction (Cébron et al., 2013). This difference could be caused by different PAH availability levels including bioavailable (Tenax available) and chemoavailable (deduced by FL oxidation) probably due to higher accessibility for chemical oxidation as suggested elsewhere (Cébron et al., 2013). Very few studies used the extraction pre-treatment to re-incorporate PAHs in same soil as fresh contaminants (Usman et al., 2012; Cébron et al., 2013). Owing to its strong efficiency to improve PAH availability, the *in-situ* feasibility of extraction pretreatment merits to be explored at field scale. It should be noted that cyclodextrin or surfactant-enhanced *in-situ* flushing and extraction of hydrocarbons compounds have been tested at the field scale (McCray and Brusseau, 1999; McCray et al., 2000; Bardi et al., 2007; Chong et al., 2014). However to the best of our knowledge, the use of solubility-enhancement agents in combination with chemical oxidation has been never tested at field scale.

2.2.3 Thermal pretreatment

Pre-heating PAH contaminated soils also improved the remediation efficiency of successive Fenton-like treatment (Biache et al., 2015; Usman et al., 2016). Recently, Usman and co-workers (2016) pre-heated the two ancient coking plant soils at different temperatures (60,

100, and 150 °C for 1 week under inert atmosphere). Pre-heating resulted in slight removal of PAHs (6% for A soil and 9% for B soil even at 150 °C) but successive FL oxidation (H_2O_2 :Fe of 20:1, 10% magnetite w/w, neutral pH) in pre-heated soils showed significant PAH degradation (19, 29, and 43 % in A soil and 31, 36, and 47 % in B soil pre-treated at 60, 100, and 150 °C, respectively, reference for degradation were pre-heated samples) (Fig. 5). No PAH degradation was reported when soils were subjected to chemical oxidation without pre-treatment (Usman et al., 2016). Similarly, Biache and co-workers (2015) reported an increase in PAH degradation in preheated soil samples (100 °C for 1 week under inert atmosphere) by H_2O_2 oxidation systems (including FL oxidation by 14.5 mL of H_2O_2 (30%wt), 500 mg magnetite for 5 g soil, neutral pH). Their experiments were based on three kinds of soils that were sampled from former coking plant, former gas plant and an active wood treating facility. Both coking and gas plant soils were characterized by similar EOM (10.3 and 11.5 mg/g, respectively) and PAH content (1146 and 1781 $\mu\text{g/g}$, respectively) while soil from wood-treating facility represented higher amount of EOM (81.94 mg/g) and PAHs (c.a. 12,000 $\mu\text{g.g}^{-1}$). The PAH distribution varied for each sample origin and was dominated by 4- and 5-ring PAHs (coking plant soil) or 2- to 4-ring PAHs (gas plant soil) or by pyrene and fluoranthene (wood treatment facility). Their results indicate that efficiency of FL oxidation varied according to the type of the soil. They did not report any PAH degradation by direct FL oxidation in coking plant soil unless the sample was pre-heated (20% degradation by successive FL oxidation). On the other hand, FL oxidation in the unheated and pre-heated gas plant soils showed the highest degradation rates (78% and 92%, respectively). The oxidative degradation of PAHs reached 49% and 50%, for the unheated and pre-heated wood-treating facility soils, respectively. Results obtained by Biache and co-workers (2015) clearly demonstrate that PAH unavailability was mainly related in coking plant soil due to its long-term aging (Ouvrard et al., 2011; Usman et al., 2012; Usman et al.,

2016). It was reported elsewhere that efficiency of Fenton oxidation was negatively correlated to the age of the PAH contamination when experiments were performed by using ten different soil samples originating from former gas plant, former wood-treating facility and active coke production site and characterized by different contamination age ranging from 40 to 71 years (Jonsson et al., 2006; Jonsson et al., 2007). In aforementioned coking plant soil (Biache et al. 2015, Usman et al. 2016), no such unavailability constraint was reported during air oxidation of PAHs at 100 °C (Biache et al., 2011) which highlights the role of temperature to control PAH availability. Similarly, heating aged PAH contaminated soils at higher temperature (120 °C) prior to microbial incubation enhanced the PAH degradation due to an increase in their bioavailability (Bonten et al., 1999). Mechanisms governing thermal pretreatment enhanced PAH availability has not been yet explored due to the limited number of relevant studies. Moreover, further studies are required to test the upscaling of thermal pretreatment for field applications while nature of the soil should be considered.

2.3. Deterioration of soil quality and impacts on microbial community

Fenton-based oxidation treatments helps to decrease the PAH load in contaminated soils but it could negatively affect the soil quality and microbial community. This section illustrates the effect of various Fenton-based strategies being used to remove PAHs from contaminated soils.

Impact of Fenton oxidation (without pH adjustment or pH adjusted at 3 with H₂SO₄ 1 N) onto soil quality was investigated by Sirguey and co-workers (2008) in spiked agricultural soil and coking plant soil. Significant degradation (75-97%) of PAHs was reported whereas associated decrease in soil pH was found to be the limiting factor for plant growth. This treatment resulted in a significant decline of organic C and N along with strong changes in

nutrient pool. Water retention capacity was however improved upon treatment that could encourage re-vegetation (Sirguey et al., 2008).

Laurent and co-workers (2012) applied MF oxidation ($\text{H}_2\text{O}_2 + \text{Fe}^{\text{II}}$ without pH adjustment) under unsaturated conditions in industrial PAH-contaminated soil. A degradation of 14% and 22% of PAHs was obtained with addition of H_2O_2 at 6 and 65 g kg⁻¹, respectively. The results obtained at higher oxidant dose (65 g kg⁻¹) indicated a significant decrease in soil pH (4.8 decreased from 7.3), cation exchange capacity (2 units) and extractable phosphorus content (77% of initial content). Although drop in pH was not as harsh as it could be if initial acidification (low to 3) was adopted, it contributed along with higher oxidant amount to decrease bacterial, fungal, and PAH degrading bacteria population that rebounded after five weeks of incubation (Fig. 6). Plant growth was also negatively affected at higher oxidant dose. Treated soil also displayed a significant delay in seed germination and a decrease in root and shoot biomass (Laurent et al., 2012).

A column study was conducted by Palmroth et al. (Palmroth et al., 2006a) where soil pH was not acidified prior to MF oxidation (0.4 g H_2O_2 /g soil without any ferrous iron addition) to minimize the stress on soil bacteria. Obtained results indicated a partial removal of PAHs (up to 52%). But toxicity of column leachate was increased towards *Vibrio fischeri* that was used to assess the effect of MF oxidation on microbial activity. Microbial activity and abundance of viable cells was lower in incubated MF-treated soil than in untreated soil. Still, incubation of MF-treated soil slightly improved the PAH-removal than in untreated soil highlighting its compatibility as also reported in another study (Palmroth et al., 2006b).

The high concentration of H_2O_2 used and the radicals formed in the Fenton's reaction are toxic to soil bacteria and, therefore, reduce the abundance of viable cells (Büyüksönmez et al., 1998). Apart from that, the acidic pH of 2–3 required for the traditional Fenton oxidation can disturb the soil ecosystem and thus, may be incompatible with biological treatment (Nam

et al. 2001). But application of this treatment at near neutral pH could be compatible with bioremediation (Nam et al., 2001). Ecological impacts and environmental concerns due to the presence of H₂O₂ can be significantly addressed by coupling with biological treatment. Moreover, combination of Fenton oxidation and bioremediation was found more efficient to degrade PAHs than sole process and thus, coupling was suggested to improve the removal efficiency and to minimize associated risks (Kulik et al., 2006; Palmroth et al., 2006a; Palmroth et al., 2006b; Valderrama et al., 2009; Venny et al., 2012c). A pilot scale study designed to treat MGP soil revealed that use of Fenton oxidation as the pre-treatment followed by biological treatment using an aerobic biosystem can decrease the PAHs to meet environmental standards (Liu et al., 1993). Moreover, no volatile organic compounds (VOCs) were observed during air monitoring from these experiments implying the safety of this process for environment. Integrated bioremediation and Fenton oxidation in bioslurry reactors also showed more than 80% degradation of PAHs while total microbial counts ranged up to 10⁴ CFU/mL despite the severe oxidation conditions (Brown, 1997). Experimental conditions should be carefully optimized to improve compatibility of both processes as suggested in above studies. Moreover, it should be noted that microbes tend to restore their population and activity after few days (Sahl and Munakata-Marr, 2006; Sirguey et al., 2008). But, too much H₂O₂ during Fenton oxidation may potentially lead to undesired soil sterilization as it destroys the native micro-biota during the treatment (Valderrama et al., 2009).

Venny and co-workers (2012c) demonstrated the strong compatibility of MF oxidation (Fe^{III} + SP as chelating agent) with bioremediation (natural attenuation and biostimulation) for an efficient degradation of PAHs in column experiments simulating in-situ soil flushing. These authors, in another study in packed columns (Venny et al., 2014), described the impact of SP modified Fenton oxidation on soil functionality during remediation of PAHs in spiked and

aged soils. Use of optimized operating parameters during MF oxidation ($\text{H}_2\text{O}_2/\text{soil}=0.081$, $\text{Fe}^{\text{III}}/\text{soil}=0.024$, $\text{SP}/\text{soil}=0.024$, pH of SP solution=7.73) could make this treatment compatible with bioremediation and crop growth as the obtained values for pH, electrical conductivity, iron precipitation and soil respiration were within acceptable limits.

Gan et al. (Gan et al., 2013) compared the impact of ethyl lactate based Fenton (ELF) treatment onto soil quality with that of conventional water based Fenton treatment (CWF). It was found that ELF caused higher loss of soil organic matter than CWF that was correlated to its high efficiency. Moreover, a decrease in CEC, available P, Cu and Zn was observed in both treatments but to a greater extent in CWF, while ELF caused higher decrease in specific surface area and loss of N. Both treatments, when conducted at low pH (3-4) and high Fe^{II} dosage ($\text{H}_2\text{O}_2:\text{Fe}^{\text{II}} = 10:1$), caused Fe and Al toxicity. But no excessive Fe or Al was formed when ELF was employed at higher pH (≥ 6.2) and lower amounts of added Fe^{II} . This study suggested that ELF adopted for soil with such higher native pH could be compatible for revegetation purposes (Gan et al., 2013).

In addition, traditional Fenton oxidation has been known to frequently generate toxic transformation products due to partial mineralization of PAHs that could also contribute towards soil toxicity (Lee and Hosomi, 2001a; Flotron et al., 2005; Lundstedt et al., 2006). It has been reported that anthracene and benzo[a]anthracene were degraded by the Fenton oxidation by 97% and each target substance was further transformed to oxygenated PAHs by 68% (Lee and Hosomi, 2001b) and 39% (Lee et al., 1998), respectively. It has been reported that oxy-PAHs are more persistent than the target PAHs and some of them were not degraded any further during Fenton oxidation (Lundstedt et al., 2006). Moreover, owing to the higher solubility, these oxygenated PAHs (dibenzofuran, 9-H fluorenone and anthraquinone) are known to be more mobile than PAHs in US-EPA list (Lundstedt et al., 2014). Thus, their distribution and behavior was also focused along with target PAHs in many chemical

oxidation studies (Choi et al., 2014; Biache et al., 2015; Usman et al., 2016). No such by-products were detected during magnetite-catalyzed FL oxidation (Usman et al., 2016) or pyrite-Fenton system (Choi et al., 2014). A similar evolution observed for oxy-PAHs and target PAHs suggested a non-preferential removal by FL oxidation (Usman et al., 2016).

3. Conclusions and Outlook

Fenton oxidation is widely used technique to remediate PAHs in contaminated soils but various limitations are associated with its environmental application. This review is intended to provide the readers a critical illustration of various limitations of this process along with a brief discussion of strategies to overcome these limitations. The practical applicability of this method strongly depends on the soil type and structure, history of contamination, and concentration and physical state of the contaminants. A rigorous investigation of all these parameters is a pre-requisite for effective application in soil remediation.

Considering the main drawback that its optimal pH is around 3, the traditional Fenton treatment is costly and impractical due to high buffering capacity of soils and associated hazardous effects. Use of modified Fenton oxidation by using chelating agents can allow its use at circumneutral pH but chelating agents can also compete for $\cdot\text{OH}$ radical with PAHs and thus significantly affecting the oxidation efficiency. A recent approach in this regard is the use of inorganic chelating agents that offers advantages over organic chelating agents for oxidant consumption. Another possibility is the use of iron minerals, which are able to catalyze Fenton-like oxidation over a wide range of pH. Due to the presence of structural Fe^{II} , magnetite showed strong reactivity to catalyze Fenton-like oxidation of PAHs at circumneutral pH. Moreover, its catalytic stability can be maintained for multiple oxidation cycles and it can be a promising treatment. Use of iron minerals or nanoparticles of ZVI could provide a cost-effective alternative to the use of more than one reagent (e.g. chelating

agent + soluble Fe^{II}). Further studies are needed for a better understanding of Fenton-like application in real field conditions, as injection of iron particles in unsaturated zone such as vadose zone requires many investigations. Mobility, transport and fate of such particles should be addressed on context of reactivity, stability and eco-toxicity. Detailed studies are required to exploit the potential of endogenous iron minerals to catalyze chemical oxidation. Another major limitation is the PAH unavailability during Fenton oxidation. Many strategies are discussed including the use of availability enhancement agents, extraction or thermal pretreatment. PAH availability is the crucial factor controlling its degradation in aged soils and that should be considered while designing a remediation treatment. Use of suitable pretreatment to enhance PAH availability can enhance the treatment efficiency. Most of the studies are based on the batch experiments that should be up-scaled to column experiments under saturated and unsaturated conditions to get closer to real field conditions. The field application of certain pretreatments (extraction or thermal) seems difficult to implement that merits to be resolved but they can serve as good approach for *ex-situ* application. Integration of Fenton's treatment with bioremediation could become a suitable option to address the detrimental effects associated with Fenton's application. The development of new modeling approaches that account for the up-scaling of different involved reactions and complexity of soil system becomes urgent in soil remediation science. More experimental work is also needed to develop strategies with high remediation efficiency and yet are suitable for soil quality so that contaminated sites can be restored to productive use.

Acknowledgment

This work was supported by a post-doctoral research fellowship from the Alexander von Humboldt Foundation, which is gratefully acknowledged.

References

- Bardi L, Mattei A, Steffan S, Marzona M. Hydrocarbon degradation by a soil microbial population with [beta]-cyclodextrin as surfactant to enhance bioavailability. *Enzyme Microb. Technol.* 2000; 27: 709-713.
- Bardi L, Martini C, Opsi F, Bertolone E, Belviso S, Masoero G, Marzona M, Marsan FA. Cyclodextrin-enhanced in situ bioremediation of polyaromatic hydrocarbons-contaminated soils and plant uptake. *J. Incl. Phenom. Macro. Chem.* 2007; 57: 439-444.
- Biache C, Ghislain T, Faure P, Mansuy-Huault L. Low temperature oxidation of a coking plant soil organic matter and its major constituents: An experimental approach to simulate a long term evolution. *J. Hazard. Mater.* 2011; 188: 221-230.
- Biache C, Lorgeoux C, Andriatsihoarana S, Colombano S, Faure P. Effect of pre-heating on the chemical oxidation efficiency: Implications for the PAH availability measurement in contaminated soils. *J. Hazard. Mater.* 2015; 286: 55-63.
- Bogan BW, Trbovic V, Paterek JR. Inclusion of vegetable oils in Fenton's chemistry for remediation of PAH-contaminated soils. *Chemosphere* 2003; 50: 15-21.
- Bonten LTC, Grotenbuis TC, Rulkens WH. Enhancement of PAH biodegradation in soil by physicochemical pretreatment. *Chemosphere* 1999; 38: 3627-3636.
- Brown K. Innovative methods for bioslurry treatment. Site Remediation Technology Report. National Risk Management Research Laboratory. Office of Research and Development. US Environment Protection Agency, Cincinnati, Ohio 1997; 45268.
- Büyüksönmez F, Hess TF, Crawford RL, Watts RJ. Toxic effects of modified Fenton reactions on *Xanthobacter flavus* FB71. *Appl. Environ. Microb.* 1998; 64: 3759-3764.
- Catrouillet C, Davranche M, Dia A, Bouhnik-Le Coz M, Marsac R, Pourret O, Gruau G. Geochemical modeling of Fe(II) binding to humic and fulvic acids. *Chem. Geol.* 2014; 372: 109-118.
- Cébron A, Louvel B, Faure P, France-Lanord C, Chen Y, Murrell JC, Leyval C. Root exudates modify bacterial diversity of phenanthrene degraders in PAH-polluted soil but not phenanthrene degradation rates. *Environ. Microbiol.* 2011; 13: 722-736.
- Cébron A, Faure P, Lorgeoux C, Ouvrard S, Leyval C. Experimental increase in availability of a PAH complex organic contamination from an aged contaminated soil: Consequences on biodegradation. *Environ. Pollut.* 2013; 177: 98-105.
- Choi K, Bae S, Lee W. Degradation of pyrene in cetylpyridinium chloride-aided soil washing wastewater by pyrite Fenton reaction. *Chem. Eng. J.* 2014; 249: 34-41.
- Chong Z-Y, Liao X-Y, Yan X-L, Sun L, Zhao D, Liang T. Enhanced Desorption of PAHs from Manufactured Gas Plant Soils Using Different Types of Surfactants. *Pedosphere* 2014; 24: 209-219.
- Cornell RM, Schwertmann U. The Iron Oxides: Structure, Properties, Reactions, Occurrence and Uses: Wiley-VCH, 2006.
- E.E.A. <http://www.eea.europa.eu/data-and-maps/figures/overview-ofcontaminants-affecting-soil-and-groundwater-in-europe>, 2012.
- Ershadi L, Ebadi T, Ershadi V, Rabbani AR. Chemical oxidation of crude oil in oil contaminated soil by Fenton process using nano zero valent Iron. 2nd International Conference on Environmental Science and Technology. 6, 2011, pp. V1-89.
- Fenton HJH. Oxidation of tartaric acid in presence of iron. *J. Chem. Soc. T.* 1894; 65: 899-910.
- Ferrarese E, Andreottola G, Oprea IA. Remediation of PAH-contaminated sediments by chemical oxidation. *J. Hazard. Mater.* 2008; 152: 128-139.

- Flotron V, Delteil C, Padellec Y, Camel V. Removal of sorbed polycyclic aromatic hydrocarbons from soil, sludge and sediment samples using the Fenton's reagent process. *Chemosphere* 2005; 59: 1427-1437.
- Gallard H, de Laat J, Legube B. Effect of pH on the oxidation rate of organic compounds by Fe-II/H₂O₂. Mechanisms and simulation. *New J. Chem.* 1998; 22: 263-268.
- Gan S, Lau EV, Ng HK. Remediation of soils contaminated with polycyclic aromatic hydrocarbons (PAHs). *J. Hazard. Mater.* 2009; 172: 532-549.
- Gan S, Yap CL, Ng HK, Venny. Investigation of the impacts of ethyl lactate based Fenton treatment on soil quality for polycyclic aromatic hydrocarbons (PAHs)-contaminated soils. *J. Hazard. Mater.* 2013; 262: 691-700.
- Garrido-Ramírez EG, Theng BKG, Mora ML. Clays and oxide minerals as catalysts and nanocatalysts in Fenton-like reactions — A review. *Appl. Clay Sci.* 2010; 47: 182-192.
- Georgi A, Schierz A, Trommler U, Horwitz CP, Collins TJ, Kopinke FD. Humic acid modified Fenton reagent for enhancement of the working pH range. *Appl. Catal. B, Environ.* 2007; 72: 26-36.
- Gryzenia J, Cassidy D, Hampton D. Production and accumulation of surfactants during the chemical oxidation of PAH in soil. *Chemosphere* 2009; 77: 540-545.
- Hanna K, Chiron S, Oturan MA. Coupling enhanced water solubilization with cyclodextrin to indirect electrochemical treatment for pentachlorophenol contaminated soil remediation. *Water Res.* 2005; 39: 2763-2773.
- Hanna K, Kone T, Medjahdi G. Synthesis of the mixed oxides of iron and quartz and their catalytic activities for the Fenton-like oxidation. *Catal. Commun.* 2008; 9: 955-959.
- Hanna K, Usman M, Chatain V. Sorption and transport of aqueous FeII in a goethite-coated sand column under anoxic conditions. *Appl. Geochem.* 2013.
- Hannah W, Thompson PB. Nanotechnology, risk and the environment: a review. *J. Environ. Monit.* 2008; 10: 291-300.
- Haritash AK, Kaushik CP. Biodegradation aspects of Polycyclic Aromatic Hydrocarbons (PAHs): A review. *J. Hazard. Mater.* 2009; 169: 1-15.
- ITRC. Technical and Regulatory Guidance for In Situ Chemical Oxidation of Contaminated Soil and Groundwater. ITRC, ISCO Team, Washington, DC, 2005.
- Jiang X, Tong M, Li H, Yang K. Deposition kinetics of zinc oxide nanoparticles on natural organic matter coated silica surfaces. *J. Colloid Interface Sci.* 2010; 350: 427-434.
- Johnson WP, Amy GL. Facilitated Transport and Enhanced Desorption of Polycyclic Aromatic Hydrocarbons by Natural Organic Matter in Aquifer Sediments. *Environ. Sci. Technol.* 1995; 29: 807-817.
- Jonsson S, Persson Y, Frankki S, Lundstedt S, van Bavel B, Haglund P, Tysklind M. Comparison of Fenton's reagent and ozone oxidation of polycyclic aromatic hydrocarbons in aged contaminated soils. *J. Soils Sediment.* 2006; 6: 208-214.
- Jonsson S, Persson Y, Frankki S, van Bavel B, Lundstedt S, Haglund P, Tysklind M. Degradation of polycyclic aromatic hydrocarbons (PAHs) in contaminated soils by Fenton's reagent: A multivariate evaluation of the importance of soil characteristics and PAH properties. *J. Hazard. Mater.* 2007; 149: 86-96.
- Jorfi S, Rezaee A, Moheb-ali G-a, Jaafarzadeh Na. Pyrene removal from contaminated soils by modified Fenton oxidation using iron nano particles. *J. Environ. Health Sci. Eng.* 2013; 11: 17.
- Jung YS, Lim WT, Park JY, Kim YH. Effect of pH on Fenton and Fenton- like oxidation. *Environ. Technol.* 2009; 30: 183-190.

- Kanel SR, Neppolian B, Choi H, Yang JW. Heterogeneous catalytic oxidation of phenanthrene by hydrogen peroxide in soil slurry: Kinetics, mechanism, and implication. *Soil Sediment Contam.* 2003; 12: 101-117.
- Kanel SR, Neppolian B, Jung H, Choi H. Comparative removal of polycyclic aromatic hydrocarbons using iron oxide and hydrogen peroxide in soil slurries. *Environ. Eng. Sci.* 2004; 21: 741-751.
- Kong SH, Watts RJ, Choi JH. Treatment of petroleum-contaminated soils using iron mineral catalyzed hydrogen peroxide. *Chemosphere* 1998; 37: 1473-1482.
- Kulik N, Goi A, Trapido M, Tuhkanen T. Degradation of polycyclic aromatic hydrocarbons by combined chemical pre-oxidation and bioremediation in creosote contaminated soil. *J. Environ. Manag.* 2006; 78: 382-391.
- Laurent F, Cébron A, Schwartz C, Leyval C. Oxidation of a PAH polluted soil using modified Fenton reaction in unsaturated condition affects biological and physico-chemical properties. *Chemosphere* 2012; 86: 659-664.
- Lee BD, Hosomi M, Murakami A. Fenton oxidation with ethanol to degrade anthracene into biodegradable 9, 10-anthraquinone: A pretreatment method for anthracene-contaminated soil. *Water Sci. Technol.* 1998; 38: 91-97.
- Lee BD, Hosomi M. A hybrid Fenton oxidation-microbial treatment for soil highly contaminated with benz(a)anthracene. *Chemosphere* 2001a; 43: 1127-1132.
- Lee BD, Hosomi M. Fenton oxidation of ethanol-washed distillation-concentrated benzo(A)pyrene: Reaction product identification and biodegradability. *Water Res.* 2001b; 35: 2314-2319.
- Lee BD, Nakai S, Hosomi M. Application of fenton oxidation to remediate polycyclic aromatic hydrocarbons-contaminated soil. *J. Chem. Eng. Jpn.* 2002; 35: 582-586.
- Lemaire J, Buès M, Kabeche T, Hanna K, Simonnot M-O. Oxidant selection to treat an aged PAH contaminated soil by in situ chemical oxidation. *J. Environ. Chem. Eng.* 2013a; 1: 1261-1268.
- Lemaire J, Laurent F, Leyval C, Schwartz C, Buès M, Simonnot M-O. PAH oxidation in aged and spiked soils investigated by column experiments. *Chemosphere* 2013b; 91: 406-414.
- Lindsey ME, Xu G, Lu J, Tarr MA. Enhanced Fenton degradation of hydrophobic organics by simultaneous iron and pollutant complexation with cyclodextrins. *Sci. Total Environ.* 2003; 307: 215-229.
- Lingle J, Hellman E, Noel M, Wilson J. Full Scale Chemical Oxidation of Manufactured Gas Plant Hydrocarbons in Soil and Groundwater Using Fenton's Reagent, 2001.
- Liu BY, Srivastava V, Paterek J, Pradhan S, Pope J, Hayes T, Linz D, Jerger D. MGP soil remediation in a slurry-phase system: a pilot-scale test. Institute of Gas Technology, Chicago, IL (United States), 1993.
- Lowry GV, Gregory KB, Apte SC, Lead JR. Transformations of Nanomaterials in the Environment. *Environ. Sci. Technol.* 2012; 46: 6893-6899.
- Lundstedt S, Persson Y, Öberg L. Transformation of PAHs during ethanol-Fenton treatment of an aged gasworks' soil. *Chemosphere* 2006; 65: 1288-1294.
- Lundstedt S, Bandowe BAM, Wilcke W, Boll E, Christensen JH, Vila J, Grifoll M, Faure P, Biache C, Lorgeoux C. First intercomparison study on the analysis of oxygenated polycyclic aromatic hydrocarbons (oxy-PAHs) and nitrogen heterocyclic polycyclic aromatic compounds (N-PACs) in contaminated soil. *TrAC, Trends Anal. Chem.* 2014; 57: 83-92.
- Martens DA, Frankenberger Jr WT. Enhanced degradation of polycyclic aromatic hydrocarbons in soil treated with an advanced oxidative process—Fenton's Reagent. *Soil Sediment Contam.* 1995; 4: 175-190.

791 Matta R, Hanna K, Chiron S. Fenton-like oxidation of 2,4,6-trinitrotoluene using different
 792 iron minerals. *Sci. Total Environ.* 2007; 385: 242-251.

793 McCray JE, Brusseau ML. Cyclodextrin-Enhanced In Situ Flushing of Multiple-Component
 794 Immiscible Organic Liquid Contamination at the Field Scale: Analysis of Dissolution
 795 Behavior. *Environ. Sci. Technol.* 1999; 33: 89-95.

796 McCray JE, Boving TB, Brusseau ML. Cyclodextrin-Enhanced Solubilization of Organic
 797 Contaminants with Implications for Aquifer Remediation. *Ground Water Monit. R.*
 798 2000; 20: 94-103.

799 Nam K, Rodriguez W, Kukor JJ. Enhanced degradation of polycyclic aromatic hydrocarbons
 800 by biodegradation combined with a modified Fenton reaction. *Chemosphere* 2001; 45:
 801 11-20.

802 Ndjou'ou AC, Cassidy D. Surfactant production accompanying the modified Fenton
 803 oxidation of hydrocarbons in soil. *Chemosphere* 2006; 65: 1610-1615.

804 Ouvrard S, Barnier C, Bauda P, Beguiristain T, Biache C, Bonnard M, Caupert C, Cébron A,
 805 Cortet J, Cotellet S, Dazy M, Faure P, Masfaraud JF, Nahmani J, Palais F, Poupin P,
 806 Raoult N, Vasseur P, Morel JL, Leyval C. In Situ Assessment of Phytotechnologies
 807 for Multicontaminated Soil Management. *Int. J. Phytorem.* 2011; 13: 245-263.

808 Palmroth MRT, Langwaldt JH, Aunola TA, Goi A, Münster U, Puhakka JA, Tuhkanen TA.
 809 Effect of modified Fenton's reaction on microbial activity and removal of PAHs in
 810 creosote oil contaminated soil. *Biodegradation* 2006a; 17: 131-141.

811 Palmroth MRT, Langwaldt JH, Aunola TA, Goi A, Puhakka JA, Tuhkanen TA. Treatment of
 812 PAH-contaminated soil by combination of Fenton's reaction and biodegradation. *J.*
 813 *Chem. Technol. Biotechnol.* 2006b; 81: 598-607.

814 Pardo F, Rosas J, Santos A, Romero A. Remediation of a biodiesel blend-contaminated soil
 815 by using a modified Fenton process. *Environ Sci Pollut Res* 2014; 21: 12198-12207.

816 Pardo F, Peluffo M, Santos A, Romero A. Optimization of the application of the Fenton
 817 chemistry for the remediation of a contaminated soil with polycyclic aromatic
 818 hydrocarbons. *J. Chem. Technol. Biotechnol.* 2016; 91: 1763-1772.

819 Park J-Y, Kim J-H. Role of sol with iron oxyhydroxide/sodium dodecyl sulfate composites
 820 on Fenton oxidation of sorbed phenanthrene in sand. *J. Environ. Manag.* 2013; 126:
 821 72-78.

822 Pignatello JJ, Liu D, Huston P. Evidence for an Additional Oxidant in the Photoassisted
 823 Fenton Reaction. *Environ. Sci. Technol.* 1999; 33: 1832-1839.

824 Pignatello JJ, Oliveros E, MacKay A. Advanced oxidation processes for organic contaminant
 825 destruction based on the fenton reaction and related chemistry. *Crit. Rev. Environ.*
 826 *Sci. Technol.* 2006; 36: 1-84.

827 Qiang Z, Chang JH, Huang CP, Cha D. Oxidation of Selected Polycyclic Aromatic
 828 Hydrocarbons by the Fenton's Reagent: Effect of Major Factors Including Organic
 829 Solvent. *Nuclear Site Remediation*. 778. American Chemical Society, 2000, pp. 187-
 830 209.

831 Rastogi A, Al-Abed SR, Dionysiou DD. Effect of inorganic, synthetic and naturally
 832 occurring chelating agents on Fe(II) mediated advanced oxidation of chlorophenols.
 833 *Water Res.* 2009; 43: 684-694.

834 Riding MJ, Doick KJ, Martin FL, Jones KC, Semple KT. Chemical measures of
 835 bioavailability/bioaccessibility of PAHs in soil: Fundamentals to application. *J.*
 836 *Hazard. Mater.* 2013; 261: 687-700.

837 Rivas FJ. Polycyclic aromatic hydrocarbons sorbed on soils: A short review of chemical
 838 oxidation based treatments. *J. Hazard. Mater.* 2006; 138: 234-251.

839 Safarzadeh-Amiri A, Bolton JR, Cater SR. The use of iron in advanced oxidation processes.
 840 *J. Adv. Oxid. Technol.* 1996: 18-26.

- Sahl J, Munakata-Marr J. The effects of in situ chemical oxidation on microbiological processes: A review. *Remed. J.* 2006; 16: 57-70.
- Sillanpaa M, Pirkanniemi K. Recent developments in chelate degradation. *Environ. Technol.* 2001; 22: 791-801.
- Silva PTdSe, Silva VLd, Neto BdB, Simonnot MO. Phenanthrene and pyrene oxidation in contaminated soils using Fenton's reagent. *J. Hazard. Mater.* 2009; 161: 967-973.
- Sirguez C, Tereza de Souza e Silva P, Schwartz C, Simonnot M-O. Impact of chemical oxidation on soil quality. *Chemosphere* 2008; 72: 282-289.
- Sun H-w, Yan Q-s. Influence of pyrene combination state in soils on its treatment efficiency by Fenton oxidation. *J. Environ. Manag.* 2008; 88: 556-563.
- Tamura H, Kawamura S, Hagayama M. Acceleration of the oxidation of Fe²⁺ ions by Fe(III)-oxyhydroxides. *Corros. Sci.* 1980; 20: 963-971.
- Taylor KM. The Chemical Oxidation of Polycyclic Aromatic Hydrocarbons at a Former Manufactured Gas Plant in Bay Shore, New York. 2012.
- Usman M, Faure P, Ruby C, Hanna K. Remediation of PAH-contaminated soils by magnetite catalyzed Fenton-like oxidation. *Appl. Catal. B, Environ.* 2012; 117-118: 10-17.
- Usman M, Faure P, Lorgeoux C, Ruby C, Hanna K. Treatment of hydrocarbon contamination under flow through conditions by using magnetite catalyzed chemical oxidation. *Environ. Sci. Pollut. Res.* 2013; 20: 22-30.
- Usman M, Chaudhary A, Biache C, Faure P, Hanna K. Effect of thermal pre-treatment on the availability of PAHs for successive chemical oxidation in contaminated soils. *Environ. Sci. Pollut. Res.* 2016; 23: 1371-1380.
- Valderrama C, Alessandri R, Aunola T, Cortina JL, Gamisans X, Tuhkanen T. Oxidation by Fenton's reagent combined with biological treatment applied to a creosote-contaminated soil. *J. Hazard. Mater.* 2009; 166: 594-602.
- Veignie E, Rafin C, Landy D, Fourmentin S, Surpateanu G. Fenton degradation assisted by cyclodextrins of a high molecular weight polycyclic aromatic hydrocarbon benzo[a]pyrene. *J. Hazard. Mater.* 2009; 168: 1296-1301.
- Venny, Gan S, Ng HK. Current status and prospects of Fenton oxidation for the decontamination of persistent organic pollutants (POPs) in soils. *Chem. Eng. J.* 2012a; 213: 295-317.
- Venny, Gan S, Ng HK. Inorganic chelated modified-Fenton treatment of polycyclic aromatic hydrocarbon (PAH)-contaminated soils. *Chem. Eng. J.* 2012b; 180: 1-8.
- Venny, Gan S, Ng HK. Modified Fenton oxidation of polycyclic aromatic hydrocarbon (PAH)-contaminated soils and the potential of bioremediation as post-treatment. *Sci. Total Environ.* 2012c; 419: 240-249.
- Venny, Gan S, Ng H. Evaluation of in situ catalysed hydrogen peroxide propagation (CHP) for phenanthrene and fluoranthene removals from soil and its associated impacts on soil functionality. *Environ. Sci. Pollut. Res.* 2014; 21: 2888-2897.
- Viglianti C, Hanna K, de Brauer C, Germain P. Removal of polycyclic aromatic hydrocarbons from aged-contaminated soil using cyclodextrins: Experimental study. *Environ. Pollut.* 2006; 140: 427-435.
- Watts RJ, Bottenberg BC, Hess TF, Jensen MD, Teel AL. Role of reductants in the enhanced desorption and transformation of chloroaliphatic compounds by modified Fenton's reactions. *Environ. Sci. Technol.* 1999; 33: 3432-3437.
- Watts RJ, Stanton PC, Howsawkung J, Teel AL. Mineralization of a sorbed polycyclic aromatic hydrocarbon in two soils using catalyzed hydrogen peroxide. *Water Res.* 2002; 36: 4283-4292.

- 889 Xue X, Hanna K, Abdelmoula M, Deng N. Adsorption and oxidation of PCP on the surface
890 of magnetite: Kinetic experiments and spectroscopic investigations. *Appl. Catal. B,*
891 *Environ.* 2009a; 89: 432-440.
- 892 Xue X, Hanna K, Despas C, Wu F, Deng N. Effect of chelating agent on the oxidation rate of
893 PCP in the magnetite/H₂O₂ system at neutral pH. *J. Mol. Catal. A: Chem.* 2009b;
894 311: 29-35.
- 895 Yang Y, Ratté D, Smets BF, Pignatello JJ, Grasso D. Mobilization of soil organic matter by
896 complexing agents and implications for polycyclic aromatic hydrocarbon desorption.
897 *Chemosphere* 2001; 43: 1013-1021.
- 898 Yap C, Gan S, Ng H. Feasibility of treating aged polycyclic aromatic hydrocarbons (PAHs)-
899 contaminated soils using ethyl lactate-based Fenton treatment via parametric and
900 kinetic studies. *Environ. Sci. Pollut. Res.* 2015; 22: 329-342.
- 901 Yap CL, Gan S, Ng HK. Fenton based remediation of polycyclic aromatic hydrocarbons-
902 contaminated soils. *Chemosphere* 2011; 83: 1414-1430.
- 903 Yap CL, Gan S, Ng HK. Ethyl lactate-Fenton treatment of soil highly contaminated with
904 polycyclic aromatic hydrocarbons (PAHs). *Chem. Eng. J.* 2012; 200–202: 247-256.

905

906

Figure captions

Figure 1: Different phases of experimental setup used in laboratory to investigate application of Fenton oxidation to remediate PAHs in contaminated soils.

Figure 2. (a) Removal of phenanthrene (PHE) and fluoranthene (FLUT) by using modified-Fenton oxidation in the presence of inorganic chelating agent- SP (sodium pyrophosphate) and organic chelating agents including EDTA (ethylenediaminetetraacetic acid), OA (oxalic acid), SC (sodium citrate) and MA (malic acid). **(b)** H_2O_2 residual for MF treatment in the presence of inorganic (SP) and organic chelating agents (EDTA, OA, SC and MA). Adapted with permission from Figure 2 and Figure 5 of Venny et al. (2012b).

Figure 3: PAHs degradation in (a) organic extracts (spiked on sand) and (b) soil pretreated with availability enhancement agent during oxidation experiments by: H_2O_2 alone without iron activation (■-HP), H_2O_2 + soluble Fe^{II} (○-F) and H_2O_2 + magnetite (●-FL). Blank (▲) experiment was performed in the presence of magnetite but without any oxidant. This degradation is represented in terms of C_t/C_0 where C_t is the sum of 16 PAHs concentration at specified oxidation time and C_0 is their concentration at $t=0$ (before oxidation) measured by GC-MS. Adapted with permission from Fig. 2 and Fig. 6 of Usman et al. (2012). Experimental conditions were: solid matrix = 2 g, volume of solution = 20 mL, magnetite (10%, w/w) and equivalent amount of Fe^{II} .

Figure 4: Efficiency of ethanol in comparison with other PAH-availability enhancement agent to assist Fenton oxidation where F: traditional Fenton, ETF: ethanol-Fenton system, ELF = ethyl lactate-Fenton system, SDS-F = sodium dodecyl sulfate- Fenton system, CD-F = cyclodextrin-Fenton system. Instead of traditional Fenton oxidation, Usman and co-workers (2012) applied magnetite catalyzed Fenton-like oxidation (FL) assisted by ET and CD.

Figure 5: Degradation of 16 PAHs in former coking plant soil by Fenton-like oxidation assisted by various pretreatments and is represented by C/C_0 where C is the final concentration of PAHs after treatment and C_0 is the initial concentration. Where initial = PAH concentration before treatment and others represent application of Fenton-like oxidation direct (Direct oxidation = when soils were subjected to chemical oxidation without any pretreatment), spiking/FL (when EOM was extracted from soils and spiked on sand), Ext-Pt/FL (extraction pretreatment to incorporate PAHs as fresh contaminants in soil as detailed in Section 2.2.2), and FL oxidation assisted by ethanol (ET-FL) or cyclodextrin (CD) or by thermal pretreatment at 60, 100, and 150 °C.

Figure 6. Density of cultivable bacteria, 16S rDNA, PAH-RDH α and 18s rDNA gene copy numbers in oxidized by Fenton treatment at low (L) i.e. 6 g kg⁻¹ and high (H) i.e. 65 g kg⁻¹ concentration of H₂O₂ and control soil (Control) at the start (T_0) and after 5 weeks (T_5) of incubation. As the evolution of PAH-RDH α GN and GP populations was not significantly different, the two populations were added to represent the global population of degraders. Mean $\pm$ SE ($n = 3$). LD: detection limit. Different letters within category data indicate significant differences between treatments; stars indicate significant differences between T_0 and T_6 by Mann and Whitney test. Reproduced with permission from Fig. 2 of Laurent et al. (2012).

950 **Table 1:** Research studies using Fenton oxidation assisted by various kinds of chelating agents.

Focus of study	Soil	Tested chelating agents (CA)	Experimental conditions	Important results	Reference
To compare the efficiency of traditional Fenton and modified Fenton to couple with biodegradation for enhanced PAH removal	Spiked soil (PAHs = 600 mg/Kg), silt loam and soil sampled from former manufactured gas plant site, loamy sand	Catechol (CA) and gallic acid (GA)	5 g of MGP soil in 20 ml of distilled water, Fe ^{III} , 82.5 mg of catechol or 141.0 mg of gallic acid and Fe(ClO ₄) ₃ ·6H ₂ O (0.07 M), 2 g of H ₂ O ₂ , neutral pH For traditional Fenton, experiments were performed at pH = 2-3 without chelating agents	Traditional Fenton was effective but required pH (pH 2–3) for optimum efficiency made the process incompatible with biological treatment. Efficiency was maintained by using chelating agents that allowed coupling with bioremediation.	Nam et al., 2001
Production of surfactants during various chemical oxidation strategies for PAH remediation	Soil sampled from former manufactured gas plant site, having sand:silt:clay =79:20:1%	EDTA	Laboratory slurry reactors, 4 kg of soil (dry weight) and 8.5 L of water (including the oxidant solutions), 200 mg of EDTA, 1 g Fe ^{III} (as ferric sulfate), and 1 L of 50% H ₂ O ₂ , pH=8	All tested treatments resulted significant surfactant production and overall removal of PAHs. Modified Fenton with calcium peroxide exhibited better efficiency.	Gryzenia et al., 2009
Effect of various chelating agents on the efficiency of Fenton oxidation to remove PAHs	Spiked soil (loamy sand), containing 1000 mg/kg of two PAHs	Sodium pyrophosphate (SP) as organic CA, EDTA, Oxalic acid (OA),	Batch slurry system, 5 g of soil, 480 mM H ₂ O ₂ , 60 mM Fe ³⁺ , solution:soil = 15 mL:5 g. Different doses of CAs: 0.1135–0.4015 g	Inorganic SP was demonstrated to be superior in efficiency to the tested organic CAs.	Venny et al., 2012c

		sodium citrate (SC) as inorganic CAs			
Optimization of experimental conditions for efficient coupling of Fenton's reagent with bioremediation	Soil (loamy sand) spiked with two PAHs, 500 mg the PAH/kg for each PAH,	SP	Unsaturated soil columns, 1.2 Kg soil, Various doses of oxidant, SP, Fe ^{III} at different reaction times with soil pH of 4.07–5.41	An optimized operating condition of the modified Fenton was found at H ₂ O ₂ /soil 0.05, Fe ^{III} /soil 0.025, SP/soil 0.04 and 3 h reaction time with 79.42% and 68.08% removal of tested PAHs. Use of SP assisted Fenton treatment and bioremediation serves as a suitable strategy to enhance soil quality.	Venny et al., 2012b
Role of various chelating agents to improve operating range of pH for degradation of PAHs by modified Fenton oxidation using iron nano particles	Spiked soil (silty sand), containing 100 mg of pyrene per kg of soil	SP as inorganic CA and EDTA, FA, HA and SC (as organic CAs)	Batch experiments, 5 g soil, 15 mL distilled water, iron nano-oxide = 5 – 60 mM, H ₂ O ₂ = 0-500 mM, pH = 3 and 7	Application of modified Fenton treatment, with the following conditions (300 mM of H ₂ O ₂ , 30 mM of iron nano oxide, SP as CA, pH 7 after 6 h of reaction, removed 93% of tested PAHs that was higher than organic CAs.	Jorfi et al., 2013

952 **Figure 1:**

953

954

Figure 2:

959 **Figure 3:**

960

961 **Figure 4:**

962

963 **Figure 5:**

964

965 **Figure 6:**