

Aggressiveness Changes in Populations of *Didymella pinodes* over Winter and Spring Pea Cropping Seasons

G. Laloi, J. Montarry, M. Guibert, D. Andrivon, Didier Michot, Christophe
Le May

► **To cite this version:**

G. Laloi, J. Montarry, M. Guibert, D. Andrivon, Didier Michot, et al.. Aggressiveness Changes in Populations of *Didymella pinodes* over Winter and Spring Pea Cropping Seasons. *Applied and Environmental Microbiology*, American Society for Microbiology, 2016, 82 (14), pp.4330 - 4339. 10.1128/AEM.00480-16 . hal-01400496

HAL Id: hal-01400496

<https://hal-univ-rennes1.archives-ouvertes.fr/hal-01400496>

Submitted on 22 Nov 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Aggressiveness changes over time in populations of *Didymella pinodes* over**
2 **winter and spring pea cropping seasons**

3

4 Laloi G.^{1,4}, Montarry J.¹, Guibert M.¹, Andrivon D.¹, Michot D.^{2,3}, Le May C.^{1,2,*}

5 ¹ INRA, UMR1349 Institut de Génétique, Environnement et Protection des Plantes (IGEPP), Domaine de la
6 Motte, 35653 Le Rheu, France.

7 ² Agrocampus-Ouest, Laboratoire Ecologie et Santé des Plantes (ESP), 65 rue de Saint Briec, 35042 Rennes,
8 France.

9 ³ INRA, UMR1069 Sol Agro et Hydrosystème Spatialisation (SAS), 65 rue de Saint Briec, 35042 Rennes, France.

10 ⁴Present address : INRA, UMR1345 Institut de Recherche en Horticulture et Semences (IRHS), 42 rue Georges
11 Morel BP60057, 49071 Beaucozé France.

12

13 *** Corresponding author**

14 Christophe Le May

15 E-mail: lemay@agrocampus-ouest.fr

16

17 **Words in the text:** 5650

18 **Words in the abstract:** 173

19 **Number of references:** 76

20 **Number of figures:** 5

21 **Number of tables:** 3

22

23

24

25 **ABSTRACT**

26 Ascochyta blight, caused by the necrotrophic ascomycete *Didymella pinodes*, is responsible for severe
27 losses in winter and spring pea crops. Despite different climatic conditions, epidemics on winter and
28 spring crops are due to a single population of *D. pinodes*, suggesting gene flow either between the
29 two crops or from reservoir sources during the cropping season. This should lead to similar
30 pathogenicity characteristics in isolates sampled from both crops. However, these hypotheses have
31 never been formally tested. We thus sampled a total of 520 *D. pinodes* strains throughout a growing
32 season from winter and spring pea plots (WP and SP, respectively) and from trap plants (TWP and
33 TSP). AFLP markers revealed a high genetic diversity within sub-populations, whereas pathogenicity
34 tests showed that mean aggressiveness increases over the course of epidemics. These results support
35 the idea that allo-inoculum contributes to the carry-over of epidemics between winter and spring
36 crops, and that the most aggressive isolates are selected as the epidemics progress.

37

38 **IMPORTANCE**

39 *Ascochyta blight, caused *Didymella pinodes*, is responsible for severe losses in pea crops. While*
40 *previous study have showed that Ascochyta blight epidemics on winter and spring crops are due to a*
41 *single population of *D. pinodes*, suggesting that isolates from both crops present similar pathogenicity*
42 *characteristics, that hypothesis have never been tested. Genetic analysis of sub-populations sampled*
43 *throughout a growing season from winter and spring pea plots revealed a high genetic diversity*
44 *within sub-populations, whereas pathogenicity tests showed that mean aggressiveness increases*
45 *over the course of epidemics.*

46

47 **Keywords:** Ascochyta blight, genetic structure, gene flow, *Didymella pinodes*, AFLP, pea,
48 aggressiveness

49

50 **Introduction**

51 The spatio-temporal dynamics of crop diseases are simultaneously impacted by pathogens, host
52 plants, environment and human activities (**1, 2**). Indeed, whether or not hosts and pathogens interact
53 is largely determined by spatial and temporal components of host and pathogen life-history traits
54 largely (**3, 4**). These interactions can thus be conceptualized as a continuous sequence of biological
55 cycles including dormancy, growth, reproduction, dispersal, and pathogenesis (**5**).

56 Gene flow, resulting from pathogen reproduction and dispersal, can drastically increase the ex-
57 tent to which pathogen epidemics spread across a landscape (**6**). As such, it is therefore a main factor
58 in the transmission of disease to previously uninfected areas, and drives the spatial structure of
59 pathogen populations in fragmented landscapes by influencing the long-term survival and genetic
60 composition of populations (**7, 8, and 9**). Individual dispersal events, occurring over periods of days
61 or weeks during both the cropping and intercropping seasons (**2**), originate from a large number of
62 potential inoculum sources: resting structures in soil (mycelium, oospores, chlamydozoospores or sclero-
63 tia), infested stubble left on the soil surface, infested seed, and alternative hosts (wild or cultivated
64 plants, including volunteers). The degree of connectivity among host populations is thus likely to
65 influence spatial patterns of disease occurrence and persistence (**10, 11, 12, 13, and 14**). Dispersal
66 events also shape the structure and changes of population genetic variability. The role of gene flow
67 within and among plant pathogen populations is still insufficiently characterized, but is crucial to
68 understand the distribution of alleles conferring virulence or fungicide resistance within populations.
69 Disease dispersal can favor the contact between wild-cultivated or cultivated-cultivated areas during
70 the cropping and the intercropping season, and thus gene flow between these compartments can
71 influence the genetic structure of populations. By mixing populations submitted initially to different
72 selection pressures, it can also negate the effect of local selection on adaptation and result in local or
73 general maladaptation (**15**).

74 Reproduction is the other important factor which impacts the genetic structure of plant patho-
75 gens (**16, 17**). The life cycle of many fungal plant pathogens alternates asexual multiplication with

76 episodes of sexual reproduction, but their relative importance varies both within and among species
77 (18). Asexual multiplication rapidly increases the size of populations. Strains with the best reproduc-
78 tive success are amplified and tend to decrease the overall population genotypic diversity in time (19,
79 9). For instance, in the case of the grapevine powdery mildew, a positive relationship between spatial
80 and genetic distances shows that epidemics result from the spread of clones within a crop (20). Be-
81 sides, the genotypic diversity may be increased with a genetic mixing by sexual reproduction (21, 18).
82 For the potato late blight pathogen *Phytophthora infestans*, clonal reproduction has been reported in
83 populations with restricted levels of genotypic diversity from UK (22) and France (23, 15), whereas
84 sexual reproduction has been reported in populations with high genotypic diversity from Nordic
85 countries (21, 24).

86 Despite recent advances, the role of gene flow (and of its two mains triggers, reproduction and
87 dispersal) on the local evolution of pathogen populations is still poorly assessed. This is especially the
88 case for epidemics occurring on both winter and spring genotypes of the same host. This situation
89 raises specific questions, particularly regarding the respective roles of auto- and allo-inoculum ([Inocu-
90 lum fraction produced by an exogenous source to the plot \(25\)](#)) in the epidemic process and in the
91 pathogenicity changes within populations. To address these questions, we analyzed the phenotypic
92 and genotypic diversity in populations of an aggressive foliar pathogen of pea, *Didymella pinodes*
93 along the course of an epidemic. *D. pinodes* is a necrotrophic, polyphagous, polycyclic and homothal-
94 lic fungus, that causes Ascochyta blight on winter and spring pea fields worldwide (26, 27). This path-
95 ogen is known to present individuals varying widely in their ability to cause disease on pea (28, 29,
96 30, 31, and 32). Contrary to many necrotrophic fungi, *D. pinodes* can simultaneously develop its ana-
97 morph (asexual) and teleomorph (sexual) forms on the same plant during the growing season (33,
98 32). Moreover, due to the indeterminate growth of the pea crop, both pycnidia (asexual) and
99 pseudothecia (sexual) can be observed simultaneously on the same plant organs. Whereas pycnidia
100 are produced on both green and senescent plant organs, pseudothecia only appear on the senescent
101 parts (33). This fungal pathogen has different ways to persist in the environment and to disperse over

102 more or less long distances (splashing, wind) (27, 34, 35). These features are more important than for
103 pathogens where reproduction modes are dissociated, as they help to increase the population diver-
104 sity and have an impact on pathogenicity. A recent study (32) showed that epidemics on winter and
105 spring pea crops are due to a single *D. pinodes* population. Despite the long intercropping season, *D.*
106 *pinodes* populations from winter and spring pea display a high, but similar genetic variability. Contra-
107 ry to other natural plant–pathogen associations, no high annual extinction is observed (36, 37, 38,
108 39, and 40). Population turnover is thus likely to be driven by both local selection and demographic
109 stochasticity due to seasonal population growth and decline. During spring, abundant spore produc-
110 tion (41) leads to high colonization rates. The fact that winter and spring pea crops are attacked by
111 the same populations of *D. pinodes* suggests that, as observed by Laine & Hanski (42) in another as-
112 comycete plant pathogen, regional persistence of *D. pinodes* populations could occur at the scale of
113 metapopulations, consisting of many coupled populations.

114 The origin and the availability of inoculum sources during the cropping season are still
115 questionable. Do *D. pinodes* populations colonizing winter pea crops constitute the only inoculum
116 sources for the spring pea crops? Does their apparent genetic similarity translate into similar
117 pathogenicity levels? This study, designed to address these questions, is thus based on a collection of
118 *D. pinodes* isolates sampled throughout a single cropping season (January to June) on both winter
119 pea (WP) and spring pea (SP) crops, but also on trap plants (TWP and TSP) catching the allo-inoculum.
120 The specific objectives of this work were 1) to determine if and how *D. pinodes* aggressiveness evolves
121 during the cropping season, 2) to define if *D. pinodes* populations developed on winter pea crops
122 constitute the main inoculum source for spring pea crops, and 3) to and analyse the relationships
123 between crop populations and allo-inoculum caught on trap plants.

124

125

126

127 **Materials and methods**

128 **Field experiments.** Two field experiments were carried out at the INRA experimental station of Le
129 Rheu, western France (48°06 00 N, 1°48 00 W, 30m above mean sea level) in 2004-2005. Both fields,
130 separated by more than three km, shared similar pedoclimatic environments and were free from soil
131 borne inoculum, due to a rotation without pea crop during the five previous years. These fields were
132 respectively sown with a winter (Cheyenne, GAE-Recherche, France) and a spring pea cultivar
133 (Baccara, Florimond-Desprez, France). The winter cultivar was sown on October 25th, 2004, and the
134 spring crop was sown on February 24th, 2005. For both crops, the experimental design consisted of a
135 30m² (3m wide x 10m long) plot divided into thirty micro-plots (1 m² each).

136

137 **Trap plants.** Allo-inoculum, defined as the [inoculum fraction produced by an exogenous source to the](#)
138 [plot \(25\)](#) was assessed through trap plants. Each week from mid-January to the end of June 2005, one
139 tray containing 20 trap plants (Cheyenne or Baccara seedlings at the 5-leaf stage, respectively for the
140 winter and spring trials) was placed at each of the four corners of the plots. After seven days
141 exposure in the field, trap trays were brought back and incubated in a dew chamber (12h-
142 photoperiod, 20°C night/day, 100% relative humidity) for four days. Deposition of viable ascospores
143 on trap plants was assessed by isolating strains from the small, purple-black, irregular flecks on the
144 five lower stipules of the plants after incubation (41).

145

146 **Spatio-temporal disease development.** [Disease started](#) in each crop without artificial supply of
147 inoculum. Disease severity was assessed visually once a month in the different micro-plots from
148 January to June. To determine disease development at the plot level, one plant was randomly
149 collected from each micro-plot at the different sampling dates (five sampling dates for the winter pea
150 crop, and three sampling dates for the spring pea crop). Disease was scored on each stipule of the
151 sampled plants using the adapted disease scale of Tivoli (1994) (41), and a mean disease score (D_i)
152 was calculated. Disease distribution maps were created for each sampling date using the Arcmap®

153 software and a deterministic method based on the estimation of the weighted inverse distance (43).

154

155 **Fungal strains.** A total of 520 strains of *D. pinodes* collected from infected field pea plants and trap
156 plants were used in this study. Each field was divided into 30 micro-plots where one plant was
157 randomly sampled at each sampling date. One hundred and fifty strains were sampled from winter
158 pea (WP) during the winter cropping season (February to May). Ninety strains were sampled from
159 spring pea (SP) during the spring cropping season (May and June). Two hundred and fifteen strains
160 were sampled from trap plants placed at the corner of the winter pea plots (TWP), and sixty-six
161 strains were sampled from trap plants placed at the corner of the spring pea plots (TSP). For each
162 population (WP, SP, TWP and TSP), the different sub-populations correspond to the different sampling
163 dates (Table I). All the strains were cultured and single-spored before being studied. For isolation,
164 approximately 5 mm² of diseased leaf tissue was surface-sterilized for 1 min in 70% ethanol, rinsed
165 three times in sterile water, placed on sterile filter paper to remove excess water, and plated for 14
166 days on V8 medium (99 mL V8 vegetable juice [Campbell, France], 35 g agar, and 801 mL distilled
167 water, autoclaved at 105°C for 30 min) distributed in Petri dishes. Pycnidiospores from resulting
168 cultures were spread on 2% malt agar and incubated for 12 h as described by Onfroy *et al.* (44). Single
169 germinating pycnidiospores were transferred to fresh V8 plates under a dissecting microscope, and
170 cultures were incubated at 20°C with a 12 h photoperiod under cool white fluorescent lamps. Single-
171 spore cultures were then maintained on malt slants and stored in the dark at 4°C.

172

173 **DNA extraction and AFLP typing.** Each strain was grown in 75 mL of LP liquid medium (10 g tryptone,
174 5 g extract of yeast powder, 5 g NaCl, 1 L distilled water; autoclaved at 115°C for 20 min)
175 supplemented with streptomycin (1.5 g) and penicillin (0.75 g) from four pieces (approximately 1 cm²
176 each) cut from the margin of an actively growing culture on malt agar. Inoculated vials were
177 incubated, under agitation, for 14 days at 20°C with a 12 h photoperiod under cool white fluorescent
178 lamps. Mycelia were harvested by vacuum filtration through two layers of sterilized Miracloth

179 (Calbiochem CN Biosciences, Inc., La Jolla, CA), rinsed twice in sterile water, and stored at -80°C until
180 lyophilized. DNA was extracted from lyophilized mycelium as described by Lodhi *et al.* (45), quantified
181 by measuring the optical density of extracts at 260 and 280 nm with a Nanodrop 1000
182 spectrophotometer (Thermo Scientific), and adjusted to a final concentration of 100 ng.L⁻¹ for
183 amplified fragment length polymorphism (AFLP) analysis. AFLP analysis was carried out as described
184 by Vos *et al.* (46) with modifications used by Le May *et al.* (2012). AFLP reactions were performed
185 independently three times, using the same set of primers with reference strains and a random
186 sample of 10 isolates from the collection, and independent DNA preparations of the same strains to
187 estimate the repeatability of fragment scoring.

188

189 **AFLP analysis.** The raw data were analyzed with GeneMapper (Version3.5, Applied Biosystem). The
190 presence and absence of all fragments between 100 and 400 bp were scored in each of the 520
191 strains. Bands with molecular sizes exceeding 400 bp were not scored because of insufficient
192 resolution. The dataset obtained was based on the assumption that bands of the same molecular
193 weight were identical.

194

195 **Genotypic diversity.** The AFLP data were used to define multilocus genotypes (MLGs) and check for
196 repeated MLGs, , *i.e.* the strains sharing the same alleles at all loci, using the Microsoft EXCEL add-in
197 GenAEx version 6.5 (47). Genotypic diversity was calculated for each population using the Shannon &
198 Wiener's index H' (48, 49). The number of different alleles (Na), Shannon's information index (I) and
199 the unbiased diversity (uh) were computed using GenAEx for each population and sub-population.
200 Clonality was assessed using the index of association (I_A), a measure of the multilocus linkage
201 disequilibrium, calculated with the Multilocus software version 3.1b (50). [The index of association \(\$I_A\$ \)](#)
202 [is the traditional measure of multilocus linkage disequilibrium. The "distance" \(number of loci at](#)
203 [which they differ\) between all pairs of individuals is calculated, and the variance of these distances](#)
204 [compared to that expected if there is no linkage disequilibrium. Index of association is calculated](#)

205 using the following formula:

$$206 \quad I_A = \frac{V_D}{\sum var_j} - 1$$

207 where V_D is the variance of the distances two isolates over all loci (i.e., the number of loci at which
208 they differ), and var_j corresponds to the variance of the mean distance (either 0 or 1) between all
209 $n(n-1)/2$ possible pairs of isolates. The higher the I_A , the more clonal the population. Departure from
210 the null hypothesis, i.e. complete panmixia, was checked by permuting alleles between individuals
211 independently for each locus (500 permutations).

212

213 **Population differentiation.** The analysis of molecular variance (AMOVA; online as Arlequin version
214 3.1 software, hosted by the Department of Anthropology, University of Geneva, Switzerland; (51) was
215 used to partition molecular variance between populations, between sub-populations within
216 populations and within sub-populations. Pairwise F_{ST} values were calculated with Arlequin for each
217 pair of sub-populations of *D. pinodes* and compared within plots (SP-SP and WP-WP) or trap plants
218 (TSP-TSP and TWP-TWP), between plots (SP-WP) and between plots and related trap plants (SP-TSP
219 and WP-TWP).

220

221 **Genetic structure.** A principal component analysis (PCA) was performed using the procedure available
222 in the package adegenet (52) for the statistical freeware R version 3.1.1 (©2014, The R Foundation for
223 Statistical Computing). PCA has an important advantage over other methods, such as the Bayesian
224 clustering algorithm implemented in STRUCTURE (53), because it does not require strong
225 assumptions about an underlying genetic model, such as the Hardy-Weinberg equilibrium or the
226 absence of linkage disequilibrium between loci (52).

227

228 **Aggressiveness of *D. pinodes* strains.** Aggressiveness level was evaluated for a random sample of *D.*
229 *pinodes* strains sampled on winter (sampling dates: WP07, WP10, WP12, WP14) and spring (sampling
230 dates: SP01, SP03) pea plots, and of *D. pinodes* strains sampled on winter trap (sampling dates:

231 TWP20, TWP24, TWP30, TWP34) and spring trap plants (sampling dates: TSP02, TSP03, TSP06). Five
232 *D. pinodes* strains were randomly chosen within each sub-population and their aggressiveness was
233 evaluated on three pea genotypes: the winter cultivar Enduro (Florimond-Desprez, France), the
234 spring cultivar Lumina (Nickerson, France), and the spring breeding line DP (54). Enduro and Lumina
235 were chosen to replace Cheyenne and Baccara, respectively, for which seeds were no longer available
236 at the time of the pathogenicity tests. They presented a similar and high susceptibility level towards
237 the disease, while DP has a higher level of quantitative resistance. Plants were grown at 18 to 20°C
238 for three weeks, until they reached the 5- to 6-leaf stage, before inoculation. Plant preparation and
239 experimental design were as described by (44). The inoculation method was based on that proposed
240 by Onfroy *et al.* (56). Briefly, strains were grown for 10 days on V8 medium under white light with a
241 12 h photoperiod at 20°C before pycnidiospores suspensions were prepared by flooding the surface
242 of cultures with sterile distilled water, gently scraping with a glass rod, and filtering the suspension
243 through two layers of sterile cheesecloth. The spore concentration was adjusted to 5.10^4 spores.mL⁻¹,
244 and Tween 20 (VWR International SAS, Strasbourg, France) was added as a wetting agent (2 drops per
245 500 mL spore suspension). Inoculation consisted of depositing a 10 µL drop of the spore suspension
246 on the upper surface of freshly detached stipules floated, lower surface down, on tap water in a
247 compartmentalized square Petri dish (12 cm side; Gosselin, France). Drops were deposited away from
248 the main veins. To avoid drop evaporation, Petri dishes were placed into large transparent plastic
249 boxes immediately after inoculum deposition and incubated in a climate chamber for seven days with
250 a continuous cycle of 14 h of light at 20°C. Symptom development was assessed two, four, and seven
251 days after inoculation, as described by Onfroy *et al.* (55). A 0-3 semi-quantitative scale (0 = symptom
252 free, 1=flecks appearing, 2 = flecks covering half of the area of drop deposition, 3 = coalescence of the
253 flecks within the area of drop deposition) was used to score symptoms not extending past the
254 inoculation droplet. For stipules with necrosis extending beyond the borders of inoculum drops,
255 lesion diameter (mm) was measured with a graduated ruler. A visual assessment using a 0-7 scale
256 adapted from Wroth (1998), where 0 = symptom free, 1 = flecks appearing, 2 = flecks covering half of

257 the drop deposit, 3 = coalescence of the flecks in the area of the drop deposit, 4 = 3-to-6mm lesion
258 diameter, 5 = 6-to-9mm lesion diameter, 6 = 9-to-12mm lesion diameter, 7= >12mm diameter, was
259 also performed. Two stipules from each of four different plants per genotype were inoculated. For
260 each genotype, the area under the disease progress curve (AUDPC) was calculated as

$$AUDPC_i = \sum_{j=1}^m \frac{(D_{i,j} + D_{i,j+1})}{2} \times (t_{j+1} - t_j)$$

261 where $D_{i,j}$ and $D_{i,j+1}$ correspond to disease scores at two consecutive dates, t_j and t_{j+1} (56). Statistical
262 analysis of the data was performed with the R statistical software, version 3.1.1 (©2014, The R
263 Foundation for Statistical Computing). Normality and homogeneity of variances were checked by the
264 Shapiro-Wilk and the Leven tests, respectively. The effects of population (WP, SP, TWP and TSP), pea
265 genotype (DP, Lumina and Enduro) and their interaction on AUDPC values were tested through a
266 multi-way ANOVA. The effects of sub-populations, pea genotypes and their interaction were tested
267 on AUDPC values using a second ANOVA model. When significant effects were detected, mean values
268 were compared with Tukey tests (alpha = 0.05).

269

270 Results

271 *D. pinodes* displayed a homogeneous spatial development during the cropping season

272 Disease layout maps in winter and spring pea crops, as determined with the Arcmap® software, were
273 spatially similar for each set of maps in both pea crops (Fig. 1). Disease severity increased with time,
274 but was spatially homogeneously distributed through the plots, with no strong foci. The mean disease
275 scores were higher in the winter pea crop than in the spring pea crop, essentially because of the
276 longer growing season and the more conducive climatic conditions in the late winter and early spring.

277

278

279 **Genotypic flow acts on the genetic structure of *D. pinodes* population during the cropping**
280 **season**

281 The three AFLP primer sets allowed the detection of 646 loci, 388 of them were polymorphic over all
282 tested populations. The percentage of polymorphic markers in each sub-population ranged from
283 29.90% in TWP21 to 100% in WP8. No repeated MLG was found among the 520 strains, indicating a
284 very low incidence of clonality. However, the overall I_A value significantly differed from zero ($I_A =$
285 1.97), thus rejecting the null hypothesis of recombination ($P = 0.001$) (**Fig. 2**). Number of alleles (N_a),
286 Shannon's Information index (I) and unbiased diversity (uh) showed that the highest genetic diversity
287 within sub-populations was for WP12 ($I=0.61$, $uh=0.43$) and the least for TWP21 ($I=0.176$, $uh=0.151$).
288 The low genetic diversity observed for the sub-population TWP21 may be due to the low number of
289 isolates within this sub-population. At the scale of the populations (WP, SP, TWP and TSP), number of
290 alleles, Shannon's Information index and unbiased diversity showed that the genetic diversity was
291 higher for populations sampled into the plots (SP and WP) than for populations sampled on trap
292 plants (TWP and TSP) (**Table I**).

293 AMOVA revealed that 80.35 % of the total genetic variance was partitioned within sub-
294 populations (**Table II**). A relatively low proportion of genetic variability was attributable to differences
295 between populations (9.62 %) and between sub-populations within populations (10.03 %). High
296 variances of F_{ST} values were detected, ranging from -0.055 to 0.586. Pairwise F_{ST} were higher between
297 plots and trap plants (WP-TWP and SP-TSP) than within plots (WP-WP or SP-SP) (**Fig. 3**), quite low
298 between plots (SP-WP) and within spring trap plants (TSP-TSP), but quite high within winter trap
299 plants (TWP-TWP).

300 Principal Component Analysis (PCA) failed to separate into different groups the strains of the
301 sub-populations sampled on WP, SP, TWP and TSP (**Fig. 4**). Moreover, the low percentage of genetic
302 diversity explained by the two principal axis of the PCA (11% and 9.95%, respectively) suggested that
303 these different sub-populations did not constitute distinct genetic groups and that all the strains
304 belonged to a single population.

305 **Aggressiveness changes during the cropping season differed between winter and spring**
306 **pea populations**

307 The aggressiveness level differed according to test host genotypes (pea genotype effect, **Table III**): as
308 expected, AUDPC values were significantly lower for the resistant pea genotype DP than for the pea
309 genotype Enduro, and intermediate (and not significantly different from the two other genotype) for
310 the pea genotype Lumina. However, the lack of significant interaction between populations or sub-
311 populations and pea genotypes (**Table III**) showed that all populations shared the same ranking
312 across hosts.

313 Overall, strains sampled on winter plots (WP) and on trap plants (TWP and TSP) were significantly
314 more aggressive than strains sampled on spring plots (SP) (**Fig. 5A**). Moreover, at the sub-population
315 level, the aggressiveness of SP strains increased strongly during the growing season, whereas it
316 remained stable over time in the other three populations (**Fig. 5B**).

317 Allo-inoculum sampled on TWP or TSP showed a similar aggressiveness level than strains sampled
318 from the winter plot (**Fig. 5A, 4B**). Altogether, these results tended to show that trap plants were
319 always infected from the winter populations, and that the spring plots initially select against
320 aggressive isolates.

321

322 **Discussion**

323 One of the objectives of this study was to determine how *D. pinodes* population structures change
324 during a growing season. The amount of genetic variation was similar within *D. pinodes* sub-
325 populations sampled from winter or spring pea plots ($H' = 0.83$, and 0.86 respectively). The genetic
326 variability observed at the beginning of the season remained roughly the same throughout the
327 growing season. The main part of this genetic variability was observed within sub-populations (80%),
328 which confirms the PCA result of an absence of a clear genetic structure over time, and suggests that
329 significant gene flow occurs between sub-populations. The low pairwise F_{ST} values observed within

330 (SP-SP and WP-WP) and between plots (SP-WP) suggest also that sub-populations were not
331 genetically differentiated. Previous works released on *D. pinodes*/pea pathosystems showed that *D.*
332 *pinodes* populations from Canada, France, and Algeria displayed a high degree of genetic variability
333 (32, 57). This variation estimated to a different scale than in our study, is illustrated by the high value
334 of Shannon's information index and Nei's gene diversity, and by the high number of distinct
335 haplotypes. Our study showed that population diversity estimated at a field level, and during the
336 growing season is changing, but slightly higher to the diversity estimated among countries or
337 locations within a country. Indeed, H' value in our study ranged from 0.70 to 0.83, whereas it ranged
338 from 0.39 to 0.47 to *D. pinodes* populations collected from different locations in Algeria (57).
339 Moreover, H' value was closely similar to those obtained for fungi showing great genetic diversity, as
340 in *A. rabiei* [$H'=0.58$ (58)] *Mycosphaerella fijiensis* [$H'=0.58$ (59)] or *Botrytis cinerea* [H' ranging from
341 0.401 to 0.518 (60)]. This change of diversity level observed during the growing season reinforced the
342 fact that sexual reproduction was important in the life cycle of *D. pinodes* and the predominance of
343 this mode of reproduction seemed to affect earlier the population structure.

344 On trap plants, pairwise F_{ST} were higher within winter than within spring sub-populations,
345 suggesting that the same allo-inoculum sources must be active during all the whole spring growing
346 season, whereas other allo-inoculum sources must be active during the winter cropping season. This
347 result suggests that epidemics on winter crops are initiated from a large number of sources producing
348 pseudothecia and releasing ascospores, whereas the release of ascospores, probably from the
349 infected winter crops, fuel the spring crop epidemics with a steadier and genetically similar source of
350 inoculum. It also suggests that asexual inoculum is not an important component of epidemic
351 extension, although it might play a major role in local disease intensification. The spatially
352 homogeneous disease severity observed throughout the winter and spring plots sampled here also
353 point to an homogeneous distribution of initial inoculum consistent with ascospore showers from
354 external sources rather than with short distance splashes of asexual pycniospores. If ascospore
355 release indeed constitute the main mechanism fueling the epidemic development and spread, it

356 would provide also a good explanation for the connected genetic structures of pathogen populations
357 sampled in different plots. As indicated in the material and methods section, our experiment was
358 realized only once, but we have assessed disease development and genetic diversity evolution on two
359 different crops. Past study showed that *D. pinodes* disease developed on winter and spring pea was
360 initiated by a single population, whose pathogenicity is a plastic trait modulated by the physiological
361 status of the host plant. The fact that only a single year was conducted suggests different limitations.
362 A second year of experiment will help us to see if a similar pool of allo-inoculum would be mobilized
363 during the cropping season. Indeed a second year of field experiments exposed us to different
364 constraints: *i)* climatic change that will modify plant development, plant receptivity, availability of
365 different sources of inoculum, *ii)* a second year of field experiment will also suggest that diversity of
366 agrosystem will change (change in prevalence of legume species, diversity of cultivated species in a
367 neighboring environment). As described by Plantegenest *et al.* (62) the composition of landscape
368 determines the local abundance of potential reservoirs of inoculum that may obviously influence the
369 global propagule pressure and hence the risk of infection of a plant. Those reservoirs may, in
370 particular, consist of diseased individuals of the same host species or of alternative, either cultivated
371 or wild, hosts. Moreover, change in landscape composition and diversity will change dynamics of
372 pathogen.

373 The initial, natural hypothesis is that each type of pea crop would serve as the main inoculum
374 source for the other, via exchanges between plots. However, since ascospores are most likely the
375 main inoculum form starting and fueling the epidemics, and given the large dispersal capabilities of
376 wind-borne ascospores, sources other than plants growing in adjacent or neighboring plots need to
377 be considered as inoculum reservoirs. Indeed, our results suggest that various parts of the metapop-
378 ulation are mobilized during the growing season, particularly during winter growing season. Two
379 further sources of inoculum could be involved: alternative host and chlamydo-spores (35, 61).

380 Compared to infected pea debris, alternative hosts are generally considered of minor importance
381 in the epidemiology of *Ascochyta* blight (63). Recently, Trapero-Casas & Kaiser (64) showed that *A.*

382 *rabiei* is capable of infecting different plant species, suggesting that these naturally infected alterna-
383 tive hosts could serve as significant sources of inoculum to initiate disease epidemics on cultivated
384 chickpea. In a recent work, Le May *et al.* (61) showed that the asexual stage of *D. pinodes* could be
385 observed on pea, common vetch and clover, fuelling the suggestion that other legume species may
386 also act as inoculum sources for epidemics on peas (65, 63, 66, 67). Legume species like vetch or clo-
387 ver generally sympatric to pea crops during the growing season, but also persist during the intercrop-
388 ping season. As shown by Savage *et al.* (6), long distance dispersal generally results in rapid transmis-
389 sion of disease to previously uninfected areas, and can facilitate genetic interaction between spatially
390 separated populations, resulting in the introduction of new virulent alleles into existing populations
391 (68). Thus, a more extensive knowledge of the host range of *D. pinodes* and of the relatedness be-
392 tween populations from peas and other hosts could help to estimate the risk of Ascochyta blight epi-
393 demics of pea arising from alternate hosts, as in other pathosystems where cultivated and wild hosts
394 grow sympatrically (69,70).

395 *D. pinodes* can also survive in soil as mycelium or chlamydospores (71, 72). Davidson *et al.* (35)
396 investigated the survival of ascochyta blight pathogens in soils of commercial pea-cropping paddocks,
397 and showed that the level of pathogen populations in the soil was related to the severity of the epi-
398 demic in the last pea crop grown. Although airborne ascospores of *D. pinodes* appear to be the pri-
399 mary inoculum during establishment of field pea crops, soil borne inoculum has also been associated
400 with disease (35). To evaluate the impact of this inoculum source on the genetic structure of *D. pi-*
401 *nodes* populations, it would be interesting to develop a genetic approach to characterize *D. pinodes*
402 genetic population variability before and after harvest. It would be particularly interesting to define
403 which level of diversity is maintained between two growing season.

404 Understanding how this gene flow influences the aggressiveness of pathogen populations during
405 the season is crucial for the development of sustainable strategies. In our study, two scenarios for the
406 evolution of the aggressiveness of *D. pinodes* strains installed in the crop were observed: i) an almost
407 constant level of aggressiveness maintained throughout the season in the winter pea crops, and ii) an

408 increase in the average aggressiveness of *D. pinodes* strains installed on spring peas, following initially
409 low aggressiveness levels. The 'spring peas' scenario was quite surprising, as the inoculum released
410 by the winter pea crop was rather aggressive. The low initial aggressiveness in the spring pea plots
411 might therefore sign other infection sources, such as alternate hosts and /or soil, as discussed above.
412 Moreover, the fact that *D. pinodes* population sampled at the beginning of the cropping season
413 displays a lower aggressiveness level suggests that an alternative hypothesis would be that spring
414 crops initially select the least aggressive components in the allo-inoculum. However, there is no
415 strong evidence or rationale to support this hypothesis. Interestingly, the fact that aggressiveness
416 levels in spring crops later increased to levels similar to those observed in trap plants and in winter
417 plots suggests either a later influx of aggressive isolates from nearby winter pea fields and/or active
418 selection of more aggressive isolates by the spring plants over the course of the epidemic – which in
419 turn would imply a polycyclic epidemic development within plots. The uniform genetic structure of *D.*
420 *pinodes* populations however does not allow to distinguish between both hypotheses, as ascospores
421 generating the late infections could come either from within or from outside the canopy.

422 This study shows that a monitoring over time allows identifying inoculum sources (auto or allo-
423 inoculum) all along the dynamic of the epidemic. Studying the temporal changes in gene and
424 genotypic diversity between the beginning and the end of an epidemic would thus be informative
425 about the different epidemiological processes involved (73). Many fungal plant pathogens alternate
426 rounds of asexual multiplication with a single annual episode of sexual reproduction (74). The
427 number of rounds of asexual multiplication has not only a demographic impact as it corresponds
428 mostly to the epidemic phase of the disease but will also have drastic consequences on the genetic
429 characteristics of pathogen populations. As a result, one would expect that both the genetic structure
430 and the gene and genotypic diversity of plant pathogens should change during the asexual phase of
431 the life cycle (74). Founder effects resulting from the colonization by a few sexually derived spores
432 and the subsequent asexual reproduction may result in reduced genotypic diversity at the beginning
433 of an epidemic. A few population genetic studies of airborne plant pathogens have used such a

434 nested hierarchical sampling strategy. Gobbin *et al.* (75) explained the arrival of new genotypes and
435 the erosion of clonal structure they observed in *Plasmopara viticola* populations by the continual
436 input of sexual spores. Examining the changes in clonal structure during the epidemic season can
437 provide insights into the balance between auto- and allo-infection processes, thus ideally
438 complementing direct epidemiological observations (i.e. disease monitoring) dedicated to the
439 quantification of the auto-infection process only (76). Contrary to a large number of necrotrophic
440 fungi, *D. pinodes* can simultaneously developed its anamorph and teleomorph forms on the same
441 plant during the growing season (32, 33). Due to the indeterminate growth of pea plant, during crop
442 growth, both pycnidia and pseudothecia can be observed on the same plant organs (33). Thus,
443 beyond these qualitative predictions, estimating the population genetic consequences of
444 intermediate rates of asexual or sexual reproduction remains a challenging task, especially when the
445 organism's life cycle consists of alternate phases of sexual and asexual reproduction. In this study we
446 particularly show that the allo-inoculum, essentially constituted by ascospores, is a driving force in
447 the epidemic dynamics of ascochyta blight of peas. As described by Savage *et al.* (6), by connecting
448 wild and cultivated compartments, this allo-inoculum allows to maintain a high level of genetic
449 variability, and to modulate pest pressure during the growing season. Our study also shows that
450 origin of this allo-inoculum still is a central question. Indeed, in the intensive French agriculture, and
451 contrary to more extensive cropping like in Australia or Canada (72, 35), the main sources of
452 inoculum are well managed by the growers. Hypotheses explaining the availability in ascospores
453 throughout the cropping season of both winter and spring peas thus need to include ascospores
454 released from stubbles or volunteers before burying, which persist for a long time in the atmosphere
455 (34), or potential reservoirs of inoculum, especially in the case of winter peas. Efficient control
456 strategies should reduce the production of allo-inoculum, and therefore to encompass all the
457 possible alternative sources, probably dispersed through the agricultural landscape. Thus, landscape
458 composition and characteristics are important to consider as they may influence pathogen ecology
459 and especially the range of hosts simultaneously or successively exploited (62), but additional

460 knowledge on the possible role, dynamics and characteristics of inoculum sources other than pea
461 (alternate hosts, soil) is also required.

462

463 **Acknowledgements**

464 Financial support to this work by the ARIMNET project “Medileg” (2012-2015; Proposal ID 396:
465 Breeding, agronomic and biotechnological approaches for reintegration and revalorization of legumes
466 in Mediterranean agriculture), INRA institute, and AGROCAMPUS Ouest, are gratefully acknowledged.

467

468 **Bibliography**

469 **1. Zadoks JC, Schein RD.** 1979. Epidemiology and plant disease management. New York, USA: Oxford
470 University Press Inc.

471 **2. Cook RJ, Veseth RJ.** 2001. *Wheat Health Management*. St. Paul, MN, USA: APS Press.

472 **3. Burdon JJ, Thrall PH.** 2008. Pathogen evolution across the agro-ecological interface: implications
473 for disease management. *Evol Appl* **1**: 57-65.

474 **4. Zhao J, Wang L, Wang Z, Chen X, Zhang H, Yao J, Zhan G, Chen W, Huang L, Kang Z.** 2013.
475 Identification of Eighteen Berberis Species as Alternate Hosts of *Puccinia striiformis* f. sp. *tritici* and
476 Virulence Variation in the Pathogen Isolates from Natural Infection of Barberry Plants in China.
477 *Phytopathol* **103**: 927-934.

478 **5. De Wolf ED, Isard SA.** 2007. Disease cycle approach to plant disease prediction. *Annu Rev*
479 *Phytopathol* **45**: 203-220.

480 **6. Savage D, Barbetti MJ, MacLeod WJ, Salam MU, Renton M.** 2012. Seasonal and diurnal patterns
481 of spore release can significantly affect the proportion of spores expected to undergo long-distance
482 dispersal. *Microb Ecol* **63**: 578-585.

483 **7. Brown JKM, Hovmøller MS.** 2002. Aerial dispersal of pathogens on the global and continental
484 scales and its impact on plant disease. *Science* **297**: 537-541.

485 **8. Nathan R, Sapir N, Trakhtenbrot A, Katul GG.** 2005. Long-distance biological transport processes
486 through the air: can nature's complexity be unfolded in silico. *Divers Distrib* **11**: 131-137.

487 **9. Barrès B, Dutech C, Andrieux A, Halkett F, Frey P.** 2012. Exploring the role of asexual multiplication
488 in poplar rust epidemics: impact on diversity and genetic structure. *Mol Ecol* **21**: 4996-5008.

- 489 **10. Thrall PH, Burdon JJ.** 1997. Host-pathogen dynamics in a metapopulation context: the ecological
490 and evolutionary consequences of being spatial. *J Ecol* **85**: 743-753.
- 491 **11. Aylor DE.** 2003. Spread of plant disease on a continental scale: Role of aerial dispersal of
492 pathogens. *Ecology* **84**: 1989-1997.
- 493 **12. Agrios GN.** 2004. *Plant Pathology* (5th Ed.). London, UK: Elsevier Academic Press.
- 494 **13. Burie JB, Langlais M, Calon nec A.** 2011. Switching from a mechanistic model to a continuous
495 model to study at different scales the effect of vine growth on the dynamic of a powdery mildew
496 epidemic. *Ann Bot* **107**: 885-895.
- 497 **14. Chakraborty, S, Newton AC.** 2011. Climate change, plant diseases and food security: an overview.
498 *Plant Pathol* **60**: 2-14.
- 499 **15. Glais I, Montarry J, Corbière R, Pasco C, Marquer B, Magalon H, Andrivon D.** 2014. Long-distance
500 gene flow outweighs a century of local selection and prevents local adaptation in the Irish famine
501 pathogen *Phytophthora infestans*. *Evol Appl* **7**: 442-452.
- 502 **16. Chen RS, McDonald BA.** 1996. Sexual reproduction plays a major role in the genetic structure of
503 populations of the fungus *Mycosphaerella graminicola*. *Genetics* **142**: 1119-1127.
- 504 **17. McDonald BA, Linde C.** 2002. Pathogen population genetics, evolutionary potential, and durable
505 resistance. *Annu Rev Phytopathol* **40**: 349-+.
- 506 **18. Burdon JJ, Silk J.** 1997. Sources and patterns of diversity in plant-pathogenic fungi. *Phytopathol*
507 **87**: 664-669.
- 508 **19. Lannou C, Soubeyrand S, Frezal L, Chadoeuf J.** 2008. Autoinfection in wheat leaf rust epidemics.
509 *New Phytol* **177**: 1001-1011.
- 510 **20. Brewer MT, Frenkel O, Milgroom MG.** 2012. Linkage disequilibrium and spatial aggregation of

511 genotypes in sexually reproducing populations of *Erysiphe necator*. *Phytopathol* **102**: 997-1005.

512 **21. Brurberg MB, Elameen A, Le VH, Naerstad R, Hermansen A, Lehtinen A, Hannukkala A, Nielsen**
513 **B, Hansen J, Andersson B.** 2011. Genetic analysis of *Phytophthora infestans* populations in the Nordic
514 European countries reveals high genetic variability. *Fungal Biol* **115**: 335-342.

515 **22. Cooke B, Jones M, Gareth D, Kaye B.** 2006. The Epidemiology of Plant Diseases, Second
516 Edition. Chapter 2, Disease assessment and yield loss. Springer, Printed in the Netherlands, pp: 583.

517 **23. Montarry J, Andrivon D, Glais I, Corbière R, Mialdea G, Delmotte F.** 2010. Microsatellite markers
518 reveal two admixed genetic groups and an ongoing displacement within the French population of the
519 invasive plant pathogen *Phytophthora infestans*. *Mol Ecol* **19**: 1965-1977.

520 **24. Sjöholm L, Anderson B, Hogberg N, Widmark AK, Yuen J.** 2013. Genotypic diversity and migra-
521 tion patterns of *Phytophthora infestans* in the Nordic countries. *Fungal Biol* **117**: 722-730.

522 **25. Lepoivre P.** 2005. Epidemiology, chapter 9, pp193-209, In : *Phytopathologie, Bases moléculaires*
523 *et biologiques des pathosystèmes et fondements des stratégies de lutte*. Les Presses de Gembloux,
524 Belgium : Ed. De Boeck, ISBN : 2-8041-4115-2.

525 **26. Bretag TW, Ramsey M.** 2001. Foliar diseases caused by fungi: Ascochyta spp. (J.M. Kraft & F.L.
526 Pflieger). In: *Compendium of pea disease and pests*. APS press, Second edition 24-28.

527 **27. Tivoli B, Banniza S.** 2007. Comparison of the epidemiology of ascochyta blights on grain legumes.
528 *Eur J Plant Pathol* **119**: 59-76.

529 **28. Ali MS, Paterson J, Crosby J.** 1982. A standard technique for detecting seed-borne pathogens in
530 peas, chemical control, and testing commercial seed in South Australia. *Aust J Exp Agric.* **32**: 1121-
531 1125.

532 **29. Nasir M, Hoppe HH, & Ebrahim Nesbat F.** 1992. The development of different pathotype groups

533 of *Mycosphaerella pinodes* in susceptible and partially resistant pea leaves. Plant Pathol **41**: 187-194.

534 **30. Xue AG, Warkentin TD.** 2001. Partial resistance to *Mycosphaerella pinodes* in field pea. Can J
535 Plant Sci **81**: 535-540.

536 **31. Zhang JX, Fernando WGD, Xue AG.** 2004. « Temporal and spatial dynamics of mycosphaerella
537 blight [*Mycosphaerella pinodes*] in field pea ». Can J Plant Pathol **26**: 522-532.

538 **32. Le May C, Guibert M, Leclerc A, Andrivon D, Tivoli B.** 2012. A Single, Plastic Population of
539 *Mycosphaerella pinodes* Causes Ascochyta Blight on Winter and Spring Peas (*Pisum sativum*) in
540 France. Appl Environ Microbiol **78**: 8431-8440.

541 **33. Roger C, Tivoli B.** 1996. Spatio-temporal development of pycnidia and perithecia and
542 dissemination of spores of *Mycosphaerella pinodes* on pea (*Pisum sativum*). Plant Pathol **45**: 518-528.

543 **34. Salam MU, Galloway J, Diggle AJ, MacLeod WJ, Maling T.** 2011. Predicting regional-scale spread
544 of ascospores of *Didymella pinodes* causing ascochyta blight disease on field pea. Australas Plant
545 Pathol **40**: 640-647.

546 **35. Davidson JA, Krysinska-Kaczmarek M, Wilmshurst CJ, McKay A, Scott ES.** 2011. Distribution and
547 survival of ascochyta blight pathogens in field pea cropping soils of Australia. Plant Dis **95**: 1217-1223.

548 **36. Jarosz AM, Burdon JJ.** 1991. Host-pathogen interactions in natural populations of *Linum mar-*
549 *ginale* and *Melampsora lini*: II. Local and regional variation in patterns of resistance and racial struc-
550 ture. Evolution **45**: 1618-1627.

551 **37. Antonovics J, Iwasa Y, Hassell MP.** 1995. A generalized model of parasitoid, venereal, and vector-
552 based transmission processes. Am Nat **145**: 661-675.

553 **38. Burdon JJ, Thompson JN.** 1995. Changed patterns of resistance in a population of *Linum*
554 *marginale* attacked by the rust pathogen *Melampsora lini*. J Ecol **83**: 199-206.

- 555 **39. Ericson L, Burdon JJ, Müller WJ.** 1999. Spatial and temporal dynamics of epidemics of the rust
556 fungus *Uromyces valerianae* on populations of its host *Valeriana salina*. *J Ecol* **87**: 649-658.
- 557 **40. Smith DL, Ericson L, Burdon JJ.** 2003. Epidemiological patterns at multiple spatial scales: an 11-
558 year old study of a *Triphragmium ulmariae*–*Filipendula ulmaria* metapopulation. *J Ecol* **91**: 890-903.
- 559 **41. Schoeny A, Jumel S, Rouault F, May CL, Tivoli B.** 2007. Assessment of airborne primary inoculum
560 availability and modelling of disease onset of ascochyta blight in field peas. *Eur J Plant Pathol.* **119**:
561 87-97.
- 562 **42. Laine AL, Hanski I.** 2006. Large-scale spatial dynamics of a specialist plant pathogen in a
563 fragmented landscape. *J Ecol* **94**: 217-226.
- 564 **43. Fisher N I, Lewis T, Embleton BJJ.** 1987. Statistical analysis of spherical Data. Cambridge, UK:
565 University Press, 329 pp.
- 566 **44. Onfroy C, Tivoli B, Corbière R, Bouznad Z.** 1999. Cultural, molecular and pathogenic variability of
567 *Mycosphaerella pinodes* and *Phoma medicaginis* var. *pinodella* isolates from dried pea (*Pisum*
568 *sativum*) in France. *Plant Pathol* **48**: 218-229.
- 569 **45. Lodhi MA, Ye GN, Weeden NF, Reisch BI.** 1994. A simple and efficient method for DNA extraction
570 from grapevine cultivars and *Vitis* species. *Plant Mol Biol Report.* **12**.
- 571 **46. Vos P, Hogers R, Bleeker M, Reijans M, Vandelee T, Hornes M, Frijters A, Pot J, Peleman J, Kuiper**
572 **M.** 1995. Aflp - a New Technique for Dna-Fingerprinting. *Nucleic Acids Res* **23**: 4407-4414.
- 573 **47. Peakall R, Smouse PE.** 2006. GENALAEX 6: genetic analysis in Excel. Population genetic software
574 for teaching and research. *Mol Ecol Res* **6**: 288-295.
- 575 **48. Shannon C E, Weaver W.** 1949. The Mathematical Theory of Communication. University of Illinois
576 Press, Urbana, IL.

577 **49. Grünwald NJ, Goodwin SB, Milgroom MG, Fry WE.** 2003. Analysis of genotypic diversity data for
578 populations of microorganisms. *Phytopathol* **93**: 738-746.

579 **50. Agapow PM, Burst A.** 2001. Indices of multilocus lineage disequilibrium. *Mol Ecol Notes* **1**: 101-
580 102.

581 **51. Excoffier L, Laval G, Schneider S.** 2005. Arlequin (version 3.0): An integrated software package for
582 population genetics data analysis. *Evol Bioinforma* **1**: 47-50.

583 **52. Jombart T, Devillard S, Dufour AB, Pontier D.** 2008. Revealing cryptic spatial patterns in genetic
584 variability by a new multivariate method. *Heredity* **101**: 92-103.

585 **53. Pritchard JK, Stephens M, Donnelly P.** 2000. Inference of population structure using multilocus
586 genotype data. *Genetics* **155**: 945-959.

587 **54. Baranger A, Aubert G, Arnau G, Lainé AL, Deniot G, Potier J, Weinachter C, Lejeune I, Lallemand**
588 **J, Burstin J.** 2004. Genetic diversity within *Pisum sativum* using protein and PCR-based markers. *Theor*
589 *Appl Genet* **108**: 1309-1321

590 **55. Onfroy C, Baranger A, Tivoli B.** 2007. Biotic factors affecting the expression of partial resistance in
591 pea to ascochyta blight in a detached stipule assay. *Eur J Plant Pathol* **119**: 13–27.

592 **56. Shaner G, Finney RE.** 1977. The effect of nitrogen fertilization on the expression of slow-
593 mildewing resistance in Knox wheat. *Phytopathol* **67**: 1051–6.

594 **57. Setti B, Bencheikh M, Henni DE, Neema C.** 2012. Genetic variability and population structure of
595 *Mycosphaerella pinodes* in western Algeria using aflp fingerprinting. *J Plant Pathol* **94**: 127-133.

596 **58. Rhaïem A, Cherif M, Peever TL, Dyer PS.** 2008. Population structure and mating system of
597 *Ascochyta rabiei* in Tunisia: evidence for the recent introduction of mating type 2. *Plant Pathol* **57**:
598 540-551.

599 **59. Carlier J, Lebrun MH, Zapater MF, Dubois C, Mourichon X.** 1996. Genetic structure of the global
600 population of banana black leaf streak fungus *Mycosphaerella fijiensis*. *Mol Ecol* **5**: 499-510.

601 **60. Alfonso C, Rapso R, Malgarejo P.** 2000. Genetic diversity in *Botrytis cinerea* populations on
602 vegetable crops in greenhouses in south-eastern Spain. *Plant Pathol* **49**: 243-251.

603 **61. Le May C, Guibert M, Baranger A, Tivoli B.** 2014. A wide range of cultivated legume species act as
604 alternative hosts for the pea ascochyta blight fungus, *Didymella pinodes*. *Plant Pathol* **63**: 877-887.

605 **62. Plantegenest M, Le May C, & Fabre F.** 2007. Landscape epidemiology of plant diseases. *J Roy Soc*
606 *Inter* **4**: 963-972.

607 **63. Lawyer AS.** 1984. Diseases caused by *Ascochyta* spp. *In* Compendium of pea diseases. *Edited by*
608 J.M. Kraft and F.L. Pflieger. APS Press, St Paul, Minn. pp. 11–15.

609 **64. Trapero-Casas A, Kaiser WJ.** 2009. Alternative hosts and plant tissues for the survival, sporulation
610 and spread of the Ascochyta blight pathogen of chickpea. *Eur J Plant Pathol* **125**: 573–587.

611 **65. Weimer JL.** 1947. Resistance of *Lathyrus* spp. and *Pisum* spp. to *Ascochyta pinodella* and
612 *Mycosphaerella pinodes*. *J Agr Res* **75**: 181-190.

613 **66. Gurung AM, Pang ECK, Taylor PWJ.** 2002. Examination of *Pisum* and *Lathyrus* species a sources of
614 ascochyta blight resistance for field pea (*Pisum sativum*). *Australas Plant Pathol* **31**: 41-45.

615 **67. Bretag TW.** 2004. Review: Ascochyta blight in field peas. Victorian Department of Primary
616 Industries: Horsham, Vic.

617 **68. Singh AB, Babu CR.** 1980. Diurnal periodicity of common allergenic pollen. *Allergy* **35**: 311–317.

618 **69. Abbo S, Frenkel O, Sherman A, Shtienberg D.** 2007. The sympatric *Ascochyta* pathosystems of
619 Near Eastern legumes, a key for better understanding of pathogen biology. *Eur J Plant Pathol* **119**:
620 111-118.

621 **70. Akisanmi OA, Chakraborty S, Backhouse D, Simpfendorfer S.** 2007. Passage through alternative
622 hosts changes the fitness of *Fusarium graminearum* and *Fusarium pseudograminearum*. Environ
623 Microbiol **9**: 512-520.

624 **71. Hare WW, Walker JC.** 1944. Ascochyta diseases of canning pea. Wisc Res Bul **150**: 1–31.

625 **72. Wallen VR, Jeun J.** 1968. Factors limiting the survival of Ascochyta spp of peas in soil. Can J Bot
626 **46**: 1279-1286.

627 **73. Prugnolle F, De Meeûs T.** 2010. Apparent high recombination rates in clonal parasitic organisms
628 due to inappropriate sampling design. Heredity **104**: 135-140.

629 **74. Barrès B, Dutech C, Andrieux A, Halkett F, Frey P.** 2012. Exploring the role of asexual
630 multiplication in poplar rust epidemics: impact on diversity and genetic structure. Mol Ecol **21**: 4996-
631 5008.

632 **75. Gobbin D, Jermini M, Loskill B, Pertot I, Raynal M, Gessler C.** 2005. Importance of secondary
633 inoculum of *Plasmopara viticola* to epidemics of grapevine downy mildew. Plant Pathol **54**: 522-534.

634 **76. Lannou C, Soubeyrand S, Frezal L, Chadoeuf J.** 2008. Autoinfection in wheat leaf rust epidemics.
635 New Phytol **177**: 1001-1011.

636

1 **Table I:** Genetic characterization of *Didymella pinodes* strains sampled on winter or spring pea plots
 2 (WP or SP) and on Trap winter plants (TWP) or Trap spring plants (TSP). Na = No. of different alleles, I =
 3 Shannon's information index ($= -1\sum(p \cdot \ln(p) + q \cdot \ln(q))$), h = Diversity ($= 1 - (p^2 + q^2)$), uh = Unbiased diversity ($=$
 4 $(N/(N-1)) h$), where p = Band freq., and q = 1 - p (GenAlEx v6.5)

5

| Source | Sub-populations | Sample date (d/m) | n | Na | I | uh | H' |
|--------|-----------------|-------------------|------|------|------|------|------|
| WP | WP7 | 17 February | 30 | 1.97 | 0.51 | 0.35 | 0.83 |
| | WP8 | 28 February | 30 | 2.00 | 0.58 | 0.41 | |
| | WP10 | 29 March | 30 | 1.96 | 0.49 | 0.34 | |
| | WP12 | 25 April | 30 | 1.99 | 0.61 | 0.43 | |
| | WP14 | 23 May | 30 | 1.99 | 0.56 | 0.39 | |
| SP | SP1 | 30 May | 30 | 1.98 | 0.59 | 0.42 | 0.86 |
| | SP2 | 13 June | 30 | 1.97 | 0.57 | 0.40 | |
| | SP3 | 27 June | 30 | 1.99 | 0.58 | 0.41 | |
| TWP | TWP19 | 28 January | 13 | 1.83 | 0.50 | 0.37 | 0.70 |
| | TWP20 | 04 February | 14 | 1.77 | 0.43 | 0.31 | |
| | TWP21 | 11 February | 5 | 1.06 | 0.18 | 0.15 | |
| | TWP22 | 18 February | 15 | 1.94 | 0.55 | 0.40 | |
| | TWP23 | 25 February | 13 | 1.81 | 0.44 | 0.31 | |
| | TWP24 | 04 March | 12 | 1.75 | 0.46 | 0.34 | |
| | TWP28 | 01 April | 15 | 1.82 | 0.48 | 0.35 | |
| | TWP29 | 08 April | 15 | 1.88 | 0.53 | 0.39 | |
| | TWP30 | 15 April | 14 | 1.82 | 0.48 | 0.35 | |
| | TWP31 | 22 April | 15 | 1.79 | 0.39 | 0.28 | |
| | TWP32 | 29 April | 15 | 1.48 | 0.28 | 0.20 | |
| | TWP33 | 05 May | 13 | 1.86 | 0.48 | 0.35 | |
| | TWP34 | 16 May | 12 | 1.49 | 0.31 | 0.23 | |
| | TWP35 | 20 May | 13 | 1.85 | 0.52 | 0.38 | |
| TWP36 | 27 May | 15 | 1.41 | 0.28 | 0.20 | | |
| TSP | TWP37 | 03 June | 15 | 1.86 | 0.47 | 0.34 | 0.76 |
| | TSP2 | 20 May | 4 | 1.06 | 0.25 | 0.23 | |
| | TSP3 | 27 May | 9 | 1.86 | 0.51 | 0.39 | |
| | TSP4 | 03 June | 10 | 1.84 | 0.49 | 0.37 | |
| | TSP5 | 10 June | 14 | 1.84 | 0.48 | 0.35 | |
| | TSP6 | 20 June | 14 | 1.83 | 0.46 | 0.33 | |
| | TSP7 | 27 June | 15 | 1.78 | 0.45 | 0.33 | |

6

7

8 **Table II:** Analysis of molecular variance (AMOVA) for *D. pinodes* strains sampled from different plots
 9 (WP, SP) and trap plants (TWP, TSP) based on AFLP markers.

| Source of variation | d.f. | Sum of square | Variance components | Variation (%) | Fixation Indices | P-value* |
|-----------------------------------------|------|---------------|---------------------|---------------|------------------|-----------|
| Among populations | 3 | 3781.378 | 8.26273 | 9.62 | FCT 0.09616 | <0.000001 |
| Among sub-population within populations | 26 | 5535.163 | 8.62056 | 10.03 | FSC 0.11100 | <0.000001 |
| Within sub-populations | 490 | 33830.767 | 69.04238 | 80.35 | FST 0.19649 | <0.000001 |

10
 11

12 **Table III:** ANOVAs on AUDPC values at the population (A) and at the sub-population (B) levels.
13 Sources of variation are *D. pinodes* population (or sub-population), pea genotype and the
14 corresponding two-way interaction between these variables. Statistically significant effects are
15 indicated by asterisks (*, $P < 0.001$). df: degrees of freedom.

16

17 **A**

| Source of variation | df | F value | P>F |
|-------------------------------|-----|---------|----------|
| Sub-population | 3 | 28.41 | <0,0001* |
| Pea genotype | 2 | 7.71 | 0.0006* |
| Sub-population * Pea genotype | 6 | 0.11 | 0.9955 |
| Error | 180 | | |

18

19

20 **B**

| Source of variation | df | F value | P>F |
|-------------------------------|-----|---------|----------|
| Sub-population | 12 | 11.22 | <0,0001* |
| Pea genotype | 2 | 8.50 | 0.0003* |
| Sub-population * Pea genotype | 24 | 0.23 | 0.9999 |
| Error | 153 | | |

21

22




Fig. 1: Maps representing the evolution of disease severity due to ascochyta blight within winter and spring pea plots and for different sampling dates (**Winter sampling dates:** **A:** 17 February, **B:** 28 February, **C:** 29 March, **D:** 25 April, **E:** 23 May; **Spring sampling dates:** **F:** 30 May, **G:** 13 June, **H:** 27 June). Each plot was divided into 1m² micro-plots (A1 to C10) where one plant was randomly sampled at each sampling date. Disease was scored on each stipule of the sampled plants using the adapted disease scale of Tivoli (1994) (40), and a mean disease score (D_j) was calculated. Disease distribution maps were created for each sampling date using the Arcmap® software.




Fig. 2: Standardized index of association (I_A) calculated from the clone-corrected dataset, rejecting the absence of linkage disequilibrium among microsatellite loci ($P\text{-value} < 0.001$).




Fig. 3: Box plots representing the distribution of pairwise F_{ST} within sub-population (sp-sp, wp-wp, twp-twp, tsp-tsp) and among each sub-population (wp-sp, wp-twp, sp-tsp).




Fig. 4: Principal component analysis (PCA) realized on the *D. pinodes* sub-populations respectively sampled within winter pea plot (WP, 150 strains), spring pea plot (SP, 90 strains), trap winter pea (TWP, 214 strains), and trap spring plant (TSP, 66 strains).




Fig. 5: Mean aggressiveness level (AUDPC) for each *D. pinodes* population **(A)** and for each *D. pinodes* sub-population **(B)**. AUDPC was measured on detached stipules of three pea genotypes (Lumina, Enduro and DP) for *D. pinodes* strains sampled at the different sampling dates on winter (WP), spring (SP) pea plots, and for *D. pinodes* strains sampled on winter (TWP) and spring (TSP) trap plants. AUDPC was calculated as described by Shaner & Finney (55) by estimating the integral of the disease progress curve including assessment of the disease at 2, 4 and 7 days after inoculation.