Synthesis, structure and photophysical properties of NIR aza-BODIPYs with -F/-N-3/NH₂ groups at 1,7-positions

Xin-Dong Jiang, Jian Guan, Jiuli Zhao, Boris Le Guennic, Denis Jacquemin, Zhigang Zhang, Shangdong Chen, Linjiu Xiao

► To cite this version:

Xin-Dong Jiang, Jian Guan, Jiuli Zhao, Boris Le Guennic, Denis Jacquemin, et al.. Synthesis, structure and photophysical properties of NIR aza-BODIPYs with -F/-N-3/NH₂ groups at 1,7-positions. *Dyes and Pigments*, 2017, 136, pp.619-626. 10.1016/j.dyepig.2016.09.019 . hal-01438101

HAL Id: hal-01438101

<https://hal-univ-rennes1.archives-ouvertes.fr/hal-01438101>

Submitted on 18 May 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Synthesis, structure and photophysical properties of NIR aza-BODIPYs with –F/-N₃/–NH₂ groups at 1,7-positions

Xin-Dong Jiang,^{a,b,*} Jian Guan,^a Jiuli zhao,^a Boris Le Guennic,^{c,*} Denis Jacquemin,^d Zhigang Zhang^a Shangdong Chen^a and Linjiu Xiao^a

^aCollege of Applied Chemistry, Shenyang University of Chemical Technology, Shenyang, 110142, China. E-mail: xdjiang@syuct.edu.cn; Tel: +86 024 89387219.

^bState Key Laboratory of Fine Chemicals, Dalian University of Technology, Dalian 116024, China.

^cInstitut des Sciences Chimiques de Rennes, UMR 6226 CNRS, Université de Rennes1, 263 Av. du Général Leclerc, 35042 Cedex Rennes, France. E-mail: boris.leguennic@univ-rennes1.fr

^dLaboratoire CEISAM, UMR CNRS 6230, Université de Nantes, 2 Rue de la Houssinière, BP92208, 44322 Cedex 3 Nantes, France

Corresponding author: Tel: +86 024 89387219. Fax: +86 024 89388211

E-mail: xdjiang@syuct.edu.cn

Abstract: With 4-fluorostyrene as the starting material, symmetric/asymmetric fluorine-containing aza-difluoroboradiaza-*s*-indacenes (aza-BODIPYs) **3-6** at 1/7-position were successfully prepared. NaN₃ undertook nucleophilic aromatic substitution via replacement of F atoms of **4** to form the N₃/NH₂-containing aza-BODIPYs **9** and **10**. The solid-state structure of aza-BODIPY **4** was confirmed by single crystal X-ray analysis, the F-C_{Ph} bond distance of which was found to be shorter than those of the reported fluoride. These aza-BODIPYs absorb and emit in

the NIR region. Surprisingly, although the withdrawing group was introduced at 1/7 positions of **3-6**, the effect of the fluorine atoms to the photophysical properties is trifling. However, **9** and **10** with $-N_3/-NH_2$ groups resulted in a remarkable bathochromic shift. Aza-BODIPY **10** could be also used as a turn-on fluorescent probe for pH value. Moreover, the main photophysical data are well supported and explained by cLR-PCM($CHCl_3$)-M06-2X/6-311+G(2d,p)/SOS-CIS(D) calculations.

Keywords: aza-BODIPY • NIR • fluorine • fluorescence • TD-DFT calculations

1. Introduction

Near-infrared (NIR) fluorescent dyes can greatly reduce background absorption, fluorescence, light scattering, and can be used as building blocks to improve the sensitivity of the probes and sensors [1]. More importantly, NIR fluorescent dyes with strong absorption and emission in the low energy regions (650-900 nm) have deep penetration in tissues which allows more efficient in vivo imaging and photodynamic therapy [2]. The safe, non-invasive nature of NIR fluorescent imaging has attracted great interests in the design and synthesis of novel NIR compounds [3]. Indeed, NIR derivatives such as azo, cyanine, quinone, and phthalocyanine dyes have been developed though, in most cases, insufficient photo-stability and/or difficult modification obstructed their applications [3]. Most NIR fluorescent dyes possess highly π -conjugated systems which lead to significantly reduced fluorescence quantum yields due to increased internal conversion. However, BODIPY and their derivatives are well-known to be highly fluorescent and very stable dyes, presenting tunable emission wavelengths and an exceptional insensitivity to the polarity of the medium [4]. Therefore, they are very promising candidates for biological labeling applications, and are already used in those fields [4,5].

By using an imine instead of a methane in BODIPY system to narrow the HOMO–LUMO gap one obtains aza-BODIPY that are known to be an attractive platform to reach NIR absorption (Fig. 1a) [6]. Furthermore, by introducing an electron-donating/withdrawing group one can further reduce the HOMO–LUMO gap by increasing the HOMO and/or decreasing the LUMO energy and reach more redshifted absorption and emission bands [7]. For instance, aza-BODIPYs with an electron-donating group such as $-(p-NMe_2)Ph/-(p-OMe)Ph$ at 3/5-positions (Fig. 1b) exhibit bathochromic shift of the absorption/emission bands [8]. By contrast, aza-BODIPYs bearing an electron-withdrawing group (such as the $-CN$ group) at 1/7-positions (Fig. 1c) are also known to undergo a redshift of the absorption/emission bands [9]. Since fluorine is a strong inductive acceptor, introduction of fluorine atom(s) in aza-BODIPY was considered in the present work. Importantly, the introduction of a fluorine atom into an organic molecule often induces significant, if not remarkable, variations of its chemical, physical and biological properties, due to the lipophilizing nature, strong electron-withdrawing nature and specific steric repulsion of fluorine [10]. In line with their widespread use

in both medicinal chemistry and in materials science, fluorinated molecules have attracted a considerable interest in the framework of molecular design. Therefore, studies on the characteristic of a fluorine-containing aza-BODIPY at 1/7-positions are of importance. We have been already exploring fluorides [11], and our most recent research interest lies in novel BODIPY/aza-BODIPY family of fluorescent dyes [12].

<Fig. 1>

2. Experimental methods

2.1 General: The melting points were measured using a SGW® X-4 melting point apparatus. ^1H NMR spectra were recorded on a Bruker AVANCE III 500 MHz spectrometer. ^1H NMR chemical shifts (δ) are given in ppm downfield from Me_4Si , determined by chloroform ($\delta = 7.26$ ppm). ^{13}C NMR spectra were recorded on a Bruker AVANCE III 125 MHz spectrometer. ^{13}C NMR chemical shifts (δ) are reported in ppm with the internal CDCl_3 at δ 77.0 ppm as standard. HRMS (ESI) was measured by micro TOF-Q II.

Tetrahydrofuran (THF) was freshly distilled from Na/benzophenone, *n*-hexane was distilled over Na, and other solvents were distilled over CaH_2 . Merck silica gel 60 was used for the column chromatography. Fluorescence spectra were recorded on F-4600 spectrophotometer. UV/Vis spectra were recorded on UV-2550 spectrophotometer at room temperature. The refractive index of the medium was measured by 2 W Abbe's refractometer at 20 °C. The fluorescence quantum yields (Φ_f) of the BODIPY systems were calculated using the following relationship (equation 1):

$$\Phi_f = \Phi_{\text{ref}} F_{\text{sampl}} A_{\text{ref}} n_{\text{sampl}}^2 / F_{\text{ref}} A_{\text{sampl}} n_{\text{ref}}^2 \quad (1)$$

Here, F denotes the integral of the corrected fluorescence spectrum, A is the absorbance at the excitation wavelength, ref and sampl denote parameters from the reference and unknown experimental samples, respectively. The reference systems used was boronazadipyrrromethene dye [13] as standard ($\Phi_f = 0.36$ in chloroform, $\lambda_{\text{abs}} = 688$ nm) for **3-6**, **9** and **10**.

2.2 Computational details: DFT, and TD-DFT calculations on molecules were carried out with the latest version of the Gaussian 09 program package [14], applying both a tightened self-consistent field convergence criterion (10^{-9} – 10^{-10} au) and an improved optimization threshold (10^{-5} au on average forces). For each molecule, we have optimized the geometry of both the ground and the first excited states, as well as computed the vibrational spectra of both states. The same DFT integration grid, namely the so-called ultrafine pruned (99,590) grid, was used for both the ground and excited states. All DFT and TD-DFT calculations were performed with the M06-2X hybrid exchange–correlation functional, that has been shown to be an adequate choice for investigating excited-state energies and structures of many classes of molecules [15]. Atoms were described with the 6-311+G(2d,p) basis sets. Solvation effects (here CHCl_3) have been quantified using the Polarizable Continuum Model (PCM) [16].

Here, we have applied the corrected LR approach [17]. The SOS-CIS(D)/6-311+G(2d,p) energies were determined with the Q-Chem package using the resolution of identity (RI) scheme with the same basis set. The interested reader can find the whole applied computational protocol in ref. 18.

Synthesis of 1-Fluoro-4-(1,2-dibromoethyl)benzene: Under N₂, Br₂ (0.42 ml, 8.1 mmol) in CH₂Cl₂ (20 ml) was dropwise added to 4-fluorostyrene (1.0 g, 8.1 mmol) in CH₂Cl₂ (40 ml) at 0 °C for 30 min. The color of the mixture changed from red to colorless. The reaction was allowed to warm up to room temperature slowly. After removal of the solvents by evaporation, the resulting crude mixture was recrystallization (*n*-hexane) to afford 1-fluoro-4-(1,2-dibromoethyl)benzene (1.82 g, 80%) as yellow solids.

2.3 Synthesis

Synthesis of 3-(4-fluorophenyl)-2H-azirine: Under N₂, NaN₃ (840 mg, 12.9 mmol) was slowly added to 1-fluoro-4-(1,2-dibromoethyl)benzene (1.8 g, 6.3 mmol) in DMSO (20 ml) at 20 °C and stirred for 12 h at 25 °C. Then, 50% NaOH (1 ml) was added to the mixture and stirred for 24 h at 25 °C. The reaction was quenched with 10% NaHCO₃ (50 ml). The mixture was extracted with CH₂Cl₂ (2 × 30 ml), and the organic layer was washed with brine (2 × 30 ml) and dried over anhydrous MgSO₄. After removal of the solvents by evaporation, the red oil was obtained. Toluene (40 ml) was added the red oil and the mixture was refluxed for 2 h. After removal of the solvents by evaporation, the resulting crude mixture was separated by column chromatography (*n*-hexane : CH₂Cl₂ = 1 : 1) to afford 3-(4-fluorophenyl)-2H-azirine (510 mg, 60%) as yellow oil. This mixture was direct used for the next step without further purification. ¹H NMR (500 MHz, CDCl₃): δ (ppm) 7.22-7.25 (m, 2H), 7.13-7.17 (m, 2H), 2.34 (s, 2H).

Synthesis of pyrrole 1: Under N₂, 1-tetralone (71.0 mg, 0.59 mmol) in DMSO (10 ml) was added to NaH (excess) in DMSO (10 ml) at -40 °C and stirred for 10 min. Then, 3-(4-fluorophenyl)-2H-azirine (80 mg, 0.59 mmol) in DMSO (5 ml) was added and the resulting mixture was stirred for 1 h at the same temperature. The reaction was allowed to warm up to room temperature slowly. It was quenched with water, neutralized with dilute HCl to a pH about 7. The mixture was extracted with CH₂Cl₂ (2 × 30 ml), and the organic layer was washed with brine (2 × 30 ml) and dried over anhydrous MgSO₄. After removal of the solvents by evaporation, the resulting crude mixture was separated by column chromatography (*n*-hexane : CH₂Cl₂ = 1 : 1) to afford pyrrole **1** as hoary solids (76.9 mg, 55%). ¹H NMR (500 MHz, CDCl₃): δ (ppm) 8.45 (br s, 1H), 7.50-7.53 (m, 4H), 7.40 (t, ³J = 8.0 Hz, 2H), 7.25 (d, ³J = 7.5 Hz, 1H), 7.80-7.09 (m, 1H), 7.05 (t, ³J = 8.5 Hz, 2H), 6.76 (m, 1H). HRMS-MALDI (m/z): [M + H]⁺ calcd for C₁₆H₁₃FN: 238.10265; found: 238.10268.

Synthesis of pyrrole 2: 6-Methoxytetralone (218 mg, 1.45 mmol) was used as the starting materials, and pyrrole **2** was obtained green solids (232.8 mg, 60%). ¹H NMR

(500 MHz, CDCl₃): δ (ppm) 8.34 (br s, 1H), 7.51 (d, $^3J = 9.0$ Hz, 1H), 7.50 (d, $^3J = 9.0$ Hz, 1H), 7.44 (d, $^3J = 9.0$ Hz, 2H), 7.02-7.06 (m, 3H), 6.94 (d, $^3J = 8.5$ Hz, 2H), 6.65 (m, 1H), 3.84 (s, 1H). HRMS-MALDI (m/z): [M + H]⁺ calcd for C₁₇H₁₅FNO: 268.11322; found: 268.11322.

Synthesis of aza-BODIPY 3: Sodium nitrite (6.9 mg, 0.1 mmol) was added to a suspension of pyrrole **1** (47.5 mg, 0.2 mmol) in a mixture of acetic acid/anhydride (1 ml/0.4 ml) at 0 °C, and was stirred for 10 min. The color changed from colorless to brown, then green, and finally dark green was observed. After 0.5 h stirring at room temperature, the mixture was heated at 80 °C for 0.5 h. Crushed ice was added to the mixture, the resulted dark green dye was filtered, washed with water. The dark green dye was dissolved in CH₂Cl₂, filtered through a pad of alumina (activity III). Solvent was removed under reduced pressure, and the residue was dissolved in dry 1,2-dichloroethane. Triethylamine (0.28 ml, 2.0 mmol) was added, followed by dropwise addition of BF₃·Et₂O (0.50 ml, 4.0 mmol) with stirring at room temperature. The mixture was stirred for 0.5 h, then heated in 80 °C for 0.5 h. The reaction was quenched with crushed ice, extracted with CH₂Cl₂, and purified by chromatography on silica gel followed by recrystallization from CH₂Cl₂/*n*-hexane to afford aza-BODIPY **3** (24.5 mg, 46%) as coppery solids. M.p.: 240.0–240.8 °C (decomp.). ¹H NMR (500 MHz, CDCl₃): δ (ppm) 8.02-8.04 (m, 8H), 7.48-7.50 (m, 6H), 7.16 (t, $^3J = 8.5$ Hz, 4H). ¹³C NMR (125 MHz, CDCl₃): δ (ppm) 159.6, 145.4, 143.0, 131.4, 131.2, 131.1, 131.0, 129.5, 128.6, 118.8, 115.8, 115.7. HRMS-MALDI (m/z): [M + Na]⁺ calcd for C₃₂H₂₀BF₄N₃Na: 556.1584; found: 556.1583.

Synthesis of aza-BODIPY 4: Pyrrole **2** (53.6 mg, 0.2 mmol) was used as the starting materials, and aza-BODIPY **4** was obtained green solids (25.5 mg, 43%). M.p.: 290.0–291.0 °C (decomp.). ¹H NMR (500 MHz, CDCl₃): δ (ppm) 8.08 (d, $^3J = 8.5$ Hz, 4H), 8.01 (dd, $^3J = 8.5$ Hz, 4H), 7.14 (t, $^3J = 8.5$ Hz, 4H), 7.01 (d, $^3J = 8.5$ Hz, 4H), 6.99 (s, 2H), 3.89 (s, 6H). ¹³C NMR (125 MHz, CDCl₃): δ (ppm) 161.9, 158.1, 145.1, 141.9, 131.6, 131.0, 128.6, 123.9, 118.3, 115.7, 115.5, 114.2. HRMS-MALDI (m/z): [M + H]⁺ calcd for C₃₄H₂₅BF₄N₃O₂: 594.19705; found: 594.19678.

Synthesis of aza-BODIPY 5: Sodium nitrite (11.6 mg, 0.16 mmol) was added to a suspension of 2,4-diphenyl-1*H*-pyrrole (36.9 mg, 0.16 mmol) in acetic acid (1 mL) at 0 °C, and was stirred for 10 min. The color changed from colorless to brown, then green, and finally brown was observed. The second pyrrole moiety **1** (40 mg, 0.16 mmol) was added, followed by addition of acetic anhydride (0.4 mL). The mixture turned blue immediately. After 0.5 h stirring, the mixture was heated at 80 °C for 0.5 h. Crushed ice was added to the mixture, the resulted blue dye was filtered, washed with water. The blue dye was dissolved in CH₂Cl₂, filtered through a pad of alumina (activity III). Solvent was removed under reduced pressure, and the residue was dissolved in dry 1,2-dichloroethane. Triethylamine (0.28 mL, 2.0 mmol) was added, followed by dropwise addition of BF₃·Et₂O (0.50 mL, 4.0 mmol) with stirring at room temperature. The mixture was stirred for 0.5 h, then heated in 80 °C for 0.5 h. The

reaction was quenched with crushed ice, extracted with CH_2Cl_2 , and purified by chromatography on silica gel followed by recrystallization from $\text{CH}_2\text{Cl}_2/n$ -hexane to afford aza-BODIPY **5** (42.0 mg, 51%) as coppery solids. M.p.: 230–231.0 °C (decomp.). ^1H NMR (500 MHz, CDCl_3): δ (ppm) 8.01–8.07 (m, 8H), 7.46–7.49 (m, 9H), 7.14 (t, (t, $^3J = 8.5$ Hz, 2H), 7.04 (s, 1H), 6.98 (s, 1H). ^{13}C NMR (125 MHz, CDCl_3): δ (ppm) 159.6, 159.3, 145.5, 145.3, 144.2, 142.8, 132.1, 131.4, 131.2, 131.1, 131.0, 130.9, 129.6, 129.5, 129.4, 129.3, 129.2, 128.6, 128.5, 126.4, 119.2, 118.6, 115.7, 115.6. HRMS-MALDI (m/z): $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{32}\text{H}_{22}\text{BF}_3\text{N}_3$: 516.18534; found: 516.18463.

Synthesis of aza-BODIPY 6: 2-(4-methoxyphenyl)-4-phenyl-1H-pyrrole (25.0 mg, 0.093 mmol) and pyrrole **2** (24.1 mg, 0.096 mmol) was used as the starting materials, and aza-BODIPY **6** was obtained green solids (26.2 mg, 49%). M.p.: 257.0–258.0 °C (decomp.). ^1H NMR (500 MHz, CDCl_3): δ (ppm) 8.01–8.10 (m, 8H), 7.41–7.48 (m, 3H), 7.13 (t, $^3J = 8.5$ Hz, 2H), 7.04 (s, 1H), 7.01 (dd, $^3J = 8.5$ Hz, $^4J = 3.0$ Hz, 4 H), 6.98 (s, 1H), 3.89 (s, 3H), 3.88 (s, 3H). ^{13}C NMR (125 MHz, CDCl_3): δ (ppm) 161.8, 158.2, 158.0, 157.9, 145.2, 145.1, 143.2, 141.7, 132.4, 131.5, 131.0, 130.9, 129.3, 129.2, 128.6, 128.5, 124.1, 123.9, 118.7, 118.6, 118.1, 115.6, 115.5, 114.2. HRMS-MALDI (m/z): $[\text{M} + \text{Na}]^+$ calcd for $\text{C}_{34}\text{H}_{25}\text{BF}_3\text{N}_3\text{NaO}_2$: 598.1890; found: 598.1889.

Synthesis of aza-BODIPY 9: NaN_3 (8.8 mg, 0.13 mmol) was added to a solution of **4** (25.0 mg, 0.042 mmol) in DMF (4 ml), and then the mixture was stirred at 40 °C for 30 min. The reaction was quenched with water, extracted with CH_2Cl_2 , and purified by chromatography on silica gel followed by recrystallization from $\text{CH}_2\text{Cl}_2/n$ -hexane to afford aza-BODIPY **9** (15.3 mg, 57%) as coppery solids. ^1H NMR (500 MHz, CDCl_3): δ (ppm) 8.04 (d, $^3J = 8.5$ Hz, 4H), 7.79 (d, $^3J = 8.5$ Hz, 4H), 6.99 (d, $^3J = 8.5$ Hz, 4H), 6.85 (s, 2H), 6.75 (d, $^3J = 8.5$ Hz, 4H), 3.87 (s, 6H). HRMS-MALDI (m/z): $[\text{M} - 4\text{N} + 5\text{H}]^+$ calcd for $\text{C}_{34}\text{H}_{29}\text{BF}_2\text{N}_5\text{O}_2$: 588.23769; found: 588.23730.

Synthesis of aza-BODIPY 10: NaN_3 (3.0 mg, 0.046 mmol) was added to a solution of **4** (27.4 mg, 0.046 mmol) in DMF (5 ml), and then the mixture was stirred at 40 °C for 1 h. The reaction was quenched with NaHS aqueous solution, extracted with CH_2Cl_2 , and purified by chromatography on silica gel followed by recrystallization from $\text{CH}_2\text{Cl}_2/n$ -hexane to afford aza-BODIPY **10** (13.5 mg, 50%) as coppery solids. ^1H NMR (500 MHz, CDCl_3): δ (ppm) 8.08 (d, $^3J = 8.5$ Hz, 2H), 8.03 (td, $^3J = 8.5$ Hz, $^4J = 3.0$ Hz, 4H), 7.96 (d, $^3J = 8.5$ Hz, 2H), 7.15 (t, $^3J = 8.5$ Hz, 2H), 7.00 (dd, $^3J = 8.5$ Hz, $^4J = 3.0$ Hz, 4H), 6.92 (s, 1H), 6.90 (s, 1H), 6.74 (d, $^3J = 8.5$ Hz, 2H), 4.04 (br s, 2H), 3.88 (s, 6H). HRMS-MALDI (m/z): $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{34}\text{H}_{27}\text{BF}_3\text{N}_4\text{O}_2$: 591.21737; found: 591.21777.

3. Results and Discussion

To introduce the fluorine atom into the aza-BODIPY core, 4-fluorostyrene was selected as a starting material (Scheme 1). 1-Fluoro-4-(1,2-dibromoethyl)benzene was obtained in 80% yield by the reaction of 4-fluorostyrene with Br₂ at 0 °C. Followed by NaN₃, DMSO, NaOH and reflux in toluene, the yellow oil 3-(4-fluorophenyl)-2*H*-azirine can be synthesized in 60% yield. Acetophenone was employed to react with 3-(4-fluorophenyl)-2*H*-azirine to give fluoro-containing pyrroles **1** (y. 55%) and **2** (y. 60%) in the presence of NaH (Scheme 1), respectively [19]. The clear 1:10 integral areas of the hydrogen signals in pyrrole **2** between the proton of N-*H* (δ = 8.34 (br s, 1H)) and other protons of the pyrrole in the ¹H NMR spectrum were obtained (see ESI), indicating that the fluorine atom was successfully introduced into the pyrrole skeleton.

<Scheme 1>

<Scheme 2>

Using the pyrroles **1** and **2**, symmetric/asymmetric fluorine-containing aza-BODIPYs **3-6** were successfully synthesized in moderate yields under HOAc, Ac₂O and NaNO₂, followed by complexation with Et₃N-BF₃·Et₂O, as described in the literature (Scheme 2) [19]. Their structures were unambiguously confirmed by detailed characterizations including NMR and ESI-HRMS analyses (see ESI).

Furthermore, the solid-state structure of aza-BODIPY **4** was clearly confirmed by single crystal X-ray analysis (Fig. 2 and see ESI). Selected bond lengths and angles are summarized in Fig. 2. The F2-C15 distances (1.278 Å) were found to be slightly shorter than the F-C_{ph} bond of the reported fluoride (1.365 Å) by 0.09 Å [11c]. The other bond lengths for **4** are very similar to those found for **8** (R₁ = H, R₂ = OMe) (Fig. 1b) [13]. Additionally, the *sp*³ hybridized boron center for **4** appears as a slightly distorted tetrahedron with angles. The dihedral angles C(11*)-N(2)-C(11)-C(10) of 179.9° are nearly of the ideal value of 180°, thus indicating the plane skeleton of the aza-BODIPY core.

<Fig. 2>

Additionally, the derivative of the fluorine-containing aza-BODIPYs was found to be feasible (Scheme 3). NaN₃ undertook nucleophilic aromatic substitution (S_NAr) via replacement of the fluorine atoms of **4** to form N₃-containing aza-BODIPY **9** by treatment of water, whereas it provides NH₂-containing aza-BODIPY **10** by treatment of NaHS (Scheme 3) [20]. The proton of N-*H*₂ (δ = 4.04 (br s, 2H)) and other distinct hydrogen signals for the asymmetric NH₂-containing aza-BODIPY **10** in the ¹H NMR spectrum indicate clearly that the amino group was introduced into the aza-BODIPY **10** (see ESI).

<Scheme 3>

<Table 1>

The absorption and emission spectra of aza-BODIPYs **3-6** are shown in Fig. 3 whereas the photophysical data collected are listed in Table 1. Aza-BODIPYs **3-6**

have excellent optical properties. These dyes absorb and emit in the NIR region, with high molar extinction coefficients, high quantum yields and narrow full width at half maxima (FWHM). The profile curves including absorption and emission between **3/4** and **5/6** nearly overlap, and optical properties of every pair of aza-BODIPYs are very similar. Moreover, the pictures of **3-6** taken in CHCl_3 under normal conditions reveal vivid colors, and the hues between pairs of **3/4** and **5/6** are also uniform (Fig. 3). To investigate the spectroscopic effects of the withdrawing group F atom, the spectral characteristics of **3-6** were respectively compared to those of the known dyes **7** ($R_1 = R_2 = \text{H}$) [13] and **8** [13] (Fig. 1b and Table 1). By comparison with the absorption and emission maxima of **7** ($\lambda_{\text{abs}} = 650$, $\lambda_{\text{em}} = 672$ nm), it is clear that the effect of the fluorine atom in **3** and **5** is trifling. Moreover, extinction coefficients and quantum yields of dyes **3** and **5** in CHCl_3 are nearly the same to those of dye **7** ($79000 \text{ M}^{-1} \text{ cm}^{-1}$, $\Phi_f = 0.34$). In addition, the photophysical properties of mono/bis-fluoro substituted aza-BODIPYs **4** and **6** bearing the $-\text{OMe}$ group are also completely equivalent to those of aza-BODIPY **8**.

<Fig. 3>

Interestingly, the replacement of -F atoms with $-\text{N}_3$ groups in **9** ($\lambda_{\text{abs}} = 710$ nm) resulted in a remarkable bathochromic shift (22 nm) compared to that of the parent aza-BODIPY **4** ($\lambda_{\text{abs}} = 688$) (Fig. 4 and Table 1), which is well in agreement with the reported literature [9]. Although the electronegativity of N_3 group (-3.04) was well-known to be smaller than that of F atom (-4.0) [18], the effect of the N_3 group at 1/7 position to photophysical properties in aza-BODIPY **9** is notable (Fig. 4 and Table 1), comparing to F-containing aza-BODIPYs **3-6** as described above. Aza-BODIPY **9** emits at 754 nm, possesses large Stokes' shift (44 nm) and has moderate quantum yield ($\Phi_f = 0.26$). The absorption and emission maxima of **10** bearing the NH_2 -donating group ($\lambda_{\text{abs}} = 699$ nm, $\lambda_{\text{em}} = 749$ nm) was obviously red shifted compared with **4** ($\lambda_{\text{abs}} = 688$ nm, $\lambda_{\text{em}} = 715$ nm), and the fluorescence of **10** became so weak (0.03) due to the intramolecular charge transfer (ICT) effect.

<Fig. 4>

Since having a $-\text{NH}_2$ group in **10**, therefore, we continued to explore the response of the amino-containing aza-BODIPY **10** to pH value. Dimethylamino group ($-\text{NMe}_2$) is well-known to be one of the fragments for the pH-responsive fluorescent probe frequently used for the purpose of the ICT effect.[21] In this ICT process, the excited state of the fluorophore can be quenched by the electron transfer from electron donating amine to the fluorophore; upon recognition of a proton, the electron transfer is "switched off" and in turn the emission of fluorescence is "switched on". Herein, the mechanism of aza-BODIPY **10** bearing a $-\text{NH}_2$ group to pH value is same to that of the above description. Upon addition of hydrochloric acid to **10** with a $-\text{NH}_2$ group

as the pH-sensitive functionality, **10** was protonated at relatively low pH value [8,22], and a distinctly hypochromatic shift occurs at pH 2 (Fig. 5a). A stepwise increase of the absorption intensity was observed in the 700 nm band of **10-H**⁺ from the 710 nm of **10** in DMSO/H₂O (2:1, v/v) (Fig. 5a). The absorption intensity of **10-H**⁺ reached the maximum when 4 M HCl was used. The absorption band of **10-H**⁺ is blue-shifted by about 10 nm compared to that of **10**. Furthermore, with decreasing pH, a remarkable increase in fluorescence intensity at 750 nm by about 6 folds when **10** was treated with 4M HCl. (Fig. 5b). So, aza-BODIPY **10** could be used as a turn-on fluorescent probe for pH value.

<Fig. 5>

Quantum chemical calculations were performed on the four fluorine-containing aza-BODIPY compounds **3–6** to support the absorption and emission results. We followed a strategy that was developed recently (see computational details) [23] and that gave excellent result for both the BODIPY and aza-BODIPY systems [24]. The main computed photophysical data at the best level of calculation, i.e. cLR-PCM(CHCl₃)-M06-2X/6-311+G(2d,p)/SOS-CIS(D) are summarized in Table 2. The calculations confirmed the limited effect of adding a F atom at 1/7-positions, comparing for instance the calculated absorption and emission wavelengths of **4** and **6** with respect to those of **8**. This observation persists for the absorption/fluorescence crossing point energies (E^{AFCP}) that can be compared to experimental E^{0-0} energies. Fig. 6 shows the density difference plots computed for the fluorine-containing aza-BODIPY dyes and the involvement of the fluorine atom is clearly limited in the excited-states, which is consistent with the measurements. CT parameters, i.e. the CT distance (d^{CT}) and the transferred charge (q^{CT}) were also calculated through Le Bahers' model (Table 2) [25]. Again the only substantial difference emerges when comparing CT distances of parent complexes with and without the methoxy group (**3** vs. **4** for instance). Similar calculations were realized on compounds **9** and **10** (Table 2 and Fig. 6) confirming the tendencies experimentally observed, such as the redshift of the absorption/emission when going from **3** and **4** to **9** and **10**. Replacing F by N₃ (**4** vs. **9**) slightly reduces the CT distance in agreement with the smaller electronegativity of N₃ with respect to F. Finally, the small calculated d^{CT} of **10** can be attributed to the strong dissymmetry induced by the replacement of only one fluorine group by NH₂ donating group.

<Table 2>

<Fig. 6>

4. Conclusions

Utilizing 4-fluorostyrene as starting material, acetophenone was employed to react with 3-(4-fluorophenyl)-2*H*-azirine to give fluoro-containing pyrroles in the presence

of NaH. Fluorine-containing aza-BODIPYs **3-6** at 1/7-position were successfully synthesized by the reaction of fluoro-containing pyrroles **1** and **2** under HOAc, Ac₂O and NaNO₂, followed by complexation with Et₃N-BF₃ · Et₂O. NaN₃ undertook nucleophilic aromatic substitution to form N₃-containing aza-BODIPY **9** by treatment of water, whereas it gives NH₂-containing aza-BODIPY **10** by treatment of NaHS. The solid-state structure of aza-BODIPY **4** was confirmed by single crystal X-ray analysis, the F-C_{Ph} bond distance of which was found to be shorter than those of the reported fluoride. The optical properties of fluorine-containing aza-BODIPYs **3-6** are very similar to those of the parent complexes. Surprisingly, the effect of the withdrawing -F group at 1/7-position on the photophysical properties is found to be almost negligible, comparing to those of the parent aza-BODIPYs **7** and **8**. However, the replacement of -F atoms with -N₃/NH₂ groups in **9** and **10** resulted in a remarkable bathochromic shift. The amino-containing aza-BODIPY **10** could be used as a turn-on fluorescent probe for pH value. The main photophysical data are well supported and explained by cLR-PCM(CHCl₃)-M06-2X/6-311+G(2d,p)/SOS-CIS(D) calculations.

Acknowledgements

This work was supported by NNSFC (21542004), the Program for Liaoning Excellent Talents in University (LJQ2015087), Liaoning BaiQianWan Talents Program (2015-56), the State Key Laboratory of Fine Chemicals (KF1506), the Public Research Foundation of Liaoning Province for the Cause of Science (2014003009), the Scientific Research Foundation for the Returned Overseas Chinese Scholars, Liaoning Province Natural Science Foundation (2014020139), and the start-up funds from Shenyang University of Chemical Technology. D.J. acknowledges the European Research Council (ERC) and the Région des Pays de la Loire for financial support in the framework of a Starting Grant (Marches-278845) and a recrutement sur poste stratégique, respectively. This research used resources of (i) the GENCI-CINES/IDRIS, (ii) the CCIPL (Centre de Calcul Intensif des Pays de Loire), and (iii) the Troy cluster in Nantes.

References

- [1] (a) Weissleder R. A clearer vision for in vivo imaging. *Nat Biotechnol* 2001;19:316.
- (b) Tung CH, Lin Y, Moon WK, Weissleder R. A receptor-targeted near-infrared fluorescence probe for in vivo tumor imaging. *ChemBioChem* 2002;3:784.
- (c) Kim HN, Ren W, Kim JS, Yoon J. Fluorescent and colorimetric sensors for detection of lead, cadmium, and mercury ions. *Chem Soc Rev* 2012;41:3210.
- (d) Joseph R, Rao CP. Ion and molecular recognition by lower rim 1,3-diconjugates of calix[4]arene as receptors. *Chem Rev* 2011;111:4658.
- (e) Liu Z, He W, Guo Z. Metal coordination in photoluminescent sensing. *Chem*

- Soc Rev 2013;42:1568.
- (f) Ding Y, Tang Y, Zhu W, Xie Y. Fluorescent and colorimetric ion probes based on conjugated oligopyrroles. *Chem Soc Rev* 2015;44:1101.
- [2] (a) Weissleder R, Tung CH, Mahmood U, Bogdanov A. In vivo imaging of tumors with protease-activated near-infrared fluorescent probes. *Nat Biotechnol* 1999;17:375.
- (b) Becker A, Hennesius C, Licha K, Ebert B, Sukowski U, Semmler W, et al. Receptor-targeted optical imaging of tumors with near-infrared fluorescent ligands. *Nat Biotechnol* 2001;19:327.
- [3] (a) Li Y, Patrick BO, Dolphin D. Near-Infrared absorbing azo dyes: synthesis and X-ray crystallographic and spectral characterization of monoazopyrroles, bisazopyrroles, and a boron-azopyrrole complex. *J Org Chem* 2009;74:5237.
- (b) Descalzo AB, Rurack K. On the signalling pathways and Cu(II)-mediated anion indication of N-meso-substituted heptamethine cyanine dyes. *Chem Eur J* 2009;15:3173.
- (c) Kubo Y, Yoshida K, Adachi M, Nakamura S, Maeda S. Experimental and theoretical study of near-infrared absorbing naphthoquinone methide dyes with a nonplanar geometry. *J Am Chem Soc* 1991;113:2868.
- (d) Barthram AM, Cleary RL, Kowallick R, Ward MD. A new redox-tunable near-IR dye based on a trinuclear ruthenium(II) complex of hexahydroxytriphenylene. *Chem Commun* 1998;2695.
- (e) Hairong L, Ngan N, Frank RF. Syntheses and properties of octa-, tetra-, and di-hydroxy-substituted phthalocyanines. *Tetrahedron* 2009;65:3357.
- (f) Fan J, Dong H, Hu M, Wang J, Zhang H, Zhu H, et al. Fluorescence imaging lysosomal changes during cell division and apoptosis observed using Nile Blue based near-infrared emission. *Chem Commun* 2014;50:882.
- (g) Zhao C, Zhang J, Wang X, Zhang Y. Pyridone fused boron-dipyrromethenes: synthesis and properties. *Org Biomol Chem* 2013;11:372.
- (h) Miao JT, Fan C, Shi XY, Sun R, Xu YJ, Ge JF. Colorimetric and ratiometric pH responses by the protonation of phenolate within hemicyanine. *Analyst* 2014;139:6290.
- [4] (a) Rohrer H, Hofmann W. Versuche zur aktiven Immunisierung bei Maul- und Klauenseuche mit Mäusegehirnpassagevirus vom B-Typ. *Nature* 1943;151:504.
- (b) Loudet A, Burgess K. BODIPY dyes and their derivatives: syntheses and spectroscopic properties. *Chem Rev* 2007;107:4891.
- (c) Bessette A, Hanan GS. Design, synthesis and photophysical studies of dipyrromethene-based materials: insights into their applications in organic photovoltaic devices. *Chem Soc Rev* 2014;43:3342.
- (d) Jiao L, Wu Y, Ding Y, Wang S, Zhang P, Yu C, et al. Conformationally restricted aza-dipyrromethene boron difluorides (aza-BODIPYs) with high fluorescent quantum yields. *Chem Asian J* 2014;9:805.
- (e) Killoran J, O'Shea DF. Impact of a conformationally restricted receptor on the BF₂ chelated azadipyrromethene fluorosensing platform. *Chem Commun* 2006;1503.

- (f) McDonnell SO, Hall MJ, Allen LT, Byrne A, Gallagher WM, O'Shea DF. Supramolecular Photonic Therapeutic Agents. *J Am Chem Soc* 2005;127:16360.
- (g) Hall MJ, Allen LT, O'Shea DF. PET modulated fluorescent sensing from the BF₂ chelated azadipyrrromethene platform. *Org Biomol Chem* 2006;4:776.
- (h) Adarsh N, Krishnan MS, Ramaiah D. Sensitive naked eye detection of hydrogen sulfide and nitric oxide by aza-BODIPY dyes in aqueous medium. *Anal Chem* 2014;86:9335.
- [5] (a) Umezawa K, Nakamura Y, Makino H, Citterio D, Suzuki K. Bright, color-tunable fluorescent dyes in the visible-near-infrared region. *J Am Chem Soc* 2008;130:1550.
- (b) Umezawa K, Matsui A, Nakamura Y, Citterio D, Suzuki K. Bright, color-tunable fluorescent dyes in the Vis/NIR region: establishment of new "tailor-made" multicolor fluorophores based on borondipyrrromethene. *Chem Eur J* 2009;15:1096.
- (c) Awuah SG, Polreis J, Biradar V, You Y. Singlet oxygen generation by novel NIR BODIPY dyes. *Org Lett* 2011;13:3884.
- (d) Wang YW, Descalzo AB, Shen Z, You XZ, Rurack K. Dihydronaphthalene-fused boron-dipyrrromethene (BODIPY) dyes: insight into the electronic and conformational tuning modes of BODIPY fluorophores. *Chem Eur J* 2010;16:2887.
- (e) Kowada T, Yamaguchi S, Ohe K. Highly fluorescent BODIPY dyes modulated with spirofluorene moieties. *Org Lett* 2010;12:296.
- (f) Heyer E, Retailleau P, Ziessel R. α -Fused dithienyl BODIPYs synthesized by oxidative ring closure. *Org Lett* 2014;16:2330.
- (g) Wu Y, Cheng C, Jiao L, Yu C, Wang S, Hao E, et al. β -Thiophene-fused BF₂-azadipyrrromethenes as near-infrared dyes. *Org Lett* 2014;16:748.
- (h) Goze C, Ulrich G, Ziessel R. Tetrahedral boron chemistry for the preparation of highly efficient "cascadelle" devices. *J Org Chem* 2007;72:313.
- (i) Davies LH, Stewart B, Harrington RW, Clegg W, Higham LJ. Air-stable, highly fluorescent primary phosphanes. *Angew Chem Int Ed* 2012;51:4921.
- (j) Le Guennic B, Maury O, Jacquemin D. Aza-boron-dipyrrromethene dyes: TD-DFT benchmarks, spectral analysis and design of original near-IR structures. *Phys Chem Chem Phys* 2012;14:157.
- (k) Zhao W, Carreira EM. Conformationally restricted aza-bodipy: a highly fluorescent, stable, near-infrared-absorbing dye. *Angew Chem Int Ed* 2005;44:1677.
- (l) Zhao W, Carreira EM. Conformationally restricted aza-BODIPY: highly fluorescent, stable near-infrared absorbing dyes. *Chem Eur J* 2006;12:7254.
- (m) Huang L, Cui X, Therrien B, Zhao J. Energy-funneling-based broadband visible-light-absorbing bodipy-C₆₀ triads and tetrads as dual functional heavy-atom-free organic triplet photosensitizers for photocatalytic organic reactions. *Chem Eur J* 2013;19:17472.

- [6] Killoran J, Allen L, Gallagher JF, Gallagher WM, O'Shea DF. Synthesis of BF₂ chelates of tetraarylazadipyrromethenes and evidence for their photodynamic therapeutic behaviour. *Chem Commun* 2002;1862.
- [7] (a) Lu H, Mack J, Yang Y, Shen Z. Structural modification strategies for the rational design of red/NIR region BODIPYs. *Chem Soc Rev* 2014;43:4778.
 (b) Bellier Q, Pégaz S, Aronica C, Le Guennic B, Andraud C, Maury O. Near-Infrared Nitrofluorene Substituted Aza-Boron-dipyrromethenes Dyes. *Org Lett* 2011;13:22.
- [8] McDonnell SO, O'Shea DF. Near-infrared sensing properties of dimethylamino-substituted BF₂-azadipyrromethenes. *Org Lett* 2006;8:3493.
- [9] Jiao L, Wu Y, Wang S, Hu X, Hao E, Vicente MG, et al. Accessing near-infrared-absorbing BF₂-azadipyrromethenes via a push-pull effect. *J Org Chem* 2014;79:1830.
- [10] Valero G, Companyó X, Rios R. Enantioselective organocatalytic synthesis of fluorinated molecules. *Chem Eur J* 2011;17:2018.
- [11](a) Jiang XD, Kakuda K, Matsukawa S, Yamamichi H, Kojima S, Yamamoto Y. Synthesis and application of a bidentate ligand based on decafluoro-3-phenyl-3-pentanol: steric effect of pentafluoroethyl groups on the stereomutation of O-equatorial C-apical spirophosphoranes. *Chem Asian J* 2007;2:314.
 (b) Jiang XD, Matsukawa S, Yamamichi H, Yamamoto Y. First isolation and kinetic study of hypervalent 10-As-5 organoarsenic compounds with a C-apical, O-equatorial configuration. *Inorg Chem* 2007;46:5480.
 (c) Jiang XD, Matsukawa S, Yamamoto Y. Synthesis, structure and isomerization of arylphosphoranes with anti-apicophilic bonding modes using a novel bidentate ligand with two C₂F₅ groups. *Dalton Trans* 2008;28:3678.
 (d) Jiang XD, Matsukawa S, Yamamichi H, Kakuda K, Kojima S, Yamamoto Y. Stereomutation and Experimental Determination of the Relative Stability of Diastereomeric O-Equatorial Anti-Apicophilic Spirophosphoranes. *Eur J Org Chem* 2008;1392.
 (e) Jiang XD, Matsukawa S, Fukuzaki Y, Yamamoto Y. Formation of epoxides from pentacoordinated organoarsenic compounds with a β -hydroxyethyl group. *New J Chem* 2010;34:1623.
 (f) Jiang XD, Matsukawa S, Kakuda K, Fukuzaki Y, Kojima S, Yamamoto Y, et al. Efficient synthesis of tetradecafluoro-4-phenylheptan-4-ol by a Cannizzaro-type reaction and application of the alcohol as a bulky Martin ligand variant for a new anti-apicophilic phosphorane. *Dalton Trans* 2010;39:9823.
 (g) Jiang XD, Matsukawa S, Kojima S, Yamamoto Y. Synthesis and characterization of antiapicophilic arsoranes and related compounds. *Inorg Chem* 2012;51:10996.
- [12] (a) Jiang XD, Zhang J, Furuyama T, Zhao W. Development of mono- and di-AcO substituted BODIPYs on the boron center. *Org Lett* 2012;14:248.
 (b) Jiang XD, Zhang H, Zhang Y, Zhao W. Development of non-symmetric thiophene-fused BODIPYs. *Tetrahedron* 2012;68:9795.

- (c) Jiang XD, Zhang J, Shao X, Zhao W. A selective fluorescent turn-on NIR probe for cysteine. *Org Biomol Chem* 2012;10:1966.
- (d) Kang R, Shao X, Peng F, Zhang Y, Sun GT, Jiang XD, et al. A highly selective turn-on fluorescent chemosensor for Al^{3+} imaging in living cells via through-bond energy Transfer. *RSC Adv* 2013;3:21033.
- (e) Jiang XD, Xi D, Sun C, Guan J, He M, Xiao LJ. Synthesis of a pyrene-fused aza-BODIPY as a near-infrared dye having the absorption maximum at 746 nm. *Tetrahedron Lett* **2015**;56:4868.
- (f) Jiang XD, Zhao J, Xi D, Yu H, Guan J, Xiao LJ, et al. A new water-soluble phosphorus-dipyrromethene and phosphorus-azadipyrromethene dye: PODIPY/aza-PODIPY. *Chem Eur J* 2015;21:6079.
- (g) Jiang XD, Li S, Guan J, Fang T, Liu X, Xiao LJ. Recent advances of the near-infrared fluorescent aza-BODIPY dyes. *Curr Org Chem* 2016;20:1736.
- (h) Jiang XD, Guan J, Li Q, Sun C. New near-infrared fluorescent aza-BODIPY dyes with 1-methyl-1H-pyrrolyl substituents at 3,5-positions. *Asian J Org Chem* 2016;5:1063.
- [13] Gorman A, Killoran J, O'Shea C, Kenna T, Gallagher WM, O'Shea DF. In Vitro Demonstration of the Heavy-Atom Effect for Photodynamic Therapy. *J Am Chem Soc* 2004;126:10619.
- [14] Frisch MJ, Trucks GW, Schlegel HB, Scuseria GE, Robb MA, Cheeseman JR, et al. Gaussian 09W, revision A.1; Gaussian Inc.: Wallingford, CT, 2009.
- [15](a) Jacquemin D, Planchat A, Adamo C, Mennucci B. TD-DFT Assessment of Functionals for Optical 0-0 Transitions in Solvated Dyes. *J Chem Theory Comput* 2012;8:2359.
- (b) Zhao Y, Truhlar DG. The M06 suite of density functionals for main group thermochemistry, thermochemical kinetics, noncovalent interactions, excited states, and transition elements: two new functionals and systematic testing of four M06-class functionals and 12 other functionals. *Theor Chem Acc* 2008;120:215.
- (c) Li R, Zheng J, Truhlar DG. Density functional approximations for charge transfer excitations with intermediate spatial overlap. *Phys Chem Chem Phys* 2010;12:12697.
- (d) Isegawa M, Peverati R, Truhlar DG. Performance of recent and high-performance approximate density functionals for time-dependent density functional theory calculations of valence and Rydberg electronic transition energies. *J Chem Phys* 2012;137:244104.
- (e) Leang SS, Zahariev F, Gordon MS. Benchmarking the performance of time-dependent density functional methods. *J Chem Phys* 2012;136:104101.
- (f) Charaf-Eddin A, Planchat A, Mennucci B, Adamo C, Jacquemin D. Choosing a Functional for Computing Absorption and Fluorescence Band Shapes with TD-DFT. *J Chem Theory Comput* 2013;9:2749.
- [16] Cossi M, Barone V. Time-dependent density functional theory for molecules in liquid solutions. *J Chem Phys* 2001;115:4708.

- [17] Impropa R, Barone V, Scalmani G, Frisch MJ. A state-specific polarizable continuum model time dependent density functional theory method for excited state calculations in solution. *J Chem Phys* 2006;125:054103.
- [18] (a) Sanderson RT. Polar Covalence, New York: Academic Press, 1983, 37.
(b) Bratsch SG. A group electronegativity method with Pauling units. *J Chem Educ* 1985;62:101.
- [19](a) Jiang XD, Gao R, Yue Y, Sun GT, Zhao W. A NIR BODIPY dye bearing 3,4,4a-trihydroxanthene moieties. *Org Biomol Chem* 2012;10:6861.
(b) Jiang XD, Xi D, Zhao J, Yu H, Sun GT, Xiao LJ. A styryl-containing aza-BODIPY as a near-infrared dye. *RSC Adv* 2014;4:60970.
(c) Jiang XD, Zhao J, Li Q, Sun CL, Guan J, Xiao LJ, et al. Synthesis of NIR fluorescent thienyl-containing aza-BODIPY and its application for detection of Hg^{2+} : Electron transfer by bonding with Hg^{2+} . *Dyes Pigm* 2016;125:136.
- [20] Lippert AR, New EJ, Chang CJ. Reaction-based fluorescent probes for selective imaging of hydrogen sulfide in living cells. *J Am Chem Soc* 2011;133:10078.
- [21](a) Thivierge C, Han J, Jenkins RM, Burgess K. Fluorescent Proton Sensors Based on Energy Transfer. *J Org Chem* 2011;76:5219.
(b) Czarnik AW. Chemical Communication in Water Using Fluorescent Chemosensors. *Acc Chem Res* 1994;27:302.
(c) Jager WF, Hammink TS, van den Berg O, Grozema FC. Highly Sensitive Water-Soluble Fluorescent pH Sensors Based on the 7-Amino-1-methylquinolinium Chromophore. *J Org Chem* 2010;75:2169.
- [22] Rurack K, Kollmannsberger M, Daub J. Molecular Switching in the Near Infrared (NIR) with a Functionalized Boron–Dipyrromethene Dye. *Angew Chem Int Ed* 2001;40:385.
- [23] Chibani S, Laurent AD, Le Guennic B, Jacquemin D. Improving the Accuracy of Excited-State Simulations of BODIPY and Aza-BODIPY Dyes with a Joint SOS-CIS(D) and TD-DFT Approach. *J Chem Theory Comput* 2014;10:4574.
- [24] Le Guennic B, Jacquemin D. Taking up the cyanine challenge with quantum tools. *Acc Chem Res* 2015;48:530.
- [25](a) Le Bahers T, Adamo C, Ciofini I. A Qualitative Index of Spatial Extent in Charge-Transfer Excitations. *J Chem Theory Comput* 2011;7:2498.
(b) Jacquemin D, Le Bahers T, Adamo C, Ciofini I. What is the "best" atomic charge model to describe through-space charge-transfer excitations? *Phys Chem Chem Phys* 2012;14:5383.

Fig. 1. (a) BODIPY vs. aza-BODIPY. (b) Structures of some previously reported aza-BODIPYs [6,8,9].

Scheme 1 Synthesis of fluorine-containing pyrroles **1** and **2**.

Scheme 2 Synthesis of fluorine-containing aza-BODIPYs **3-6**.

Fig. 2. ORTEP diagram of **4** showing the thermal ellipsoids at the 30% probability level. All hydrogen atoms are omitted for clarity. Selected bond lengths (Å) and angles (°): B1-F1, 1.349(4); B1-N1, 1.518(5); N1-C8, 1.344(5); N1-C11, 1.365(5); N2-C11, 1.301(5); F2-C15, 1.278(9); C(11*)-N(2)-C(11)-C(10), 179.9(5).

Scheme 3 Synthesis of aza-BODIPYs **9** and **10**.

Table 1 Photophysical properties of aza-BODIPYs **3-10** in CHCl₃ at 298 K.

Fig. 3. (a) Normalized absorption and (b) fluorescence spectra of aza-BODIPYs **3** (black curve), **4** (blue curve), **5** (red curve) and **6** (green curve) in CHCl₃ at 298 K. Photoimages of **3-6** under normal room illumination in CHCl₃ at 298 K.

Fig. 4. (a) Normalized absorption and (b) fluorescence spectra of aza-BODIPYs **9** (red curve) and **10** (black curve) in CHCl₃ at 298 K.

Fig. 5. Absorption (a) and fluorescence (b) spectra (pH 7, 6, 5, 4, 3, 2, 1.8, 1.6, 1.4, 1.2, 1 and 1 M, 2 M, 4 M, 6 M of HCl) of 2 μM dye **10** in DMSO/H₂O (2:1, v/v) as a function of pH. $\lambda_{\text{ex}} = 700 \text{ nm}$.

Table 2 Computed λ_{abs} and λ_{em} (in nm), AFCP energy (E^{AFCP} , in nm), charge transfer distances (d^{CT} , in Å), and transferred charges (q^{CT} , in e) of aza-BODIPYs **3**, **4**, **5**, **6**, **8**, **9** and **10** at the cLR-PCM(CHCl₃)-M06-2X/6-311+G(2d,p)/SOS-CIS(D).

Fig. 6. Density difference plots for (from left to right and top to bottom) aza-BODIPYs **3**, **4**, **5**, **6**, **9** and **10**. The blue (red) zones indicate density decrease (increase) upon transition.

Fig. 1 (a) BODIPY vs. aza-BODIPY. (b) Structures of some previously reported aza-BODIPYs [6,8,9].

Scheme 1 Synthesis of fluorine-containing pyrroles **1** and **2**.

Scheme 2 Synthesis of fluorine-containing aza-BODIPYs 3-6.

Fig. 2. ORTEP diagram of **4** showing the thermal ellipsoids at the 30% probability level. All hydrogen atoms are omitted for clarity. Selected bond lengths (Å) and angles (°): B1-F1, 1.349(4); B1-N1, 1.518(5); N1-C8, 1.344(5); N1-C11, 1.365(5); N2-C11, 1.301(5); F2-C15, 1.278(9); C(11*)-N(2)-C(11)-C(10), 179.9(5).

Scheme 3 Synthesis of aza-BODIPYs **9** and **10**.

Table 1 Photophysical properties of aza-BODIPYs **3-10** in CHCl₃ at 298 K.

Dye	$\lambda_{\text{abs}}/\lambda_{\text{em}}$ [nm]	Stokes' shift [cm ⁻¹]	FWHM [nm]	ϵ [M ⁻¹ cm ⁻¹]	Φ_f
3	650/674	25	49	80000	0.34
4	688/715	27	55	86000	0.36
5	650/672	25	49	79000	0.34
6	688/715	27	55	85000	0.36
7 [13]	650/672	25	49	79000	0.34
8 [13]	688/715	27	55	85000	0.36
9	710/754	44	86	67000	0.26
10	699/749	50	68	52000	0.03

Fig. 3. (a) Normalized absorption and (b) fluorescence spectra of aza-BODIPYs **3** (black curve), **4** (blue curve), **5** (red curve) and **6** (green curve) in CHCl_3 at 298 K. Photoimages of **3-6** under normal room illumination in CHCl_3 at 298 K.

Fig. 4. (a) Normalized absorption and (b) fluorescence spectra of aza-BODIPYs **9** (red curve) and **10** (black curve) in CHCl_3 at 298 K.

Fig. 5. Absorption (a) and fluorescence (b) spectra (pH 7, 6, 5, 4, 3, 2, 1.8, 1.6, 1.4, 1.2, 1 and 1 M, 2 M, 4 M, 6 M of HCl) of 3 μ M dye **10** in DMSO/H₂O (2:1, v/v) as a function of pH. $\lambda_{\text{ex}} = 700$ nm.

Table 2 Computed λ_{abs} and λ_{em} (in nm), AFCP energy (E^{AFCP} , in nm), charge transfer distances (d^{CT} , in Å), and transferred charges (q^{CT} , in e) of aza-BODIPYs **3**, **4**, **5**, **6**, **8**, **9** and **10** at the cLR-PCM(CHCl_3)-M06-2X/6-311+G(2d,p)/SOS-CIS(D).

Dye	λ_{abs} [nm]	λ_{em} [nm]	E^{AFCP} [nm]	d^{CT} [Å]	q^{CT} [e]
3	664	717	706	0.56	0.34
4	704	770	755	1.44	0.39
5	664	717	710	0.61	0.34
6	703	769	752	1.44	0.39
8	702	768	752	1.47	0.39
9	714	783	769	1.11	0.38
10	716	798	770	0.56	0.39

Fig. 6. Density difference plots for (from left to right and top to bottom) aza-BODIPYs **3**, **4**, **5**, **6**, **9** and **10**. The blue (red) zones indicate density decrease (increase) upon transition.

ESI

**Synthesis, structure and photophysical properties of NIR
aza-BODIPYs with –F/–N₃/–NH₂ groups at 1,7-positions**

**Xin-Dong Jiang,^{a,b,*} Jian Guan,^a Jiuli zhao,^a Boris Le Guennic,^{c,*} Denis
Jacquemin,^d Zhigang Zhang^a Shangdong Chen^a and Linjiu Xiao^a**

1 NMR

2 HRMS

3 X-ray single crystal diffraction data of **4**

4 Ground state geometries in gas phase

1. NMR

¹H NMR of 3-(4-fluorophenyl)-2H-azirine

^1H NMR of pyrrole 1

¹H NMR of pyrrole 2

¹H NMR of aza-BODIPY 3

^{13}C NMR of aza-BODIPY 3

¹H NMR of aza-BODIPY 4

^{13}C NMR of aza-BODIPY 4

¹H NMR of aza-BODIPY 5

¹³C NMR of aza-BODIPY 5

¹H NMR of aza-BODIPY 6

^{13}C NMR of aza-BODIPY 6

¹H NMR of aza-BODIPY 9

¹H NMR of aza-BODIPY 10

2. HRMS

使用仪器：

Thermo Scientific LTQ Orbitrap XL

离子源：ESI

检测结果：

1. HRMS (ESI) m/z calcd for $C_{16}H_{13}FN^+$ ($M+H$) $^+$ 238.10265, found 238.10268.

HRMS for 1

2. HRMS (ESI) m/z calcd for $C_{17}H_{15}FNO^+$ ($M+H$) $^+$ 268.11322, found 268.11322.

HRMS for 2

Mass Spectrum Molecular Formula Report

Analysis Info D:\Data\20150115CEYANG\2.d
Analysis Name tune_wide_pos.m
Method 2
Sample Name Comment
Acquisition Date 1/15/2015 11:13:26 AM
Operator microTOF-Q
Instrument / Ser# Bruker Customer 125

Acquisition Parameter

Source Type	ESI	Ion Polarity	Positive	Set Nebulizer	0.3 Bar
Focus	Active	Set Capillary	4500 V	Set Dry Heater	180 °C
Scan Begin	50 m/z	Set End Plate Offset	-500 V	Set Dry Gas	4.0 l/min
Scan End	3000 m/z	Set Collision Cell RF	600.0 Vpp	Set Divert Valve	Source

Generate Molecular Formula Parameter

Formula, min.	C32H20B1F4N3Na				
Formula, max.					
Measured m/z	556.158	Tolerance	5 ppm	Charge	1
Check Valence	no	Minimum	0	Maximum	0
Nitrogen Rule	no	Electron Configuration	both		
Filter H/C Ratio	no	Minimum	0	Maximum	3
Estimate Carbon	yes

3. HRMS (ESI) m/z calcd for $C_{34}H_{25}BF_4N_3O_2^+$ (M+H) $^+$ 594.19705, found 594.19678.

001#23 RT: 0.67 AV: 1 NL: 1.87E4
T: FTMS + p ESI Full ms [100.00-1050.00]

HRMS for 4**4. HRMS (ESI) m/z calcd for $C_{32}H_{22}BF_3N_3^+$ (M+H) $^+$ 516.18534, found 516.18463.**

002#39 RT: 1.15 AV: 1 NL: 1.53E4
T: FTMS + p ESI Full ms [100.00-1050.00]

HRMS for 5

Mass Spectrum Molecular Formula Report

Analysis Info D:\Data\20150115CEYANG\3.d

Analysis Name

Acquisition Date 1/15/2015 11:16:31 AM

Method tune_wide_pos.m

Operator micrOTOF-Q

Sample Name 3

Instrument / Ser# Bruker Customer 125

Comment

Acquisition Parameter

Source Type	ESI	Ion Polarity	Positive	Set Nebulizer	0.3 Bar
Focus	Active	Set Capillary	4500 V	Set Dry Heater	180 °C
Scan Begin	50 m/z	Set End Plate Offset	-500 V	Set Dry Gas	4.0 l/min
Scan End	3000 m/z	Set Collision Cell RF	600.0 Vpp	Set Divert Valve	Source

Generate Molecular Formula Parameter

Formula, min. C34H25B1F3N3O2Na

Formula, max.

Measured m/z 598.189

Tolerance 5 ppm

Charge 1

Check Valence no

Minimum 0

Maximum 0

Nitrogen Rule no

Electron Configuration both

Filter H/C Ratio no

Minimum 0

Maximum 3

Estimate Carbon yes

Bruker Daltonics DataAnalysis 3.4

printed: 1/15/2015 6:55:40 PM

Page 1 of 1

HRMS for 6

5. HRMS (ESI) m/z calcd for $C_{34}H_{29}BF_2N_5O_2^+$ (M+H) $^+$ 588.23769, found 588.23730.

HRMS for 9

6. HRMS (ESI) m/z calcd for $C_{34}H_{27}BF_3N_4O_2^+$ (M+H) $^+$ 591.21737, found 591.21777.

HRMS for 10

3. X-ray single crystal diffraction data of 4

Crystals suitable for the X-ray structural determination were mounted on a Mac Science DIP2030 imaging plate diffractometer and irradiated with graphite monochromated Mo- $K\alpha$ radiation ($\lambda = 0.71073 \text{ \AA}$) for the data collection. The unit cell parameters were determined by separately autoindexing several images in each data set using the DENZO program (MAC Science). For each data set, the rotation images were collected in 3 degree increments with a total rotation of 180 deg about the ϕ axis. The data were processed using SCALEPACK. The structures were solved by a direct method with the SHELX-97 program. Refinement on F^2 was carried out using the full-matrix least-squares by the SHELX-97 program. All non-hydrogen atoms were refined using the anisotropic thermal parameters. The hydrogen atoms were included in the refinement along with the isotropic thermal parameters (Table S1-S5).

Table S1. Crystal data and structure refinement.

Identification code	shelx	
Empirical formula	C ₃₄ H ₂₄ B F ₄ N ₃ O ₂	
Formula weight	593.37	
Temperature	293(2) K	
Wavelength	0.71073 Å	
Crystal system	?	
Space group	Ia-3d	
Unit cell dimensions	a = 16.1853(2) Å	$\alpha = 90^\circ$.
	b = 16.1853(2) Å	$\beta = 90^\circ$.
	c = 19.9274(6) Å	$\gamma = 90^\circ$.
Volume	5220.26(18) Å ³	
Z	48	
Density (calculated)	9.060 Mg/m ³	
Absorption coefficient	0.683 mm ⁻¹	
F(000)	14688	
Crystal size	0.40 x 0.30 x 0.20 mm ³	
Theta range for data collection	3.24 to 28.38°.	
Index ranges	-21 ≤ h ≤ 21, -21 ≤ k ≤ 21, -26 ≤ l ≤ 26	
Reflections collected	23013	
Independent reflections	3276 [R(int) = 0.0867]	
Completeness to theta = 28.38°	99.8 %	
Max. and min. transmission	0.8756 and 0.7719	
Refinement method	Full-matrix least-squares on F ²	
Data / restraints / parameters	3276 / 0 / 202	
Goodness-of-fit on F ²	2.181	
Final R indices [I > 2σ(I)]	R1 = 0.1437, wR2 = 0.3623	
R indices (all data)	R1 = 0.1832, wR2 = 0.4016	
Extinction coefficient	0.0002(6)	
Largest diff. peak and hole	1.077 and -0.724 e.Å ⁻³	

Table S2. Atomic coordinates ($\text{\AA} \times 10^4$) and equivalent isotropic displacement parameters ($\text{\AA}^2 \times 10^3$).
 $U(\text{eq})$ is defined as one third of the trace of the orthogonalized U^{ij} tensor.

	x	y	z	$U(\text{eq})$
F(1)	4356(2)	2252(2)	52(1)	68(1)
N(1)	4724(2)	3202(2)	885(2)	56(1)
N(2)	5000	2500	1895(2)	57(1)
C(11)	4754(3)	3150(3)	1568(2)	57(1)
C(8)	4405(3)	3946(3)	732(2)	58(1)
C(5)	4206(3)	4249(3)	81(2)	59(1)
C(10)	4471(3)	3893(3)	1848(2)	59(1)
O(1)	3564(2)	5205(2)	-1736(2)	89(1)
C(9)	4269(3)	4355(3)	1333(2)	65(1)
C(12)	4440(3)	4104(3)	2551(2)	60(1)
C(2)	3755(3)	4932(3)	-1130(2)	69(1)
C(3)	4209(3)	4242(3)	-1115(2)	72(1)
C(4)	4431(3)	3912(3)	-523(2)	71(1)
C(6)	3730(3)	4966(3)	42(2)	79(1)
C(15)	4374(4)	4560(4)	3832(2)	87(2)
C(13)	4194(3)	3561(3)	3029(3)	79(1)
C(17)	4604(3)	4870(3)	2739(3)	85(2)
C(7)	3504(4)	5296(3)	-541(2)	81(2)
B(1)	5000	2500	432(3)	55(2)
C(16)	4576(4)	5107(3)	3395(3)	90(2)
C(14)	4152(4)	3771(4)	3686(2)	88(2)
F(2)	4452(7)	4791(5)	4443(4)	270(4)
C(1)	3093(6)	5936(5)	-1790(4)	142(3)

Table S3. Bond lengths [Å] and angles [°].

F(1)-B(1)	1.349(4)
N(1)-C(8)	1.344(5)
N(1)-C(11)	1.365(5)
N(1)-B(1)	1.518(5)
N(2)-C(11)	1.301(5)
N(2)-C(11)#1	1.301(5)
C(11)-C(10)	1.403(6)
C(8)-C(9)	1.387(6)
C(8)-C(5)	1.425(6)
C(5)-C(4)	1.370(6)
C(5)-C(6)	1.396(6)
C(10)-C(9)	1.311(6)
C(10)-C(12)	1.442(6)
O(1)-C(2)	1.322(5)
O(1)-C(1)	1.411(8)
C(12)-C(17)	1.322(7)
C(12)-C(13)	1.356(7)
C(2)-C(3)	1.338(7)
C(2)-C(7)	1.376(7)
C(3)-C(4)	1.343(6)
C(6)-C(7)	1.330(6)
C(15)-F(2)	1.278(9)
C(15)-C(16)	1.285(8)
C(15)-C(14)	1.357(8)
C(13)-C(14)	1.355(7)
C(17)-C(16)	1.363(7)
B(1)-F(1)#1	1.349(4)
B(1)-N(1)#1	1.518(5)
C(8)-N(1)-C(11)	107.1(3)
C(8)-N(1)-B(1)	130.5(3)
C(11)-N(1)-B(1)	122.4(3)
C(11)-N(2)-C(11)#1	119.8(5)
N(2)-C(11)-N(1)	124.1(4)
N(2)-C(11)-C(10)	126.4(4)
N(1)-C(11)-C(10)	109.5(4)
N(1)-C(8)-C(9)	107.1(4)
N(1)-C(8)-C(5)	126.9(4)
C(9)-C(8)-C(5)	125.9(4)
C(4)-C(5)-C(6)	115.4(4)
C(4)-C(5)-C(8)	127.1(4)
C(6)-C(5)-C(8)	117.4(4)

C(9)-C(10)-C(11)	105.0(4)
C(9)-C(10)-C(12)	128.1(4)
C(11)-C(10)-C(12)	126.9(4)
C(2)-O(1)-C(1)	118.4(5)
C(10)-C(9)-C(8)	111.3(4)
C(17)-C(12)-C(13)	117.9(5)
C(17)-C(12)-C(10)	119.3(4)
C(13)-C(12)-C(10)	122.6(4)
O(1)-C(2)-C(3)	115.4(5)
O(1)-C(2)-C(7)	124.6(5)
C(3)-C(2)-C(7)	120.0(4)
C(2)-C(3)-C(4)	119.9(5)
C(3)-C(4)-C(5)	122.8(4)
C(7)-C(6)-C(5)	122.3(5)
F(2)-C(15)-C(16)	114.7(7)
F(2)-C(15)-C(14)	120.4(6)
C(16)-C(15)-C(14)	124.7(5)
C(14)-C(13)-C(12)	122.0(5)
C(12)-C(17)-C(16)	121.9(5)
C(6)-C(7)-C(2)	119.5(5)
F(1)-B(1)-F(1)#1	111.8(5)
F(1)-B(1)-N(1)	109.19(16)
F(1)#1-B(1)-N(1)	109.74(16)
F(1)-B(1)-N(1)#1	109.74(16)
F(1)#1-B(1)-N(1)#1	109.18(16)
N(1)-B(1)-N(1)#1	107.1(5)
C(15)-C(16)-C(17)	117.7(5)
C(13)-C(14)-C(15)	115.5(5)

Symmetry transformations used to generate equivalent atoms:

#1 -x+1,-y+1/2,z+0

Table S4. Anisotropic displacement parameters ($\text{\AA}^2 \times 10^3$). The anisotropic displacement factor exponent takes the form: $-2\pi^2 [h^2 a^{*2} U^{11} + \dots + 2 h k a^* b^* U^{12}]$

	U^{11}	U^{22}	U^{33}	U^{23}	U^{13}	U^{12}
F(1)	69(2)	65(2)	70(2)	-1(1)	-10(1)	2(1)
N(1)	65(2)	49(2)	53(2)	2(1)	1(2)	3(1)
N(2)	58(3)	48(2)	65(3)	0	0	0(2)
C(11)	64(2)	57(2)	48(2)	-1(2)	1(2)	3(2)
C(8)	61(2)	53(2)	61(2)	4(2)	4(2)	3(2)
C(5)	65(2)	54(2)	59(2)	5(2)	-6(2)	7(2)
C(10)	65(3)	49(2)	61(2)	4(2)	6(2)	1(2)
O(1)	103(3)	92(3)	72(2)	16(2)	-15(2)	6(2)
C(9)	77(3)	54(2)	63(2)	-2(2)	2(2)	7(2)
C(12)	66(3)	58(2)	56(2)	-1(2)	6(2)	7(2)
C(2)	72(3)	61(3)	73(3)	11(2)	-8(2)	-2(2)
C(3)	88(3)	66(3)	60(2)	3(2)	4(2)	4(2)
C(4)	86(3)	64(3)	62(3)	2(2)	1(2)	15(2)
C(6)	99(4)	71(3)	66(3)	1(2)	-1(3)	27(3)
C(15)	123(5)	88(4)	51(2)	-10(3)	9(3)	15(3)
C(13)	86(4)	76(3)	74(3)	5(2)	3(3)	4(3)
C(17)	115(4)	67(3)	71(3)	2(2)	-6(3)	7(3)
C(7)	94(4)	70(3)	78(3)	11(3)	-5(3)	29(3)
B(1)	53(3)	61(4)	52(3)	0	0	8(3)
C(16)	133(5)	69(3)	68(3)	-14(3)	-4(3)	21(3)
C(14)	111(4)	89(4)	62(3)	7(3)	10(3)	0(3)
F(2)	415(14)	190(7)	204(7)	-10(6)	73(7)	67(8)
C(1)	156(8)	139(6)	132(6)	54(5)	-13(5)	64(6)

Table S5. Hydrogen coordinates ($\text{\AA} \times 10^4$) and isotropic displacement parameters ($\text{\AA}^2 \times 10^3$).

	x	y	z	U(eq)
H(9)	4060	4889	1367	78
H(3)	4370	3990	-1513	86
H(4)	4752	3436	-524	85
H(6)	3565	5222	438	94
H(13)	4050	3027	2902	95
H(17)	4743	5259	2414	101
H(7)	3179	5770	-549	97
H(16)	4698	5646	3522	108
H(14)	3983	3401	4015	105
H(1A)	3426	6401	-1664	213
H(1B)	2622	5901	-1498	213
H(1C)	2909	6003	-2245	213

Table S6. Torsion angles [°].

C(11)#1-N(2)-C(11)-N(1)	0.8(3)
C(11)#1-N(2)-C(11)-C(10)	179.9(5)
C(8)-N(1)-C(11)-N(2)	177.2(4)
B(1)-N(1)-C(11)-N(2)	-1.5(6)
C(8)-N(1)-C(11)-C(10)	-2.0(5)
B(1)-N(1)-C(11)-C(10)	179.2(3)
C(11)-N(1)-C(8)-C(9)	2.1(5)
B(1)-N(1)-C(8)-C(9)	-179.3(3)
C(11)-N(1)-C(8)-C(5)	-175.4(4)
B(1)-N(1)-C(8)-C(5)	3.3(7)
N(1)-C(8)-C(5)-C(4)	-12.5(8)
C(9)-C(8)-C(5)-C(4)	170.5(5)
N(1)-C(8)-C(5)-C(6)	167.4(5)
C(9)-C(8)-C(5)-C(6)	-9.6(7)
N(2)-C(11)-C(10)-C(9)	-178.1(4)
N(1)-C(11)-C(10)-C(9)	1.1(5)
N(2)-C(11)-C(10)-C(12)	3.8(7)
N(1)-C(11)-C(10)-C(12)	-176.9(4)
C(11)-C(10)-C(9)-C(8)	0.2(6)
C(12)-C(10)-C(9)-C(8)	178.2(5)
N(1)-C(8)-C(9)-C(10)	-1.4(6)
C(5)-C(8)-C(9)-C(10)	176.0(4)
C(9)-C(10)-C(12)-C(17)	-34.6(8)
C(11)-C(10)-C(12)-C(17)	143.0(5)
C(9)-C(10)-C(12)-C(13)	140.8(5)
C(11)-C(10)-C(12)-C(13)	-41.5(7)
C(1)-O(1)-C(2)-C(3)	-179.4(6)
C(1)-O(1)-C(2)-C(7)	2.2(8)
O(1)-C(2)-C(3)-C(4)	-179.6(5)
C(7)-C(2)-C(3)-C(4)	-1.2(8)
C(2)-C(3)-C(4)-C(5)	0.7(8)
C(6)-C(5)-C(4)-C(3)	-0.4(8)
C(8)-C(5)-C(4)-C(3)	179.5(5)
C(4)-C(5)-C(6)-C(7)	0.5(8)
C(8)-C(5)-C(6)-C(7)	-179.4(5)
C(17)-C(12)-C(13)-C(14)	-3.2(8)
C(10)-C(12)-C(13)-C(14)	-178.6(5)
C(13)-C(12)-C(17)-C(16)	3.6(8)
C(10)-C(12)-C(17)-C(16)	179.2(5)
C(5)-C(6)-C(7)-C(2)	-1.0(9)
O(1)-C(2)-C(7)-C(6)	179.6(5)
C(3)-C(2)-C(7)-C(6)	1.3(8)

C(8)-N(1)-B(1)-F(1)	-59.0(5)
C(11)-N(1)-B(1)-F(1)	119.5(4)
C(8)-N(1)-B(1)-F(1)#1	63.8(5)
C(11)-N(1)-B(1)-F(1)#1	-117.7(4)
C(8)-N(1)-B(1)-N(1)#1	-177.8(4)
C(11)-N(1)-B(1)-N(1)#1	0.7(3)
F(2)-C(15)-C(16)-C(17)	172.1(7)
C(14)-C(15)-C(16)-C(17)	-3.1(10)
C(12)-C(17)-C(16)-C(15)	-0.6(9)
C(12)-C(13)-C(14)-C(15)	-0.2(8)
F(2)-C(15)-C(14)-C(13)	-171.5(8)
C(16)-C(15)-C(14)-C(13)	3.4(10)

Symmetry transformations used to generate equivalent atoms:

#1 -x+1,-y+1/2,z+0

4. Ground state geometries in gas phase (M062X/6-311+G(2d,p))

Compound 3 E= -1819.96788515 a.u.

5	-2.0188870	0.0138560	0.3350060
9	-3.0272740	0.0012940	-0.5961060
9	-2.5129360	0.0474810	1.6404940
7	-1.0960220	-1.2445260	0.1814100
6	0.2921760	-1.1519410	0.2082710
7	0.9587420	-0.0164310	0.1877800
6	0.3141300	1.1296400	0.1231910
7	-1.0732940	1.2479980	0.1240010
6	-1.4122420	-2.5482270	0.0326460
6	-0.2245080	-3.3207940	-0.0368340
6	0.8550800	-2.4661510	0.0929870
6	-1.3685150	2.5540520	-0.0447040
6	-0.1679310	3.3023500	-0.1490980
6	0.8988090	2.4297490	-0.0378740
6	2.2809550	-2.7960780	0.0727170
6	2.7155920	-3.9553720	-0.5873310
1	1.9928900	-4.5834250	-1.0990190
6	4.0601840	-4.2989880	-0.6222400
1	4.4100290	-5.1879280	-1.1347680
6	4.9720160	-3.4661390	0.0107130
6	4.5830720	-2.3120230	0.6749250
1	5.3303210	-1.6962980	1.1631050
6	3.2337860	-1.9825340	0.7069900
1	2.9134640	-1.0898400	1.2313400
6	2.3286280	2.7507620	-0.0571120
6	3.2955780	1.8170870	-0.4643660
1	2.9875520	0.8267220	-0.7775630
6	4.6442640	2.1506470	-0.4747810
1	5.4015070	1.4413710	-0.7902960
6	5.0203160	3.4260920	-0.0804410
6	4.0950040	4.3777020	0.3237250
1	4.4337530	5.3607660	0.6305750
6	2.7507290	4.0310130	0.3327110
1	2.0179190	4.7575270	0.6699120
6	-2.7698250	-3.0990940	-0.0335910
6	-2.9885080	-4.2258970	-0.8405540
6	-4.2485860	-4.8060200	-0.9151610
1	-4.4074570	-5.6692630	-1.5533020
6	-5.3039470	-4.2761640	-0.1747830
6	-5.0912000	-3.1640110	0.6377100

1	-5.9081720	-2.7550160	1.2234600
6	-3.8347450	-2.5740630	0.7112450
1	-3.6713140	-1.7222550	1.3599050
6	-2.7165050	3.1278230	-0.1253640
6	-2.9140810	4.2366160	-0.9625800
6	-4.1631740	4.8375490	-1.0546580
1	-4.3052180	5.6858690	-1.7163550
6	-5.2291240	4.3473350	-0.3022430
6	-5.0374720	3.2537180	0.5398660
1	-5.8623730	2.8751300	1.1349020
6	-3.7920930	2.6428740	0.6312440
1	-3.6460660	1.8058900	1.3025780
1	-2.0896060	4.6089000	-1.5630850
1	-2.1709050	-4.6289810	-1.4305910
1	-0.1258300	4.3736710	-0.2877690
1	-0.2011370	-4.3993820	-0.1079480
9	6.3217680	3.7514770	-0.0896120
1	-6.2884870	-4.7298990	-0.2294940
1	-6.2050900	4.8174110	-0.3708510
9	6.2734970	-3.7889890	-0.0207170

Compound 4 E= -2049.00336427 a.u.

5	-1.3053530	0.0084970	0.4414670
9	-2.3325010	-0.0000150	-0.4713860
9	-1.7812100	0.0373940	1.7554680
7	-0.3893860	-1.2500670	0.2669160
6	0.9990330	-1.1578480	0.2497870
7	1.6662870	-0.0229310	0.2131600
6	1.0201350	1.1236980	0.1749680
7	-0.3664650	1.2435030	0.2191270
6	-0.7132420	-2.5547170	0.1256580
6	0.4741620	-3.3258650	0.0169240
6	1.5566540	-2.4716160	0.1126310
6	-0.6678160	2.5522150	0.0674590
6	0.5326270	3.2983160	-0.0687700
6	1.6002680	2.4239010	0.0036240
6	2.9815560	-2.8020070	0.0459330
6	3.3952810	-3.9585820	-0.6318780
1	2.6566430	-4.5847320	-1.1228110
6	4.7382340	-4.3021580	-0.7106500
1	5.0715080	-5.1892070	-1.2373730
6	5.6698110	-3.4717240	-0.1040610
6	5.3020510	-2.3201960	0.5762910

1	6.0644150	-1.7058620	1.0424310
6	3.9543090	-1.9910290	0.6524690
1	3.6505290	-1.0997820	1.1889790
6	3.0298350	2.7436710	-0.0558640
6	3.9824480	1.8126350	-0.5008980
1	3.6630810	0.8251800	-0.8119710
6	5.3308660	2.1444620	-0.5500600
1	6.0768490	1.4367620	-0.8946780
6	5.7211000	3.4157980	-0.1565660
6	4.8102310	4.3647280	0.2845130
1	5.1597840	5.3445850	0.5896100
6	3.4661770	4.0195240	0.3322410
1	2.7453460	4.7438650	0.6987430
6	-2.0668750	-3.1052050	0.0991910
6	-2.3045050	-4.2559290	-0.6626050
6	-3.5601100	-4.8493440	-0.7143680
1	-3.7082890	-5.7293400	-1.3281250
6	-4.6100100	-4.2935090	0.0228290
6	-4.3832220	-3.1491560	0.8005810
1	-5.2096250	-2.7442820	1.3749400
6	-3.1337550	-2.5618650	0.8378810
1	-2.9716460	-1.6919520	1.4619440
8	-5.8678130	-4.7872380	0.0518850
6	-6.1393700	-5.9469900	-0.7154620
1	-5.9643120	-5.7641040	-1.7811800
1	-5.5254480	-6.7913850	-0.3836780
1	-7.1917690	-6.1752600	-0.5529840
6	-2.0114450	3.1278000	0.0296300
6	-2.2259380	4.2682360	-0.7545960
6	-3.4698820	4.8842870	-0.8209850
1	-3.5996860	5.7548220	-1.4521720
6	-4.5320500	4.3627580	-0.0764090
6	-4.3285850	3.2295290	0.7234750
1	-5.1638300	2.8510090	1.3030330
6	-3.0905360	2.6193490	0.7753940
1	-2.9475130	1.7585960	1.4162160
8	-5.7804540	4.8804440	-0.0606460
6	-6.0284970	6.0290820	-0.8522860
1	-5.8533440	5.8214760	-1.9134500
1	-5.4004630	6.8688710	-0.5355670
1	-7.0770960	6.2795890	-0.6982200
1	-1.4123440	4.6691380	-1.3514590
1	-1.4991350	-4.6835160	-1.2521910
1	0.5744830	4.3707730	-0.1982660

1	0.4974810	-4.4044660	-0.0533070
9	7.0226010	3.7395990	-0.2030500
9	6.9700640	-3.7943040	-0.1783930

Compound 5 E= -1720.72334179 a.u.

5	-1.7906740	0.3791090	0.3377880
9	-2.7614470	0.6732920	-0.5869760
9	-2.2695680	0.4758930	1.6459610
7	-1.2218180	-1.0660150	0.1174950
6	0.1447270	-1.3346570	0.1180150
7	1.0795610	-0.4102030	0.1838640
6	0.7511150	0.8650830	0.2054450
7	-0.5583690	1.3357570	0.1828330
6	-1.8653500	-2.2394210	-0.0550650
6	-0.9171720	-3.2897250	-0.1592480
6	0.3483750	-2.7456120	-0.0439660
6	-0.5048250	2.6764920	0.0370110
6	0.8488920	3.0930430	-0.0352470
6	1.6529910	1.9743660	0.0895370
6	1.6341800	-3.4508080	-0.0592210
6	1.6859120	-4.7902490	0.3543350
1	0.7804270	-5.2756660	0.7066290
6	2.8868110	-5.4893040	0.3468610
1	2.9100180	-6.5231650	0.6765460
6	4.0581700	-4.8612980	-0.0737660
6	4.0178760	-3.5314720	-0.4883610
1	4.9251250	-3.0377400	-0.8229560
6	2.8178560	-2.8283890	-0.4852290
1	2.7912220	-1.7972320	-0.8175590
6	3.1148210	1.9021420	0.0635860
6	3.8115110	0.8493240	0.6786570
1	3.2614940	0.0687790	1.1909240
6	5.1993440	0.7982850	0.6407710
1	5.7514050	-0.0063330	1.1136390
6	5.8865280	1.8125610	-0.0096880
6	5.2347730	2.8728060	-0.6234690
1	5.8123520	3.6404250	-1.1259260
6	3.8475960	2.9089920	-0.5830390
1	3.3227700	3.7190440	-1.0799930
6	-3.3191810	-2.4184810	-0.1389590
6	-3.8136130	-3.4264920	-0.9807900
6	-5.1799420	-3.6589920	-1.0760200
1	-5.5494230	-4.4327490	-1.7411870

6	-6.0704500	-2.8968420	-0.3219010
6	-5.5857820	-1.9022450	0.5253090
1	-6.2750310	-1.3134750	1.1220700
6	-4.2203540	-1.6590670	0.6196330
1	-3.8501330	-0.8985230	1.2957020
6	-1.6594090	3.5790270	-0.0227300
6	-1.5637640	4.7255060	-0.8258720
6	-2.6162380	5.6299450	-0.8919450
1	-2.5346060	6.5061180	-1.5270150
6	-3.7730740	5.4077910	-0.1469450
6	-3.8704140	4.2768450	0.6615960
1	-4.7655180	4.1057890	1.2509610
6	-2.8241070	3.3640960	0.7267130
1	-2.8982050	2.4977190	1.3724470
1	-0.6695720	4.8907250	-1.4194080
1	-3.1227500	-4.0097440	-1.5819810
1	1.1670900	4.1238980	-0.1052250
1	-1.1707670	-4.3310890	-0.3005900
9	7.2267230	1.7676650	-0.0473140
1	4.9969850	-5.4061470	-0.0783910
1	-7.1381420	-3.0790170	-0.3929190
1	-4.5951740	6.1149240	-0.1951140

Compound 6 E= -1949.75869866 a.u.

5	-1.1062760	0.1520520	0.4435550
9	-2.1100610	0.4046320	-0.4604470
9	-1.5667080	0.2295880	1.7613570
7	-0.4839410	-1.2669410	0.2112740
6	0.8923930	-1.4759130	0.1660760
7	1.7886700	-0.5122390	0.2028400
6	1.4053200	0.7476600	0.2405760
7	0.0769190	1.1614490	0.2642420
6	-1.0843960	-2.4675360	0.0571510
6	-0.0926320	-3.4748260	-0.0800820
6	1.1505250	-2.8764080	-0.0059380
6	0.0653160	2.5058910	0.1258360
6	1.3988360	2.9789550	0.0118530
6	2.2531120	1.8953630	0.1010280
6	2.4645380	-3.5264100	-0.0636940
6	2.5861030	-4.8639480	0.3411950
1	1.7135170	-5.3884390	0.7194730
6	3.8142880	-5.5122570	0.2919540
1	3.8909000	-6.5455930	0.6153460

6	4.9445190	-4.8343320	-0.1621090
6	4.8351680	-3.5056370	-0.5676880
1	5.7099040	-2.9726410	-0.9277710
6	3.6073570	-2.8534570	-0.5229790
1	3.5270960	-1.8226180	-0.8476250
6	3.7155350	1.8874450	0.0263510
6	4.4785250	0.8653430	0.6137200
1	3.9806660	0.0592130	1.1396060
6	5.8654470	0.8766620	0.5302940
1	6.4680300	0.0960590	0.9813250
6	6.4851050	1.9224320	-0.1377440
6	5.7667880	2.9536320	-0.7257970
1	6.2926860	3.7476540	-1.2437910
6	4.3813200	2.9271320	-0.6401700
1	3.8045220	3.7136420	-1.1169280
6	-2.5258570	-2.7097760	0.0181850
6	-3.0034420	-3.7646880	-0.7696280
6	-4.3578740	-4.0685590	-0.8379560
1	-4.6895350	-4.8817200	-1.4718630
6	-5.2668930	-3.3128100	-0.0916490
6	-4.8015190	-2.2630710	0.7127350
1	-5.5239450	-1.6997470	1.2937080
6	-3.4542910	-1.9634240	0.7664710
1	-3.1116400	-1.1636450	1.4106550
8	-6.6025290	-3.5198850	-0.0779590
6	-7.1153230	-4.5715770	-0.8769880
1	-6.8940700	-4.4048350	-1.9367270
1	-6.7055110	-5.5386460	-0.5658660
1	-8.1936680	-4.5666010	-0.7246000
6	-1.1234900	3.3562280	0.1071450
6	-1.0920770	4.5311350	-0.6539180
6	-2.1750480	5.4012340	-0.6963880
1	-2.1182890	6.2921530	-1.3096690
6	-3.3198170	5.1050070	0.0494530
6	-3.3603040	3.9384390	0.8261360
1	-4.2538820	3.7368750	1.4071730
6	-2.2819750	3.0758910	0.8542680
1	-2.3222880	2.1913790	1.4775890
8	-4.4275710	5.8786390	0.0883350
6	-4.4261420	7.0719140	-0.6755470
1	-4.3069780	6.8563910	-1.7428380
1	-3.6292040	7.7483000	-0.3479010
1	-5.3946380	7.5396080	-0.5042030
1	-0.2137760	4.7596020	-1.2500760

1	-2.3073300	-4.3449030	-1.3676060
1	1.6725690	4.0226140	-0.0573450
1	-0.3030730	-4.5270750	-0.2113840
9	7.8244550	1.9381850	-0.2188570
1	5.9047190	-5.3392380	-0.1993650

Compound 8 E= -1850.51390976 a.u.

5	-0.9147740	0.0099800	0.4371810
9	-1.9478460	0.0046140	-0.4693430
9	-1.3832000	0.0405250	1.7540870
7	-0.0040940	-1.2512460	0.2572740
6	1.3848300	-1.1637900	0.2312380
7	2.0549010	-0.0306410	0.1917220
6	1.4125390	1.1181170	0.1577440
7	0.0262220	1.2419120	0.2092020
6	-0.3333710	-2.5546730	0.1183510
6	0.8506060	-3.3296070	0.0013730
6	1.9370890	-2.4794690	0.0893910
6	-0.2721840	2.5513040	0.0587780
6	0.9296930	3.2939000	-0.0831980
6	1.9955670	2.4168930	-0.0158560
6	3.3607050	-2.8185460	0.0127550
6	3.7604980	-3.9789610	-0.6661280
1	3.0114890	-4.5987380	-1.1504680
6	5.1037850	-4.3251620	-0.7486940
1	5.3965890	-5.2238420	-1.2824430
6	6.0718720	-3.5153750	-0.1563730
6	5.6859550	-2.3602870	0.5212120
1	6.4339820	-1.7281980	0.9898440
6	4.3420860	-2.0130740	0.6109350
1	4.0460350	-1.1186390	1.1470030
6	3.4261450	2.7357700	-0.0833530
6	4.3744810	1.8057340	-0.5371680
1	4.0511250	0.8192550	-0.8481900
6	5.7217670	2.1472930	-0.5917110
1	6.4430910	1.4174200	-0.9465390
6	6.1454450	3.4163510	-0.2021970
6	5.2105930	4.3481500	0.2462790
1	5.5318330	5.3372170	0.5574400
6	3.8638900	4.0104810	0.3057240
1	3.1428130	4.7315670	0.6795310
6	-1.6893150	-3.1004320	0.1001900
6	-1.9359200	-4.2494130	-0.6612580
6	-3.1939400	-4.8385340	-0.7054530

1	-3.3490410	-5.7173970	-1.3191450
6	-4.2370710	-4.2799460	0.0390850
6	-4.0012840	-3.1372720	0.8165020
1	-4.8225560	-2.7300580	1.3965610
6	-2.7495010	-2.5543100	0.8463100
1	-2.5802980	-1.6854150	1.4699290
8	-5.4966050	-4.7694730	0.0757600
6	-5.7769740	-5.9269120	-0.6917040
1	-5.6082290	-5.7428730	-1.7582680
1	-5.1637580	-6.7739890	-0.3654400
1	-6.8290660	-6.1520330	-0.5228230
6	-1.6145510	3.1306930	0.0279670
6	-1.8300960	4.2711960	-0.7557510
6	-3.0726670	4.8908510	-0.8157040
1	-3.2033290	5.7614900	-1.4465840
6	-4.1322300	4.3727820	-0.0651470
6	-3.9276220	3.2395700	0.7343640
1	-4.7607900	2.8637240	1.3186580
6	-2.6909700	2.6259160	0.7799860
1	-2.5468250	1.7651240	1.4205270
8	-5.3793070	4.8940410	-0.0430490
6	-5.6282660	6.0423770	-0.8346530
1	-5.4592650	5.8331430	-1.8965190
1	-4.9962500	6.8808340	-0.5223020
1	-6.6753450	6.2960690	-0.6754170
1	-1.0184760	4.6694160	-1.3570830
1	-1.1357890	-4.6790990	-1.2564290
1	0.9744190	4.3662190	-0.2127350
1	0.8700870	-4.4081830	-0.0700000
1	7.1978960	3.6784880	-0.2471800
1	7.1215240	-3.7840830	-0.2226090

Compound 9 E= -2177.67866629 a.u.

5	-1.8822680	0.0262760	0.4515520
9	-2.9023820	0.0057230	-0.4690720
9	-2.3691160	0.0850420	1.7607230
7	-0.9738150	-1.2414000	0.3101760
6	0.4150280	-1.1594760	0.3076780
7	1.0907110	-0.0300580	0.2550570
6	0.4529420	1.1199680	0.1854110
7	-0.9329920	1.2497030	0.2120730
6	-1.3060250	-2.5464590	0.1896960
6	-0.1241720	-3.3280760	0.1096470

6	0.9645660	-2.4801190	0.2025840
6	-1.2241330	2.5573950	0.0301860
6	-0.0183020	3.2926440	-0.1079120
6	1.0437850	2.4130370	-0.0060030
6	2.3856250	-2.8230130	0.1614040
6	2.7997190	-4.0043510	-0.4706160
1	2.0628280	-4.6437770	-0.9472150
6	4.1387400	-4.3608590	-0.5257180
1	4.4339730	-5.2780210	-1.0273110
6	5.0983030	-3.5278140	0.0574070
6	4.7049070	-2.3481380	0.6934860
1	5.4625010	-1.7185210	1.1479820
6	3.3639710	-2.0026380	0.7479260
1	3.0665440	-1.0907430	1.2524410
6	2.4739370	2.7235720	-0.0566800
6	3.4343280	1.7752690	-0.4479660
1	3.1198520	0.7763910	-0.7257900
6	4.7801370	2.1039430	-0.4873030
1	5.5228640	1.3741380	-0.7921460
6	5.1983030	3.3905920	-0.1411700
6	4.2575250	4.3492770	0.2459980
1	4.5709680	5.3515390	0.5238560
6	2.9132410	4.0109010	0.2852910
1	2.1922390	4.7545300	0.6109550
6	-2.6638250	-3.0865180	0.1577090
6	-2.9027000	-4.2486540	-0.5860130
6	-4.1625610	-4.8328040	-0.6407680
1	-4.3114850	-5.7222350	-1.2405880
6	-5.2155810	-4.2555720	0.0751400
6	-4.9877890	-3.0992060	0.8346130
1	-5.8169520	-2.6774120	1.3925680
6	-3.7339960	-2.5213650	0.8748660
1	-3.5712280	-1.6414670	1.4846490
8	-6.4776170	-4.7388850	0.0996480
6	-6.7499290	-5.9107370	-0.6487600
1	-6.5624650	-5.7491460	-1.7158000
1	-6.1458140	-6.7532590	-0.2948780
1	-7.8056480	-6.1282390	-0.4931570
6	-2.5637530	3.1403000	-0.0337170
6	-2.7633670	4.2655910	-0.8432690
6	-4.0025910	4.8880930	-0.9341960
1	-4.1207080	5.7462500	-1.5843700
6	-5.0751710	4.3887150	-0.1893790
6	-4.8867520	3.2708130	0.6353360

1	-5.7299630	2.9095540	1.2143750
6	-3.6532040	2.6541010	0.7117790
1	-3.5219410	1.8058440	1.3715030
8	-6.3202920	4.9146260	-0.1960540
6	-6.5529220	6.0492910	-1.0120750
1	-6.3691260	5.8200590	-2.0673010
1	-5.9221970	6.8908800	-0.7056140
1	-7.6012030	6.3097440	-0.8730090
1	-1.9414840	4.6490030	-1.4402800
1	-2.0945790	-4.6930660	-1.1591610
1	0.0313260	4.3619040	-0.2589490
1	-0.1087220	-4.4079460	0.0606120
7	6.5935850	3.6323920	-0.2118740
7	6.4898880	-3.7977810	0.0533500
7	6.8433560	-4.8430990	-0.5062600
7	6.9676950	4.7722210	0.0903550
7	7.2731860	-5.7673270	-0.9890610
7	7.4153580	5.7758610	0.3444120

Compound 10 E= -2005.11678847 a.u.

5	-1.3069270	0.0239550	0.4386450
9	-2.3368660	0.0427130	-0.4725540
9	-1.7841970	0.0596430	1.7536190
7	-0.4192180	-1.2501210	0.2594150
6	0.9684350	-1.1885520	0.2454300
7	1.6628550	-0.0616120	0.2115640
6	1.0439340	1.0930360	0.1767420
7	-0.3432010	1.2399330	0.2217540
6	-0.7725080	-2.5501500	0.1177850
6	0.3948660	-3.3429610	0.0117670
6	1.4974020	-2.5077260	0.1087080
6	-0.6160250	2.5549600	0.0774330
6	0.5960890	3.2806560	-0.0540940
6	1.6524970	2.3904140	0.0126590
6	2.9154530	-2.8694820	0.0455720
6	3.3086570	-4.0272790	-0.6419020
1	2.5591740	-4.6321040	-1.1431610
6	4.6446390	-4.3994580	-0.7165660
1	4.9614020	-5.2883760	-1.2504220
6	5.5905300	-3.5961490	-0.0964270
6	5.2435330	-2.4445760	0.5944250
1	6.0157900	-1.8520600	1.0725330
6	3.9024670	-2.0876390	0.6669970

1	3.6145140	-1.1967540	1.2129850
6	3.0790360	2.6888110	-0.0470350
6	4.0420150	1.7283940	-0.4053240
1	3.7261360	0.7213360	-0.6514400
6	5.3872840	2.0503070	-0.4622080
1	6.1122560	1.2923780	-0.7476540
6	5.8306050	3.3495060	-0.1630030
6	4.8738180	4.3151580	0.1948540
1	5.1984750	5.3222890	0.4410640
6	3.5315130	3.9864150	0.2471830
1	2.8161620	4.7471190	0.5464150
6	-2.1409570	-3.0685450	0.0893980
6	-2.4100500	-4.2026830	-0.6855170
6	-3.6791530	-4.7687040	-0.7349050
1	-3.8505220	-5.6377230	-1.3583570
6	-4.7114170	-4.1995830	0.0158930
6	-4.4540670	-3.0700770	0.8047700
1	-5.2675610	-2.6530310	1.3889240
6	-3.1906780	-2.5117070	0.8414260
1	-3.0052080	-1.6521770	1.4734850
8	-5.9811500	-4.6665750	0.0479160
6	-6.2812080	-5.8103210	-0.7314680
1	-6.1095910	-5.6183540	-1.7962980
1	-5.6825150	-6.6715450	-0.4149750
1	-7.3370080	-6.0197180	-0.5648980
6	-1.9482110	3.1568410	0.0393460
6	-2.1380420	4.3051590	-0.7397520
6	-3.3697290	4.9451420	-0.8073930
1	-3.4808570	5.8208800	-1.4349380
6	-4.4439100	4.4406320	-0.0683080
6	-4.2648020	3.3001180	0.7270490
1	-5.1090170	2.9351400	1.3022620
6	-3.0391750	2.6654580	0.7792670
1	-2.9146880	1.7987090	1.4157710
8	-5.6821980	4.9820780	-0.0540960
6	-5.9061710	6.1377540	-0.8425910
1	-5.7314970	5.9304280	-1.9038920
1	-5.2636150	6.9646320	-0.5210750
1	-6.9504050	6.4073120	-0.6910050
1	-1.3149020	4.6929650	-1.3320530
1	-1.6178200	-4.6399240	-1.2859160
1	0.6524410	4.3530230	-0.1762190
1	0.3979290	-4.4220130	-0.0577070
9	6.8848790	-3.9462800	-0.1669760

7	7.1785570	3.6573280	-0.1630720
1	7.4072440	4.6361280	-0.2795330
1	7.7677710	3.0552490	-0.7233960

ACCEPTED MANUSCRIPT

- Aymmetric/asymmetric fluorine-containing aza-BODIPYs at 1/7-positions were prepared.
- The effect of the withdrawing -F group at 1/7-position on the photophysical properties is found to be almost negligible.
- The replacement of -F atoms with -N₃/-NH₂ groups resulted in a remarkable bathochromic shift.
- Aza-BODIPY with a -NH₂ group could be used as a turn-on fluorescent probe for pH value.