

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23

Palladium-Catalyzed Cascade sp^2 C-H Bond Functionalizations Allowing the One Pot Access to 4-Aryl-1,2,3,4-tetrahydroquinolines from *N*-Allyl-*N*- arylsulfonamides

24 *Kedong Yuan,^[a,b] Jean-François Soulé,*^[b] Vincent Dorcet,^[b] Henri Doucet*^[b]*

25
26
27
28 [a] Tianjin Key Laboratory of Advanced Functional Porous Material, Institute for New Energy
29 Materials & Low-Carbon Technologies, School of Materials Science and Engineering, Tianjin
30 University of Technology, Tianjin 300384, China.
31
32

33
34
35
36 [b] UMR 6226 CNRS-Université de Rennes 1, 35042 Rennes, France.
37
38
39
40

41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

KEYWORDS Palladium • C-H activation • catalysis • desulfitative coupling • cascade


ABSTRACT We have developed a palladium-catalyzed cascade reaction allowing an efficient synthesis of 4-aryl-1,2,3,4-tetrahydroquinolines from *N*-allyl-*N*-arylsulfonamides and benzenesulfonyl chlorides. In this transformation, two C(sp^2)-C(sp^3) bonds were formed *via* activation of C(sp^2)-H bonds. The reaction proceeds using easily accessible PdCl₂ catalyst, with Li₂CO₃ as inexpensive base and CuBr as additive and tolerates a wide variety of substituents on both reaction partners.

Introduction

One current important area of modern synthetic chemistry is the development of methods minimizing both the requisite number of steps and formation of wastes. Among these methods, the metal-catalyzed functionalization of C-H bonds has emerged as a simpler and “greener” way for the access to useful molecules for biological or material applications, as such processes are capable of forming similar products while avoiding the use of stoichiometric organometallic reagents. Most of the examples of such C-H bond functionalizations concern the formation of C-C bonds *via* arylations, alkylations or alkenylations;¹ whereas, the formation of two C-C bonds *via* consecutive Heck type reaction followed by an sp² C-H bond activation has attracted less attention.^{2,3} Among rare examples, Fagnou et al. described the formation of indolines from 2-bromoaniline derivatives and heteroarenes *via* a domino palladium-catalyzed Heck-intermolecular direct arylation.^{3d} In 2012, Wu et al. described the synthesis of 4-polyfluoroaryl pyrrolo[1,2-a]quinolines *via* intermolecular followed by intramolecular palladium-catalyzed sp² C-H bond activations.^{3e} Recently, Zhu and co-workers reported the synthesis of [3,4]-fused oxindoles *via* a double palladium-catalyzed sp² C-H bond activation.^{3g} On the other hand, the Heck reaction^{4,5} using *N*-allylbenzenesulfonamides and aryl halides as coupling partners has been described by Y. Dong (Scheme 1 top).⁶

In this article, we report on the unexpected one pot synthesis of 4-aryl-1,2,3,4-tetrahydroquinolines, from *N*-allyl-*N*-benzenesulfonamides and ArSO₂Cl through a palladium-catalyzed cascade desulfitative addition–cyclization reaction (Scheme 1, bottom).⁷⁻⁹ It should be mentioned that 1,2,3,4-tetrahydroquinoline is an important motif found in numerous compounds with important antitumoral, antibacterial or antioxidant activities.¹⁰ Several methods are leading to the synthesis of 1,2,3,4-tetrahydroquinolines, including partial reduction of quinolines,


condensation of anilines with two molecules of an aldehyde, or from a Schiff base and an alkene.^{10,11} However, most of these methods involve a multi-steps synthesis and in several cases does not allow the introduction of specific functional groups at the desired positions.



Scheme 1.

Results and discussion

The *N*-allyl-*N*-arylsulfonamides **1-22** were first prepared in high yields from anilines, methyl- or benzene-sulfonyl chlorides and allyl bromides (Scheme 2).


Scheme 2. Structures of **1-22**

We then examined the influence of the conditions on the products formation using *N*-allyl-*N*-*p*-tolylbenzenesulfonamide **1** and 4-methylbenzenesulfonyl chloride as the coupling partners (Table 1). Using PdCl₂ catalyst and Li₂CO₃ as base in dioxane as reaction conditions, which had been previously found operative for direct arylations of heterocycles,¹² no reaction occurred. On

1
2
3 the other hand, under the same conditions, but in the presence of 1.1 equiv. of CuBr as additive,
4 the unexpected 4-tolyl-1,2,3,4-tetrahydroquinoline **24** –resulting from a formal 6-endo-trig
5 cyclization after desulfitative addition– was obtained in 75% yield with 89% conversions of **1**
6
7
8 (Table 1, entry 2). It should be noted that no formation of the expected Heck type product **23**
9
10 was observed under these conditions. The use of Pd(OAc)₂, PdCl₂(MeCN)₂, PdCl(C₃H₅)(dppb),
11
12 Pd(TFA)₂ or Pd₂(dba)₃ catalysts did not increase the yield in the desired product **24** (Table 1,
13
14 entries 3-7). With Pd₂(dba)₃ catalyst in the absence of CuBr, deallylated **1** was the major
15
16 product. With PdCl₂ catalyst, a decrease of the CuBr loading to 0.1 equiv. had almost no
17
18 influence on the yield in **24**; whereas using 1 mol% CuBr gave **24** in low yield (Table 1, entries
19
20 9-11). Then, the influence of several bases was examined using 5 mol% of PdCl₂ associated to
21
22 0.5 equiv. of CuBr. Lower yields in **24** were obtained with K₂CO₃ and Na₂CO₃; whereas,
23
24 Cs₂CO₃ or a reaction without base were completely ineffective (Table 1, entries 12-15). The
25
26 decrease of the reaction temperature to 100 °C also affords **24** in good yield (Table 1, entry 16).
27
28 The influence of a few solvents was also investigated. No reaction occurred in xylene or DMF;
29
30 whereas **1** was obtained in 65% yield in diethyl carbonate (Table 1, entries 17-19). The addition
31
32 of 1.2 equiv. of TEMPO was found to quench almost completely the reaction without its
33
34 incorporation in **1** or **2**. This result suggests that no radical species is involved in this 6-endo-trig
35
36 cyclization (Table 1, entry 20). From **1** using 5 mol% PdCl₂ catalyst in the presence of CuBr and
37
38 Li₂CO₃ at 140 °C, but without benzenesulfonyl chloride, no cyclization to afford a 1,2,3,4-
39
40 tetrahydroquinoline occurred and **1** was recovered (Table 1, entry 21). When CuBr₂ was used as
41
42 additive instead of CuBr, a low conversion of **1** was observed, and **24** was formed in very low
43
44 yield together with unidentified side-products (Table 1, entry 22). Finally, the replacement of **1**
45
46 by *tert*-butyl allyl(phenyl)carbamate led to several unidentified side-products; whereas, the
47
48
49
50
51
52
53
54
55
56
57
58
59
60

desired cyclized product was not detected by GC/MS analysis of the crude mixture (Table 1, entry 23).


Table 1. Influence of the Reaction Conditions for the Po-Catalyzed Functionalization of **1**.



Entry	Catalyst (mol%)	Base	CuBr (equiv.)	Temp (°C)	Conv. of 1 (%)	Yield in 24 (%)
1	PdCl ₂	Li ₂ CO ₃	-	140	<10	0
2	PdCl ₂	Li ₂ CO ₃	1.1	140	89	75
3	Pd(OAc) ₂	Li ₂ CO ₃	1.1	140	81	75
4	PdCl ₂ (CH ₃ CN) ₂	Li ₂ CO ₃	1.1	140	83	74
5	PdCl(C ₃ H ₅)(dppb)	Li ₂ CO ₃	1.1	140	76	68
6	Pd(TFA) ₂	Li ₂ CO ₃	1.1	140	70	67
7	Pd ₂ (dba) ₃	Li ₂ CO ₃	1.1	140	74	53
8	Pd ₂ (dba) ₃	Li ₂ CO ₃	-	140	100	trace
9	PdCl ₂	Li ₂ CO ₃	0.5	140	88	83
10	PdCl ₂	Li ₂ CO ₃	0.1	140	80	73
11	PdCl ₂	Li ₂ CO ₃	0.01	140	23	20
12	PdCl ₂	K ₂ CO ₃	0.5	140	73	31
13	PdCl ₂	Na ₂ CO ₃	0.5	140	85	42
14	PdCl ₂	Cs ₂ CO ₃	0.5	140	0	0
15	PdCl ₂	-	0.5	140	40	0
16	PdCl ₂	Li ₂ CO ₃	0.5	100	83	74
17	PdCl ₂	Li ₂ CO ₃	0.5	120	0	0 ^a
18	PdCl ₂	Li ₂ CO ₃	0.5	120	0	0 ^b
19	PdCl ₂	Li ₂ CO ₃	0.5	120	72	65 ^c
20	PdCl ₂	Li ₂ CO ₃	0.5	140	<10	<5 ^d
21	PdCl ₂	Li ₂ CO ₃	0.5	140	0	0 ^e
22	PdCl ₂	Li ₂ CO ₃	0.5 ^f	140	30	<10
23	PdCl ₂	Li ₂ CO ₃	0.5	140	<20 ^g	-

Conditions: **1** (1 equiv.), *p*-TolSO₂Cl (2 equiv.), base (3 equiv), 1,4-dioxane. ^a in xylene, ^b in DMF, ^c in diethyl carbonate, ^d *p*-TolSO₂Cl (1 equiv.) and TEMPO as additive (1.2 equiv.), ^e without *p*-TolSO₂Cl. ^f CuBr₂ instead of CuBr, ^g Using *tert*-butyl allyl(phenyl)carbamate instead of **1**.

The use of benzenesulfonyl chlorides as coupling partner is crucial for this reaction, as 4-bromotoluene under conditions *i* led to starting material **1**; whereas the use of PdCl(C₃H₅)(dppb) catalyst and K₂CO₃ as base in DMF (conditions *ii*) selectively afforded the Heck type product **23** in 77% yield (Scheme 3).




i) PdCl₂ 5 mol%, CuBr (0.5 equiv.), Li₂CO₃ (3 equiv.), 1,4-dioxane, 140 °C, 20h: **1** recovered

ii) PdCl(C₃H₅)(dppb) 2.5 mol%, K₂CO₃ (3 equiv.), DMF, 140 °C, 20h: **23** 77%

Scheme 3. Control Reaction with an Aryl Bromide

Then, the scope of the ArSO₂Cl substituent for the synthesis of 4-aryl-1,2,3,4-tetrahydroquinoline derivatives from **1** was examined using 5 mol% PdCl₂ catalyst in the presence of 0.5 equiv. of CuBr and Li₂CO₃ at 140 °C as reaction conditions (Scheme 4). Both 4-cyano- and 4-methoxybenzenesulfonyl chlorides afforded the target products **25** and **26** in 72% and 95% yields, respectively. The reaction also proceeded nicely in the presence of PhSO₂Cl or 1-naphthyl-SO₂Cl to give **27** and **28** in 75% and 71% yields, respectively. A thiophene-2-carboxylate bearing a SO₂Cl substituent at C3 was also employed. Again the expected product **29** was obtained in good yield. The influence of the SO₂R moiety on the *N*-allyl-*N*-arylsulfonamide was also investigated. A 4-bromobenzenesulfonamide was tolerated to afford **30** in 67% yield, without cleavage of the C-Br bond. The reaction also proceeds nicely with *N*-allyl-*N*-phenylmethanesulfonamide **3**, as the coupling with 4-methyl-, 4-chloro-, and 4-bromobenzenesulfonyl chlorides affords the target products **31-33** in 74-75% yields. A lower yield in **34** was obtained from 2,3,4-trifluorobenzenesulfonyl chloride and **3**. It is worth mentioning that

even 5-bromothiophene-2-sulfonyl chloride afforded the desired product **35** in 78% yield, without cleavage of the thienyl C-Br bond.




Scheme 4. Scope of the ArSO₂Cl and SO₂R¹ Substituents.

The influence of the substituents on the aniline moiety was also examined (Scheme 5). From *N*-allyl-*N*-(4-bromophenyl)methanesulfonamide **4** and *p*-TolSO₂Cl as reaction partner, **36** was obtained in 42% yield. A 4-iodo substituent on the aniline was also tolerated to afford **37** and **40** in 68% and 67% yields. In both cases, the reaction was highly chemoselective, as the C-X bonds were not involved in the cascade process. Electron-donating substituents on the aniline part were tolerated as both *N*-allyl-*N*-(4-methylphenyl)methanesulfonamide **6** and *N*-allyl-*N*-(4-methoxyphenyl)methanesulfonamide **7** led to **38** and **39** in good yields.


1
2
3 The influence of substituent on the *N*-allyl moiety was also investigated. From (*E*)-*N*-(but-2-
4 enyl)-*N*-phenylmethanesulfonamide **8** and *p*-TolSO₂Cl, **41** was obtained in 81% as a single
5 diastereomer. Similar results were obtained for the coupling of **8** and **9** with 4-fluoro- and 4-
6 methoxybenzenesulfonyl chlorides affording **42** and **43** in 78% and 88% yields, respectively.
7
8 The *trans*-stereochemistry was unambiguously assigned by X-ray analysis of **43** (see SI). The
9 regioselectivity of the reaction with aniline derivatives bearing *meta*-bromo or *meta*-methoxy
10 substituents was then studied. In both cases, the formation of mixture of 1,2,3,4-
11 tetrahydroquinolines was obtained. The major products **44a** and **45a** arises from coupling at less
12 hindered position. Similar regioselectivity was observed using 3,4-dimethylaniline, as **46a** and
13 **46b** were obtained in 12:5 ratio. The reactivity of several fluoro-containing *N*-allyl-*N*-
14 (aryl)methanesulfonamides was also examined. 4-Fluoroaniline and 3,5-difluoroaniline
15 derivatives **13** and **14** gave the 1,2,3,4-tetrahydroquinolines **47** and **48** in 72% and 77% yields,
16 respectively with formation of only trace amount of Heck type products. On the other hand, a
17 very significant effect of aniline *ortho*-fluoro or methyl substituents was observed. From both *N*-
18 (2,4-difluorophenyl)methanesulfonamide **15** and *N*-(2-methylphenyl)methanesulfonamide **16**
19 using *p*-TolSO₂Cl as the reaction partner, only the Heck type products **49** and **50** were obtained.
20 This result might be explained by lower degree of freedom of C–N bond, which prevents the
21 suitable conformation for the formal 6-endo-trig cyclization. An aniline substituted at C4 by an
22 electron-withdrawing group such as ethyl ester led to a mixture of Heck type product **51a** and
23 tetrahydroquinoline **51b** in 5:1 ratio; whereas, an aniline bearing a para-thienyl at C4 only gave
24 tetrahydroquinoline **52** in 83% yield. Finally the reactivity of *N*-allyl-*N*-
25 cyclohexylmethanesulfonamide **19** was examined. No sp³ C–H bond functionalization of the
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

cyclohexyl moiety was observed and the Heck type product **53** was selectively obtained in 78% yield.




Scheme 5. Scope of the Substituents on the Aniline and Allyl moieties.

Interestingly, the reaction is not limited to the use of aniline derivatives. From 5-aminopyrazole derivative **54** and 4-methyl- or 4-chloro-benzenesulfonyl chlorides, the expected products **55** and **56** were obtained in 83% and 81% yields, respectively (Scheme 6). Even 5-bromothiophene-2-sulfonyl chloride reacts with **54** to afford **57** in 85% yield, without cleavage of the thienyl C-Br bond.



Scheme 6. Reactivity of 5-Aminopyrazole Derivative **54**.


Then, a set of reactions was performed in order to get a better insight into the reaction mechanism. *N*-cinnamyl-*N*-phenylsulfonamide derivative **20** in the presence of CuBr and Li₂CO₃ at 140 °C, with or without Pd-catalyst was recovered unreacted (Scheme 7, a). On the other hand, from **20** and PhSO₂Cl, the Heck type product **58** was selectively obtained in 57% yield (Scheme 7, b). This is probably due to the higher acidity of the C-H bond of the CHPh₂ motif obtained after insertion of the C=C bond into the Ar-Pd bond, which favors the β-H elimination. From *N*-allyl-*N*-(2,4,6-trimethylphenyl)methanesulfonamide **21**, again only the Heck type product **59** was obtained in 38% yield (Scheme 7, c). No sp³ C-H bond activation of the aniline methyl substituents was observed. The presence of a SO₂R substituent on the aniline derivative is also crucial as *N*-allyl-*N*-ethylaniline **60** led to **61** in 68% yield (Scheme 7, d).



33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Scheme 7. Mechanistic Investigations


Finally, labelling experiments were carried out. From deuterated aniline derivative **22**, **62** was obtained in 63% yield, without any incorporation of deuterium (Scheme 8, top). An equimolar mixture of **4** and **22** led to the formation of **31** and **62** in 56:44 ratio showing no significant effect of the use of a deuterated aniline derivative on the reaction rate (Scheme 8, middle). This result suggests that the C-H bond activation step is probably not rate limiting for the catalytic cycle. From the *N*-allyl-*N*-phenylmethanesulfonamide **63** containing two deuterium atoms on the allyl moiety (Scheme 8, bottom), the formation of **64** as a mixture of two diastereomers was observed. The deuterium migration seems to confirm a mechanism involving a β -D elimination followed by a reinsertion in the Pd-D species (See scheme 9).



Scheme 8. Experiments using Deuterated Derivatives

Although the mechanism is not yet elucidated, on the basis of the preliminary mechanistic study and on the previous reports, a catalytic cycle can be proposed (Scheme 9). The first step of is probably the oxidative addition of ArSO₂Cl to a Pd(0) species and release of SO₂ to afford the Pd(II) intermediate **A**, although a Pd(II)/Pd(IV) mechanism is also possible.¹³ Then, **A** affords **B** by coordination of the *N*-allyl-*N*-phenylmethanesulfonamide. Insertion of the allyl C=C bond into the Ar-Pd bond affords **C**. Then, C-H bond activation affords the 6-membered palladacycle **D** and releases LiCl/LiHCO₃. As benzylic C-H are quite acidic, **D** might give **E** via β-H elimination. Finally, reinsertion in the Pd-D bond affords the 7-membered palladacycle **F** which

releases, *via* reductive elimination, the 1,2,3,4-tetrahydroquinoline derivative and regenerates Pd(0).



Scheme 9. Proposed Catalytic Cycle

Formation of 1,2,3,4-tetrahydroquinoline does not result from a Pd-catalyzed cyclisation of the β -elimination product (See, scheme 7, a). Currently available data do not allow to explain the remarkable effect of copper in this reaction. It might work in synergy with palladium for the desulfurative process,¹⁴ or activate palladium.

Conclusions

In summary, we report here the first palladium-catalyzed synthesis of 4-aryl-1,2,3,4-tetrahydroquinoline derivatives *via* successive insertion of an alkene into a Pd-Ar bond followed by an intramolecular sp^2 C-H functionalization. The reaction is chemo- and diastereo-selective and proceeds with an easily accessible phosphine-free air stable palladium catalyst, and Li_2CO_3 as inexpensive base associated to CuBr. The reaction can be successful for a variety of substituents both on aniline, allyl and on the benzenesulfonyl moieties. Due to the described usefulness of 1,2,3,4-tetrahydroquinoline derivatives, such simple reaction conditions offer a new attractive method for access to such structures.

Experimental Section

General procedure for the synthesis of *N*-allyl-*N*-phenylmethanesulfonamides **1-22**, **54**, **63**:

A mixture of the aniline derivative (5 mmol) and triethylamine (0.505 g, 5 mmol) was stirred in CH_2Cl_2 (15 mL) at 22 °C, then, R^2SO_2Cl (5 mmol) was slowly added to the mixture and the reaction was monitored by TLC. When a complete conversion of the starting material was observed, the mixture was concentrated then filtrated through a short silica column and evaporated to afford the desired aryl sulfamide. A mixture of this aryl sulfamide, allyl bromide derivative (7.5 mmol) and K_2CO_3 (1.725 g, 12.5 mmol) in acetone (30 mL) was refluxed overnight. After reaction cooling down, the mixture was concentrated and purified by silica gel chromatography (ethyl acetate/pentane) to afford the desired products **1-22**, **54** and **63**.

General procedure for the synthesis of **24-53**, **55-62** and **64**:

To an oven dried 25 mL Schlenk tube under argon, *N*-allyl-*N*-phenylsulfonamides (0.5 mmol), arenesulfonyl chlorides (1.0 mmol), PdCl₂ (4.4 mg, 0.0025 mmol), CuBr (0.035 g, 0.25 mmol), Li₂CO₃ (0.110 g, 1.5 mmol) and 1,4-dioxane (1.5 mL) were successively added. Then, the reaction mixture was settled in a preheated (140 °C) oil bath for 20 h under stirring. Upon the reaction finished, the crude mixture was purified on silica chromatography (ethyl ether/ pentane, ethyl acetate/ pentane).

4-*p*-Tolyl-1-tosyl-1,2,3,4-tetrahydroquinoline (24): From *N*-allyl-*N*-*p*-tolylbenzenesulfonamide **1** (0.144 g, 0.5 mmol) and 4-methylbenzenesulfonyl chloride (0.191 g, 1 mmol), **24** was obtained in 83% (0.156 g) yield as a white solid: mp 107-109 °C. ¹H NMR (400 MHz, CDCl₃): δ 7.97 (d, *J* = 8.2 Hz, 1H), 7.56 (d, *J* = 8.3 Hz, 2H), 7.29-7.20 (m, 3H), 7.02 (t, *J* = 7.6 Hz, 1H), 6.98 (d, *J* = 7.9 Hz, 2H), 6.78 (d, *J* = 7.7 Hz, 1H), 6.55 (d, *J* = 8.3 Hz, 2H), 4.15-4.07 (m, 1H), 3.83 (dd, *J* = 8.5, 7.0 Hz, 1H), 3.80-3.70 (m, 1H), 2.45 (s, 3H), 2.31 (s, 3H), 2.00-1.88 (m, 1H), 1.77-1.65 (m, 1H). ¹³C NMR (100 MHz, CDCl₃): δ 143.6, 142.0, 137.0, 136.8, 136.0, 132.9, 130.3, 129.7, 129.1, 128.1, 127.4, 126.8, 125.0, 124.6, 45.5, 43.0, 30.3, 21.6, 21.0. Elemental analysis: calcd (%) for C₂₃H₂₃NO₂S (377.50): C 73.18, H 6.14; found: C 73.30, H 6.00.

ASSOCIATED CONTENT

Supporting Information.

Reaction procedures and ¹H and ¹³C NMR of all new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.”

AUTHOR INFORMATION

Corresponding Author

E-mail: jean-francois-soule@univ-rennes1.fr; henri.doucet@univ-rennes1.fr

Author Contributions

The manuscript was written through contributions of all authors. All authors have given approval to the final version of the manuscript.

ACKNOWLEDGMENT

We thank the ministère de la recherche for a fellowship to K. Y. and the Centre National de la Recherche Scientifique and “Rennes Metropole” for providing financial support.

REFERENCES

- (1) For reviews on Pd-catalyzed C-H bond activation: (a) Alberico, D.; Scott, M. E.; Lautens, M. *Chem. Rev.* **2007**, *107*, 174-238; (b) Satoh, T.; Miura, M. *Chem. Lett.* **2007**, *36*, 200-205; (c) Lewis, C.; Bergman, R. G.; Ellman, J. A. *Acc. Chem. Res.* **2008**, *41*, 1013-1025; (d) Bellina, F.; Rossi, R. *Tetrahedron* **2009**, *65*, 10269-10310; (e) Ackermann, L.; Vicente, R.; Kapdi, A. *Angew. Chem. Int. Ed.* **2009**, *48*, 9792-9826; (f) Chen, X.; Engle, K. M.; Wang, D.-H.; Yu, J.-Q. *Angew. Chem. Int. Ed.* **2009**, *48*, 5094-5115; (g) Joucla, L.; Djakovitch, L. *Adv. Synth. Catal.* **2009**, *351*, 673-714; (h) Sun, C.-L.; Li, B.-J.; Shi, Z.-J. *Chem. Commun.* **2010**, *46*, 677-685; (i) Ackermann, L. *Chem. Rev.* **2011**, *111*, 1315-1345; (j) McMurray, L.; O'Hara, F.; Gaunt, M. J. *Chem. Soc. Rev.* **2011**, *40*, 1885-1898; (k) Kuhl, N.; Hopkinson, M. N.; Wencel-Delord, J.; Glorius, F. *Angew. Chem. Int. Ed.* **2012**, *51*, 10236-10254; (l) Yamaguchi, J.; Yamaguchi, A. D.; Itami, K. *Angew. Chem., Int. Ed.* **2012**, *51*, 8960-9009; (m) Neufeldt, S. R.; Sanford, M. S. *Acc. Chem. Res.* **2012**, *45*, 936-946; (n) Wencel-Delord, J.; Glorius, F. *Nature Chem.* **2013**, *5*, 369-375; (o) Rossi, R.; Bellina, F.; Lessi, M.; Manzini, C. *Adv. Synth. Catal.* **2014**, *356*, 17-117; (p) Zhang, M.; Zhang, Y.; Jie, X.; Zhao, H.; Li, G.; Su, W. *Org. Chem. Front.* **2014**, *1*, 843-895; (q) Schranck, J.; Tlili, A.; Beller, M. *Angew. Chem. Int. Ed.* **2014**, *53*, 9426-9428; (r) Bheeter, C. B.; Chen, L.; Soulé, J.-F.; Doucet, H. *Cat. Sci. Technol.* **2016**, *6*, 2005-2049.
- (2) For examples of synthesis of heterocycles via Pd-catalyzed molecular queuing processes: (a) Grigg, R.; Sridharan, V. *Pure. & Appl. Chem.* **1998**, *70*, 1047-1057; (b) De Meijere, A.; Von Zezschwitz, P.; Bräse, S. *Acc. Chem. Res.* **2005**, *38*, 413-422; (c) Gao, R.-D.; Xu, Q.-L.; Dai, L.-X.; You, S.-L. *Org. Biomol. Chem.* **2016**, *14*, 8044-8046. For the synthesis of phenazines via Pd-catalyzed C-H bond activation: (d) Seth, K.; Raha Roy, S.; Chakraborti, A. K. *Chem. Commun.* **2016**, *52*, 922-925.
- (3) (a) Brown, D.; Grigg, R.; Sridharan, V.; Tambyrajah, V. *Tetrahedron Lett.* **1995**, *36*, 1837-1840; (b) Ruck, R. T.; Huffman, M. A.; Kim, M. M.; Shevlin, M.; Kandur, W. V.; Davies, I. W. *Angew. Chem., Int. Ed.* **2008**, *47*, 4711-4714; (c) Satyanarayana, G.; Maichle-Mössmer, C.; Maier, M. E. *Chem. Commun.* **2009**, 1571-1573; (d) René, O.; Lapointe, D.; Fagnou, K. *Org. Lett.* **2009**, *11*, 4560-4563; (e) Ye, S.; Liu, J.; Wu, J. *Chem. Commun.* **2012**, *48*, 5028-5030; For related Pd-catalysed domino Heck C-H functionalization reactions: (f) Tietze, L. F.; Hungerland, T.; Dufert, A.; Objartel, I.; Stalke, D. *Chem. Eur. J.* **2012**, *18*, 3286-3291; (g) Bunescu, A.; Piou, T.; Wang, Q.; Zhu, J. *Org. Lett.* **2015**, *17*, 334-337.

- 1
2
3 (4) For reviews on Heck type reaction: (a) Heck, R. F. *Acc. Chem. Res.* **1979**, *12*, 146-151;
4 (b) de Vries J. G. *Can. J. Chem.* **2001**, *79*, 1086-1092; (c) The Mizoroki–Heck Reaction, ed.
5 Oestreich, M. Wiley, 2009; (d) Mc Cartney, D.; Guiry, P. J. *Chem. Soc. Rev.* **2011**, *40*, 5122-
6 5150.
7
8 (5) For Heck type reaction with benzenesulfonyl chlorides: (a) Miura, M.; Hashimoto, H.;
9 Itoh K.; Nomura, M. *Tetrahedron Lett.* **1989**, *30*, 975-976; (b) Miura, M.; Hashimoto, H.; Itoh
10 K.; Nomura, M. *J. Chem. Soc., Perkin Trans. 1* **1990**, 2207-2211; (c) Dubbaka, S. R.; Vogel, P.
11 *Chem. Eur. J.* **2005**, *11*, 2633-2641.
12
13 (6) For Heck reactions with *N*-allylsulfonamides and aryl halides: (a) Dong, Y.; Busacca, C.
14 A. *J. Org. Chem.* **1997**, *62*, 6464-6465; (b) Proudfoot, J. R.; Patel, U. R.; Dyatkin, A. B. *J. Org.*
15 *Chem.* **1997**, *62*, 1851-1853; (c) Wang, L.; Pan, Y.; Hu, H. *Heterocycl. Commun.* **2012**, *18*, 147-
16 150.
17
18 (7) For a review on transition-metal catalyzed desulfative couplings: Yuan, K.; Soulé, J. F.;
19 Doucet, H. *ACS Catal.* **2015**, *5*, 978-991.
20
21 (8) (a) Zhao, X.; Dimitrijevic E.; Dong, V. M. *J. Am. Chem. Soc.* **2009**, *131*, 3466-3467; (b)
22 Zhao X.; Dong, V. M. *Angew. Chem., Int. Ed.* **2011**, *50*, 932-934; (c) Chen, R.; Liu, S.; Liu, X.;
23 Yang L.; Deng, G.-J. *Org. Biomol. Chem.* **2011**, *9*, 7675-7679.
24
25 (9) For a review on transition-metal mediated C-S bond activation: Wang, L.; He W.; Yu, Z.
26 *Chem. Soc. Rev.* **2013**, *42*, 599-621.
27
28 (10) For a review on the synthesis of 1,2,3,4-tetrahydroquinolines: (a) Katritzky, A. R.;
29 Rachwal, S.; Rachwal, B. *Tetrahedron* **1996**, *52*, 15031-15070; For recent examples of 1,2,3,4-
30 tetrahydroquinolines synthesis: (b) Dorey, G.; Lockhart, B.; Lestage, P.; Casara, P. *Bioorg. Med.*
31 *Chem. Lett.* **2000**, *10*, 935-939; (c) Liu, H.; Dagousset, G.; Masson, G.; Retailleau, P.; Zhu, J. *J.*
32 *Am. Chem. Soc.* **2009**, *131*, 4598-4599; (d) Ramesh, E.; Manian, R. D. R. S.; Raghunathan, R.;
33 Sainath, S.; Raghunathan, M. *Bioorg. Med. Chem.* **2009**, *17*, 660-666; (e) Kouznetsov, V. V.;
34 Arenas, D. R. M.; Arvelo, F.; Forero, J. S. B.; Sojo, F.; Muñoz, A. *Lett. Drug Des. Discovery*
35 **2010**, *7*, 632-639; (f) Yan, M.; Jin, T.; Chen, Q.; Ho, H. E.; Fujita, T.; Chen, L.-Y.; Bao, M.;
36 Chen, M.-W.; Asao, N.; Yamamoto Y. *Org. Lett.*, **2013**, *15*, 1484-1487.
37
38 (11) (a) Larock, R. C.; Berrios-Pena, N. G.; Fried, C. A.; Yum, E. K.; Tu, C.; Leong, W. *J.*
39 *Org. Chem.* **1993**, *58*, 4509-4510; (b) Li, H. Y.; Horn, J.; Campbell, A.; House, D.; Nelson, A.;
40 Marsden, S. P. *Chem. Commun.* **2014**, *50*, 10222-10224; (c) Dai, W.; Jiang, X.-L.; Tao, J.-Y.;
41 Shi, F. *J. Org. Chem.* **2016**, *81*, 185-192.
42
43 (12) (a) Yuan, K.; Doucet, H. *Chem. Sci.* **2014**, *5*, 392-396; (b) Loukotova, L.; Yuan, K.;
44 Doucet, H. *ChemCatChem* **2014**, *6*, 1303-1309; (c) Jin, R.; Yuan, K.; Chatelain, E.; Soulé, J.-F.;
45 Doucet, H. *Adv. Synth. Catal.* **2014**, *356*, 3831-3841.
46
47 (13) For reviews on Pd(II)/Pd(IV) catalysis: (a) Muniz, K. *Angew. Chem. Int. Ed.* **2009**, *48*,
48 9412-9423; (b) Canty, A. J. *Dalton Trans.* **2009**, 10409-10417; (c) Lyons, T. W.; Sanford, M. S.
49 *Chem. Rev.* **2010**, *110*, 1147-1169; (d) Topczewski, J. J.; Sanford, M. S. *Chem. Sci.* **2015**, *6*, 70-
50 76.
51
52 (14) Zeng, X.; Ilies, L.; Nakamura, E. *J. Am. Chem. Soc.* **2011**, *133*, 17638-17640.
53
54
55
56
57
58
59
60

