Reactivity of dirhenium and triruthenium carbonyls toward a biphosphole ligand: M–M, P–P and C–H bonds cleavage

Yomaira Otero, David Coll, Alejandro Arce, Deisy Peña, Franmerly Fuentes, Muriel Hissler, Regis Réau, Rubén Machado, Teresa Gonzalez, Edward Ávila

► To cite this version:

Yomaira Otero, David Coll, Alejandro Arce, Deisy Peña, Franmerly Fuentes, et al.. Reactivity of dirhenium and triruthenium carbonyls toward a biphosphole ligand: M–M, P–P and C–H bonds cleavage. *Journal of Organometallic Chemistry*, 2017, 834, pp.40-46. 10.1016/j.jorganchem.2017.02.012 . hal-01468025

HAL Id: hal-01468025

<https://hal-univ-rennes1.archives-ouvertes.fr/hal-01468025>

Submitted on 4 Jul 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Reactivity of dirhenium and triruthenium carbonyls toward a biphosphole ligand:

M–M, P–P and C–H bonds cleavage

Yomaira Otero^a, David Coll^{a,b}, Alejandro Arce^{a*}, Deisy Peña^a, Franmerly Fuentes^a, Muriel Hissler^{c*}, Regis Réau^c, Rubén Machado^a, Teresa Gonzalez^a, Edward Ávila^a.

^a*Centro de Química, Instituto Venezolano de Investigaciones Científicas (IVIC), Apartado 21827, Caracas 1020-A, Venezuela.*

^b*Escuela Superior Politécnica del Litoral, Centro de Investigación y Desarrollo de Nanotecnología, Ecuador.*

^c*Institut des Sciences Chimiques de Rennes, UMR 6226, Université de Rennes 1, 35042 Rennes Cedex, France.*

*Corresponding authors:

Telephone number: +58 212 504 1322. e-mail address: aarce@ivic.gob.ve

Telephone number: +33 2 23 23 57 83. e-mail address: muriel.hissler@univ-rennes1.fr

Abstract

The reaction of $[\text{Re}_2(\text{CO})_8(\text{CH}_3\text{CN})_2]$ with bis(2-thienyl)biphosphole in refluxing *n*-octane affords the complex $[\text{Re}(\text{CO})_3(\eta^5\text{-PC}_{16}\text{H}_{14}\text{S}_2)]$ (**1**) where cleavage of the Re–Re and P–P bonds occurs; its molecular structure confirms the formation of a rhenium mononuclear compound with a η^5 -coordinated phospholyl unit and three carbonyl groups oriented in a tripod. On the other hand, bis(2-thienyl)biphosphole reacts with $[\text{Ru}_3(\text{CO})_{12}]$ in refluxing cyclohexane to affording two new compounds $[\text{Ru}_3(\text{CO})_9(\mu\text{:}\eta^1\text{:}\eta^5\text{-PC}_{16}\text{H}_{13}\text{S}_2)]$ (**2**) and $[\text{Ru}_4(\text{CO})_{13}(\mu\text{:}\eta^1\text{-PC}_{16}\text{H}_{14}\text{S}_2)_2]$ (**3**). Spectroscopic data and theoretical studies of **2** suggest the formation of an open Ru₃ cluster with two metal–metal bonds bridged by a $(\eta^1\text{:}\eta^5\text{-PC}_{16}\text{H}_{13}\text{S}_2)$ unit and a C–H bond activation of one of the thienyl substituents. X-ray structure of **3** reveals a novel tetranuclear ruthenium cluster containing 13 terminal carbonyl ligands and two phospholyl units coordinated as μ -phosphide.

Keywords: *Biphosphole; metal-metal and phosphorus-phosphorus bonds cleavage; tetranuclear cluster; η^5 -phospholyl complex; C-H activation.*

1. Introduction

Heterocyclic π -complexes have been demonstrated to be excellent ligands for asymmetric catalysis [1-5]. Particularly, phosphole ligands have been interesting for the development of complexes with specific applications since they have the ability to act as both σ - and π -donor ligands, sometimes simultaneously (Figure 1). Several phosphole-complexes have been used in catalytic transformations, such as hydroformylation, hydrosilylation or asymmetric allylic substitution [6-9]. They have also been investigated in fields as diverse as OLEDs, WOLEDs and non linear optics [10-12]. Additionally, phospholyl anions (phospholides) obtained from phosphole derivatives have unique electronic properties since they are good π -acceptors and poor σ -donors [13-16]. They are capable of η^1 - and η^5 -coordinating to metal centers (Figure 1) [17]. Particularly, η^5 -phospholyl complexes have been used in numerous catalytic [4, 13, 14, 18-20].

On the other hand, organometallic compounds that incorporate two or more metal sites in close proximity provide access to reaction pathways not available in mononuclear chemistry [21]. This has been manifested, for example, in reversible metal-metal bond cleavage [22, 23], skeletal rearrangement without degradation [24-26] and ligand activation via multisite coordination [27, 28].

Figure 1. Coordination modes of phosphole and phospholyl ligands

In this regard, our studies have been focused on the reactivity of phosphorus derivatives, such as phosphines and phospholes, toward two or more metal centers, and we have found σ,π -complexes, derivatives containing ring-opened ligands, insertion of a phospholide unit into a metal–metal bond and bridging phosphide ligands [27, 29, 30]. Recently, we have studied the reaction of a variety of π -conjugated phosphole ligands with $[\text{Os}_3(\mu\text{-H})_2(\text{CO})_{10}]$ finding that the binding mode is restricted to σ -coordination, where electronic properties and steric effects of these phosphole ligands play a role in controlling the final product distribution and giving rise to a variety of isomers [31]. We are convinced that the coordination chemistry of phospholes is still a fertile area for further research. Thus, we have extended the study to the coordination chemistry of biphosphole derivatives on direnium and triruthenium carbonyls.

2. Results and discussion

2.1. Thermal treatment of $[\text{Re}_2(\text{CO})_8(\text{CH}_3\text{CN})_2]$ with 2,2'-5,5'-tetra(2-thienyl)-1,1'-biphosphole

The reaction of $[\text{Re}_2(\text{CO})_8(\text{CH}_3\text{CN})_2]$ with biphosphole ligand in refluxing *n*-octane affords one main compound characterized as $[\text{Re}(\text{CO})_3(\eta^5\text{-PC}_{16}\text{H}_{14}\text{S}_2)]$ (**1**), (Scheme 1). Its $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum exhibits a singlet at δ -23.2, which is shifted upfield in comparison to the signal of free ligand (δ -0.1) [32], suggesting that the phosphorus atom is involved in the coordination to the metal center. The ^1H NMR spectrum does not show significant differences compared to those of the free ligand. However, in the $^{13}\text{C}\{^1\text{H}\}$ NMR spectrum, signals for the dienic system are notably upfield shifted, indicating that coordination to the metal center by the dienic system has occurred.

Scheme 1. Reaction of $[\text{Re}_2(\text{CO})_8(\text{CH}_3\text{CN})_2]$ with bis(2-thienyl)biphosphole

The molecular structure of compound **1** was determined by single crystal X-ray diffraction studies (See Figure 2, Table 1), which reveals a $(\eta^5\text{-phospholyl})\text{rhenium}$ complex obtained by cleavage of the Re–Re and P–P bonds of the starting materials. The complex is a rhenium mononuclear with three carbonyl groups in a tripodal orientation and one $\eta^5\text{-PC}_{16}\text{H}_{14}\text{S}_2$ unit, molecular species with three-legged piano-stool geometry. The $\text{PC}_{16}\text{H}_{14}\text{S}_2$ moiety is coordinated as a phospholide to the rhenium atom [$\text{Re1-C1} = 2.36$ (1) Å, $\text{Re1-C2} = 2.327(8)$ Å, $\text{Re1-C7} = 2.347$ (7) Å, $\text{Re1-C8} = 2.337$ (7) Å, $\text{Re1-P1} = 2.530$ (3) Å], resulting in a $\eta^5\text{-complex}$. Phospholyl ring can be considered planar due to that the phosphorus atom lies 0.073 Å out of the mean plane containing the dienic moiety, and the bond delocalization is evidenced by the small deviation in inter atom distances

[C1–C2 = 1.45(1) Å, C2–C7 = 1.43(1) Å, C7–C8 = 1.42(1) Å, P1–C1 = 1.817(9) Å, P1–C8 = 1.80(1) Å], similar to the reported for the analogous manganese derivative [Mn(CO)₃(η⁵-PC₉H₆)] [33].

Figure 2. Molecular structure of **1**, showing 50% probability ellipsoids. Selected bond lengths (Å) and angles (°): Re1–P1 = 2.530 (3), Re1–C1 = 2.36 (1), Re1–C2 = 2.327 (8), Re1–C7 = 2.347 (7), Re1–C8 = 2.337 (7), P1–C1 = 1.817 (9), P1–C8 = 1.80 (1), C1–C2 = 1.45 (1), C2–C7 = 1.43 (1), C7–C8 = 1.42 (1), C11A–Re1–C12A = 91.5 (4), C11A–Re1–C13A = 90.3 (4), C12A–Re1–C13A = 90.7 (4).

Additionally, the (η⁵-phospholyl)rhenium complex formation is corroborated by the presence of five bond critical points (BCP) between the rhenium atom and each element of the ring structure, as displayed in Figure 3. The electron density distribution of a free atom has spherical symmetry. Once this atom is bound, e.g. to form a molecule or a crystal, the electron density loses its spherical uniformity and this new distribution presents different types of critical points. The topological properties of a molecular or crystal charge distribution are summarized by its critical points (CP) [34]. These are points at which the gradient vector field, ∇F(r), vanishes. Among them there are maxima, minima and two types of saddle points. Every CP has a characteristic pattern of trajectories or gradient paths of F(r). The trajectories originate and end at critical points. For our purpose we are interested in two of them, the maxima CP which are associated to the nuclear positions, and one type of saddle points which is associated to the bond critical points (BCP). A BCP is found between every pair of neighboring nuclei. It represents both local maxima in two

directions and a local minimum in the third direction. The network of nuclei and BCP defines a graph, which describes the atomic connectivity and structure within a molecule or a crystal cell. In Figure 3, top and side views are presented for the molecular graph of the ring coordinated to the metal center, showing a “three legs stool” geometrical arrangement. For a better visualization, the rest of the elements in the structure are omitted. The magnitudes of the electron density over each BCP are similar, varying between 0.36 and 0.38 atomic units, which indicates that the electron density is consistently shared between the metal center and phospholyl ring. Theoretical calculations also indicate that the free energy of formation for compound **1**, based on the stoichiometrically balanced reaction, is +48 kcal/mol (Scheme 1).

Figure 3. Molecular graphs (top and side view) of the phosphorus ring coordinated to the metal center in **1**. Gray, red, blue and purple spheres represent C, O, Re and P atoms, respectively. Light green spheres represent the bond critical points (BCPs).

2.2 Thermal treatment of $[\text{Ru}_3(\text{CO})_{12}]$ with 2,2'-5,5'-tetra(2-thienyl)-1,1'-biphosphole

Two new compounds were obtained by reaction of $[\text{Ru}_3(\text{CO})_{12}]$ with 2,2'-5,5'-tetra(2-thienyl)-1,1'-biphosphole in refluxing cyclohexane, $[\text{Ru}_3(\text{CO})_9(\mu:\eta^1:\eta^5\text{-PC}_{16}\text{H}_{13}\text{S}_2)]$ (**2**) and $[\text{Ru}_4(\text{CO})_{13}(\mu:\eta^1\text{-PC}_{16}\text{H}_{14}\text{S}_2)_2]$ (**3**) (Scheme 2). The IR spectrum of compound **2** in the carbonyl region is characteristic of a nonacarbonyl trinuclear compound similar to $[\text{Ru}_3(\text{CO})_9(\mu:\eta^1:\eta^5\text{-PC}_4\text{H}_2\text{Me}_2)(\eta^1\text{-C}_6\text{H}_5)]$ [27]. Its $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum exhibits a singlet at δ 112.2, significantly downfield shifted

123 in comparison to the free ligand, similar to that observed for $[\text{Ru}_3(\text{CO})_9(\mu:\eta^1:\eta^5\text{-PC}_4\text{H}_2\text{Me}_2)(\eta^1\text{-C}_6\text{H}_5)]$ [27]. The ^1H NMR spectrum of compound **2** shows two sets of signals for the thienyl
 124 C_6H_5] [27]. The ^1H NMR spectrum of compound **2** shows two sets of signals for the thienyl
 125 protons, showing inequivalence between both thienyl rings. Absence of the thienyl proton H_3 is also
 126 observed, suggesting the occurrence of C-H bond activation. In the ^{13}C NMR spectrum, signals
 127 assigned to the dienic system of the phosphorus ring are considerably upfieldshifted, similar to
 128 those found for $[\text{Re}_2(\text{CO})_7(\mu:\eta^1:\eta^2:\eta^2\text{-PhPC}_4\text{H}_3\text{Me})]$, $[\text{Ru}_3(\text{CO})_9(\mu:\eta^1:\eta^5\text{-PC}_4\text{H}_2\text{Me}_2)(\eta^1\text{-C}_6\text{H}_5)]$ [27,
 129 29, 30], indicating the coordination of the phospholyl ring to the trinuclear complex. The signal of
 130 $\text{C}_{3\text{thienyl}}$ is downfield shifted, confirming the C-H bond activation. The ESI-MS spectrum of
 131 compound **2** displays parent peaks of high intensity corresponding to its molecular ion $[\text{M}+\text{H}]^+$ at
 132 m/z 858.79, which is in agreement with the molecular formula $[\text{Ru}_3(\text{CO})_9(\text{PC}_{16}\text{H}_{13}\text{S}_2)]$.
 133 Unfortunately, we were unable to obtain suitable crystals of compound **2** for X-ray diffraction
 134 analysis, however, based on the spectroscopic data and its comparison with the ruthenium analogue
 135 $[\text{Ru}_3(\text{CO})_9(\mu:\eta^1:\eta^5\text{-PC}_4\text{H}_2\text{Me}_2)(\eta^1\text{-C}_6\text{H}_5)]$ [27], the proposed molecular structure of
 136 $[\text{Ru}_3(\text{CO})_9(\mu:\eta^1:\eta^5\text{-PC}_{16}\text{H}_{13}\text{S}_2)]$ (**2**) is the most likely as shown in Scheme 2 . Thus, the structure
 137 consists of an open Ru_3 cluster with two metal-metal bonds bridged by a $(\eta^1:\eta^5\text{-PC}_{16}\text{H}_{13}\text{S}_2)$ unit,
 138 obtained probably by triple decarbonylation of $[\text{Ru}_3(\text{CO})_{12}]$, cleavage of a Ru-Ru and P-P bonds
 139 and C-H activation of thienyl ring. Shift to higher values of the IR $\nu(\text{CO})$ frequencies as expected
 140 for an anionic Ru_3 species was not observed for **2**, so an eight-electron donating ligand can be
 141 considered in this case, which makes compound **2** a 50-electron cluster where each ruthenium atom
 142 has the optimal 18-valence electron configuration.

Scheme 2. Reaction of $[\text{Ru}_3(\text{CO})_{12}]$ with bis(2-thienyl)biphosphole

With the aim to complement the experimental characterization, theoretical calculations were performed for compound **2**. Several possible conformers were considered, however, only the most stable conformer was taken into account (see Figure 4). Figure 4.a shows that the ligand is coordinated to one of the Ru atoms through a metal- $\text{C}_{3\text{thienyl}}$ σ -bond and by the phosphorus ring to a second metal center, which is consistent with the propose structure. The coordination by the other two carbons of the thienyl group was considered, but the geometrical stress induced by this arrangement implies the decoordination of the phosphorous ring and/or the P atom with the metal center. In the most stable geometry the phosphorus ring is coordinated in a similar way than compound **1**, through its five atoms (Figure 4.b), confirming the η^5 -coordination and the formation of a (η^5 -phospholyl)ruthenium complex. Magnitudes of the electron density over each BCP between the phosphorus ring and Ru atom vary from 0.41 to 0.49 atomic units, where the smaller value corresponds to the P-Ru interaction. Similar to compound **1**, these BCP values indicate that electron density on the phosphorus ring is noticeably shared with the metal center, which is representative of

covalent interactions. The absence of a bond critical point between the two Ru atoms bound to two and three carbonyl ligands, respectively, indicates that the triruthenium structure is open.

Figure 4. a) Most stable geometry for **2**. b) Molecular graph (side view) of the phosphorus ring coordinated to the metal center in **2**. Gray, red, blue, purple, white and yellow spheres represent C, O, Ru, P, H and S atoms, respectively. Light green spheres represent the bond critical points (BCPs).

The ^{31}P , ^1H and ^{13}C NMR spectra of **3** suggest that a μ -phosphide coordination of the ligand is obtained; its molecular structure was determined by X-ray diffraction studies of crystals obtained by slow evaporation in a cyclohexane-dichloromethane mixture (See Figure 5, Table 1). The structure reveals a novel tetranuclear ruthenium cluster with 13 terminal carbonyl ligands and two phospholyl units, what are obtained from P-P bond cleavage in the biphosphole ligand. This makes each ruthenium atom have an effective atomic number equal to 54, according to EAN rule. The metal core can be described as a kite with four Ru–Ru bonds, two pairs of almost identical sides [Ru1–Ru2 = 3.137 (1) Å, Ru1–Ru4 = 3.136 (1) Å and Ru4–Ru3 = 2.930 (1), Ru2–Ru3 = 2.920 (1) Å] with angles two pairs similar [Ru2–Ru1–Ru4 = 60.16(2)°, Ru4–Ru3–Ru2 = 65.01(2)° and Ru1–Ru4–Ru3 = 117.25(2)°, Ru1–Ru2–Ru3 = 117.51(2)]. The axial carbonyl groups are eclipsed and both $\text{PC}_{16}\text{H}_{14}\text{S}_2$ moieties are coordinated as μ -phosphides. The two phosphorus atoms are nearly positioned in the same plane of the four ruthenium atoms [0.023 Å (P1) and 0.058 Å (P2)]. The Ru2–P1 [2.344(2) Å] and Ru4–P2 [2.340(2) Å] distances are slightly shorter than Ru1–P1 [2.396(2)

179 Å] and Ru1–P2 [2.387(2) Å], indicating that Ru1–Ru2 and Ru1–Ru4 edges are asymmetrically
 180 bridged by a phospholyl unit. The P1 and P2 atoms lie out of the plane of the diene system (0.094 Å
 181 and 0.022 Å respectively), describing the phosphorus rings with an opened envelope-type
 182 conformation.

183
 184 **Figure 5.** Molecular structure of **3**, showing 50% probability ellipsoids. Selected bond lengths (Å) and angles (°): Ru1–
 185 Ru2 = 3.137 (1), Ru1–Ru4 = 3.136 (1), Ru4–Ru3 = 2.930 (1), Ru2–Ru3 = 2.920 (1), Ru2–P1 = 2.344(2) Å, Ru4–P2 =
 186 2.340(2), Ru1–P1 = 2.396(2), Ru1–P2 = 2.387(2), Ru2–Ru1–Ru4 = 60.16(2), Ru4–Ru3–Ru2 = 65.01(2), Ru1–Ru4–Ru3
 187 = 117.25(2), Ru1–Ru2–Ru3 = 117.51(2).

188
 189 On the other hand, the molecular graph containing the metal core and its neighbouring atoms for
 190 compound **3** is shown in Figure 6. This graph displays two BCPs between each P atom and the two
 191 closest ruthenium atoms, corroborating the μ -phosphide coordination of the ligand. Interestingly,
 192 and in contrast to the X-ray molecular structure shown in Figure 5, Ru2 and Ru4 are bound while
 193 Ru1–Ru2 and Ru1–Ru4 bonds are not present, even though the observed internuclear Ru2–Ru4
 194 distance [3.144(1) Å] is longer than those for Ru1–Ru2 [3.137(1) Å] and Ru1–Ru4 [3.136(1) Å].
 195 As revealed in Figure 6, the BCP electron density value of Ru2–Ru4 bond is found to be smaller
 196 ($0.174 \text{ e}/\text{\AA}^3$) than the ρ_{BCP} of Ru2–Ru3 and Ru3–Ru4 bonds (0.245 and $0.246 \text{ e}/\text{\AA}^3$, respectively),
 197 indicating that no Ru2–Ru4 bond exists. This is consistent with the corresponding internuclear
 198 distances found with X-ray diffraction analyses.

199

Figure 6. Molecular graph for the metal core of **3**. The BCP electron density values for Ru2–Ru4, Ru3–Ru4 and Ru2–Ru3 bonds are given in $e/\text{\AA}^3$. Bond distances are as presented in Table 1. Gray, red, blue and purple spheres represent C, O, Ru and P atoms, respectively. Light green spheres represent the bond critical points (BCPs).

3. Conclusions

A new (η^5 -phospholyl)rhenium complex $[\text{Re}(\text{CO})_3(\eta^5\text{-PC}_{16}\text{H}_{14}\text{S}_2)]$ (**1**) was obtained by the rhenium-rhenium and phosphorus-phosphorus bonds cleavage, when $[\text{Re}_2(\text{CO})_8(\text{CH}_3\text{CN})_2]$ reacts with bis(2-thienyl)biphosphole. Additionally, two new ruthenium carbonyl clusters were characterized as $[\text{Ru}_3(\text{CO})_9(\mu\text{-}\eta^1\text{-}\eta^5\text{-PC}_{16}\text{H}_{13}\text{S}_2)]$ (**2**) and $[\text{Ru}_4(\text{CO})_{13}(\mu\text{-}\eta^1\text{-PC}_{16}\text{H}_{14}\text{S}_2)_2]$ (**3**), wherein compound **2** represents the second example of a phospholide unit inserted into a metal–metal bond, while compound **3** is the first example of a tetranuclear cluster with two π -conjugated phospholyl units coordinated as μ -phosphides. These results open up the possibility for a facile access to new ruthenium and rhenium complexes with phospholyl units as bridging ligands which can be used in catalytic reactions.

4. Experimental section

4.1 General remarks

[Re₂(CO)₈(CH₃CN)₂] and 2,2'-5,5'-tetra(2-thienyl)-1,1'-biphosphole were synthesized as previously described [32, 35]. [Ru₃(CO)₁₂] was used as supplied by Aldrich Chemical Company. ¹H, ¹³C and ³¹P NMR spectra were recorded using Bruker Avance AM300 and AM500 spectrometers, and assignment of carbon chemical shifts was based on HMBC and HMQC experiment. Infrared spectra were recorded as cyclohexane solutions using a 2 mm CaF₂ cell on a Spectrum 100 series Perkin Elmer spectrophotometer. Mass spectra were acquired on a Thermo Scientific TSQ Quantum Ultra AM Triple Quadrupole mass spectrometer employing the Heated Electrospray Ionization (HESI) technique. Routine separation of products in air was by thin-layer chromatography on plates coated with Merck Kieselgel 60 GF₂₅₄.

4.2 Reaction of [Re₂(CO)₈(CH₃CN)₂] with 2,2'-5,5'-tetra(2-thienyl)-1,1'-biphosphole

A solution of [Re₂(CO)₈(CH₃CN)₂] (100 mg, 0.147 mmol) and 2,2'-5,5'-tetra(2-thienyl)-1,1'-biphosphole (89 mg, 0.147 mmol) in dry *n*-octane (50mL) was taken to reflux under nitrogen for 3h. After evaporation of the solvent, TLC (SiO₂) of the brown solid residue (hexane/dichloromethane, 9:1 v/v) gave [Re(CO)₃(η⁵-PC₁₆H₁₄S₂)] (**1**) (89% yield). Spectral data for **1**: IR (ν_{CO}, cyclohexane): 2026 (s), 1949 (s), 1935 (s) cm⁻¹. ¹H NMR (500 MHz, CD₂Cl₂): δ= 1.93 (m, 2H, CH₂-CH₂-C=C), 2.09 (m, 2H, CH₂-CH₂-C=C), 3.05 (m, 2H, CH₂-C=C), 3.17 (m, 2H, CH₂-C=C), 7.03 (d, ⁴J_{H5-H3} = 1.6 Hz, 2H, H₃ thienyl), 7.04 (d, ³J_{H5-H4} = 4.7 Hz, 2H, H₄ thienyl), 7.34 (dd, ³J_{H4-H5} = 4.7 Hz, ⁴J_{H3-H5} = 1.6 Hz, 2H, H₅ thienyl). ¹³C {¹H} NMR (500 MHz, CD₂Cl₂): δ= 22.6 (s, CH₂-CH₂-C=C), 26.3 (s, CH₂-C=C), 109.1 (d, J_{P-C} = 23.0 Hz, C_α), 114.6 (d, ²J_{P-C} = 4.9 Hz, C_β), 126.6 (s, C₅ thienyl), 127.3 (s, C₄ thienyl), 128.5 (d, ³J_{P-C} = 6.8 Hz, C₃ thienyl), 136.2 (d, ²J_{P-C} = 20.3 Hz, C₂ thienyl), 194.4 (s, M-CO). ³¹P {¹H} NMR (500 MHz, CDCl₃): δ= -23.2 (s). MS: *m/z* 373.24 [M+H]⁺.

4.3 Reaction of $[Ru_3(CO)_{12}]$ with 2,2'-5,5'-tetra(2-thienyl)-1,1'-biphosphole

A solution of $[Ru_3(CO)_{12}]$ (100 mg, 0.156 mmol) and 2,2'-5,5'-tetra(2-thienyl)-1,1'-biphosphole (94 mg, 0.156 mmol) in dry cyclohexane (50mL) was taken to reflux under nitrogen for 2h. After evaporation of the solvent, TLC (SiO_2) of the brown solid residue (hexane/dichloromethane, 8:2 v/v) gave two compounds $[Ru_3(CO)_9(\mu:\eta^1:\eta^5-PC_{16}H_{13}S_2)]$ (**2**) (19% yield), and $[Ru_4(CO)_{13}(\mu:\eta^1-PC_{16}H_{13}S_2)_2]$ (**3**) (35% yield). Spectral data for **2**: IR (ν_{CO} , cyclohexane): 2109 (m), 2063 (s), 2048 (s), 2026 (s), 2016 (m), 1999 (m), 1984 (w), 1949 (w) cm^{-1} . 1H NMR (300 MHz, CD_2Cl_2): δ = 2.16 (m, 2H, $CH_2-CH_2-C=C$), 2.31 (m, 2H, $CH_2-CH_2-C=C$), 2.81 (m, 1H, $CH_2-C=C$), 3.08 (m, 2H, $^*CH_2-C=C$), 3.30 (m, 1H, $CH_2-C=C$), 6.82 (dd, $^3J_{H4-H5}$ = 4.8 Hz, $^3J_{H4-H3}$ = 3.6 Hz, 1H, H_4 thienyl), 6.88 (dd, $^3J_{H3-H4}$ = 3.6 Hz, $^4J_{H3-H5}$ = 1.1 Hz, 1H, H_3 thienyl), 7.02 (d, $^3J_{H4^*-H5^*}$ = 4.9 Hz, 1H, H_{4^*} thienyl), 7.17 (dd, $^3J_{H5-H4}$ = 4.8 Hz, $^4J_{H5-H3}$ = 1.8 Hz, 1H, H_5 thienyl), 7.25 (dd, $^3J_{H5^*-H4^*}$ = 4.9 Hz, $^5J_{H5^*-P}$ = 1.3 Hz, 1H, H_{5^*} thienyl). $^{13}C\{^1H\}$ NMR (300 MHz, CD_2Cl_2): δ = 23.0 (s, $CH_2-CH_2-C=C$), 23.7 (s, $^*CH_2-CH_2-C=C$), 25.4 (d, $^2J_{P-C}$ = 5.3 Hz, $^*CH_2-C=C$), 26.2 (d, $^2J_{P-C}$ = 3.9 Hz, $CH_2-C=C$), 54.5 (s, C_{6^*} thienyl), 57.2 (s, C_6 thienyl), 98.6 (s, C_{7^*} thienyl), 98.9 (s, C_7 thienyl), 127.1 (s, C_{5^*} thienyl), 127.2 (s, C_5 thienyl), 127.7 (s, C_4 thienyl), 130.0 (d, $^3J_{P-C}$ = 7.2 Hz, C_3 thienyl), 135.6 (s, C_2 thienyl), 135.8 (s, C_{2^*} thienyl), 137.4 (s, C_4 thienyl), 160.6 (s, C_{3^*} thienyl), 195.4 (s, M-CO), 203.2 (s, M-CO), 205.9 (s, M-CO). $^{31}P\{^1H\}$ NMR (500 MHz, CD_2Cl_2): δ = 112.2 (s). MS: 858.79 m/z $[M+H]^+$. Spectral data for **3**: IR (ν_{CO} , cyclohexane): 2105 (w), 2079 (m), 2042 (s), 2015 (m), 1995 (m), 1975 (w) cm^{-1} . 1H NMR (500 MHz, CD_2Cl_2): δ = 1.73 (m, 4H, $CH_2-CH_2-C=C$), 2.44 (m, 4H, $CH_2-C=C$), 6.90 (dd, $^4J_{H5-H3}$ = 1.0 Hz, $^3J_{H4-H3}$ = 3.5 Hz, 2H, H_3 thienyl), 6.97 (dd, $^3J_{H5-H4}$ = 5.1 Hz, $^3J_{H3-H4}$ = 3.5 Hz, 2H, H_4 thienyl), 7.39 (dd, $^3J_{H4-H5}$ = 5.1 Hz, $^4J_{H3-H5}$ = 1.0 Hz, 2H, H_5 thienyl). $^{13}C\{^1H\}$ NMR (500 MHz, CD_2Cl_2): δ = 22.8 (s, $CH_2-CH_2-C=C$), 28.4 (s, $CH_2-C=C$), 127.2 (s, C_4 thienyl), 127.6 (s, C_5 thienyl), 129.6 (s, C_3 thienyl), 136.4 (s, C_2 thienyl), 139.2 (s, C_α), 150.3 (s, C_β), 206.2 (s, M-CO). $^{31}P\{^1H\}$ NMR (500 MHz, $CDCl_3$): δ = 84.4 (s). MS: m/z 1371.00 $[M+H]^+$.

266

267 *4.4 Crystal structure determination for 1 and 3*

268 Intensity data were recorded at room temperature on a Rigaku AFC-7S diffractometer equipped
269 with a Mercury CCD detector using monochromated Mo(K α) radiation ($\lambda = 0.71070$ Å). An
270 empirical absorption correction (multi-scan) was applied using the CrystalClear package. The low
271 completeness ratio is due to the experimental setup whereby by the equipment have a ω circle and
272 an added area detector (four-circle diffractometer modified with a CCD). This precludes the
273 collection of some regions of reciprocal lattice space and lowers the completeness. In order to
274 compensate, additional redundant data were measured. The structures were solved by direct
275 methods and refined by full-matrix least-squares on F² using the SHELXTL-PLUS package. All
276 non hydrogen atoms were refined anisotropically, and hydrogen atoms were added at calculated
277 positions (C-H = 0.93–0.97 Å) and refined as riding with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}$. One crystallographically
278 independent dichloromethane molecule located on special positions was found disordered. Attempts
279 were made to model this disorder or split it into two positions, but were unsuccessful.
280 PLATON/SQUEZZE routine was used to correct the data for the presence of two disordered
281 solvent. A potential solvent volume of 461 Å³ was found. The stoichiometry of the solvent was
282 calculated to be 1 molecule of dichloromethane per formula unit, which results in a total of 125
283 electrons per unit cell. This molecule was used to calculate expected molecular weight, DXcalc and
284 F(000). Additionally, we have managed to solve the disorder of the S4 atom with a thorough
285 revision of the map of Fourier differences, managing to model not only the disorder of the atom S4
286 but the complete thienyl group to which it belongs; C29-S4-C30-C31-C32 atoms, which group has
287 a 60.8% disorder. Therefore, considering this disorder we find that the bond distance for the C29-
288 C30 is 1.434 Å, whose distance is inside the maximum (1.819 Å) and minimum (1.053 Å) values
289 and average of 1.343(36) Å for similar structures in a search made with Mogul 1.7.1 build RC2.

Crystal and refinement information for compounds **1** and **3** is showed in Table 1. CCDC 1497908 and 1497909 contain the supplementary crystallographic data for compounds **1** and **3**, respectively. These data can be obtained free of charge from The Cambridge Crystallographic Data Center via www.ccdc.cam.ac.uk/data_request/cif.

Table 1. Experimental details for X-ray diffraction collected data and structural refinement for compounds **1** and **3**.

	[Re(CO)₃(η^5-PC₁₆H₁₄S₂)] (1)	[Ru₄(CO)₁₃(μ-PC₁₆H₁₃S₂)₂] (3)
Chemical formula	C ₁₉ H ₁₄ O ₃ PRE ₂ S ₂	C _{46.5} H ₃₁ Cl ₃ O ₁₃ P ₂ Ru ₄ S ₄
Formula weight	571.59	1498.52
Crystal system, Space group	Triclinic, <i>P</i> -1 (No. 2)	Triclinic, <i>P</i> -1 (No. 2)
a(Å), $\alpha(^{\circ})$	9.746(6), 98.363(9)	12.058(2), 74.21(3)
b(Å), $\beta(^{\circ})$	10.127(5), 111.609(15)	14.584(3), 71.28(2)
c(Å), $\gamma(^{\circ})$	10.340(6), 91.830(18)	17.487(3), 72.92(2)
V(Å ³), Z	934.7(9), 2	2729.8(11), 2
dx (g cm ⁻³)	2.031	1.823
Crystal size (mm)	0.05 × 0.14 × 0.16	0.44 × 0.40 × 0.36
F(000)	548	1470
μ (mm ⁻¹)	6.826	1.503
θ range ($^{\circ}$)	2.0414 – 27.4333	1.254 – 27.573
Reflections Collected	10362	25543
Reflections Unique(<i>R</i> _{int})	3359	9350
Reflections With <i>I</i> > 2 σ (<i>I</i>)	3084	7075
Number of parameters	235	629
R(<i>F</i> ²) [<i>I</i> > 2 σ (<i>I</i>)]	0.046	0.050
wR(<i>F</i> ²) [<i>I</i> > 2 σ (<i>I</i>)]	0.120	0.155
Goodness of fit on <i>F</i> ²	1.07	1.10
Max/min $\Delta\rho$ (e Å ⁻³)	1.82 – 2.43	1.60 – 0.83

4.5 Theoretical calculations

All structures were optimized using DMol³ [36–38]. This DFT based program permits determination of the relative stability of all studied species based on their electronic structure. The calculations were performed using the Kohn–Sham Hamiltonian with the Perdew–Wang 1991 gradient correction [39] and the double-zeta plus (DNP) numerical basic set [36, 37, 40], which provides good accuracy at a relatively low computational cost. The All Electron core treatment was used for all the atoms. Frequency calculations of the structures showed that all frequencies were positive

indicating that all structures are real minima. The presence and nature of the chemical bonds, has been analyzed using the quantum theory of atoms in molecules (QTAIM) [34] and the AIM-UC code [41].

Acknowledgements

We thank FONACIT and Laboratorio Nacional de Difracción de Rayos-X for funding the projects G-20050000447 and LAB-97000821, respectively. Additionally, we thank to the Programa de Cooperación de Post-grado (PCP) between France and Venezuela.

References

- [1] C. Ganter, The chemistry of chiral heterometalloenes, *J. Chem. Soc., Dalton Trans.* (24) (2001) 3541-3548.
- [2] F.O. Mathey, Phosphorus-carbon heterocyclic chemistry : the rise of a new domain, Pergamon, Amsterdam; New York, 2001.
- [3] A.J. Ashe III, S. Al-Ahmad, Diheteroferrocenes and Related Derivatives of the Group 15 Elements: Arsenic, Antimony, and Bismuth, in: A.S. F. Gordon, W. Robert (Eds.), *Adv. Organomet. Chem.*, Academic Press 1996, pp. 325-353.
- [4] F. Mathey, The chemistry of phospho- and polyphosphacyclopentadienide anions, *Coord. Chem. Rev.* 137 (1994) 1-52.
- [5] K.B. Dillon, Phosphorus : the carbon copy : from organophosphorus to phospho-organic chemistry, in: F. Mathey, J.F. Nixon (Eds.) *John Wiley*, Chichester, England ;, 1998.
- [6] G. Mora, B. Deschamps, S. van Zutphen, X.F. Le Goff, L. Ricard, P. Le Floch, Xanthene-Phosphole Ligands: Synthesis, Coordination Chemistry, and Activity in the Palladium-Catalyzed Amine Allylation, *Organometallics* 26(8) (2007) 1846-1855.
- [7] S. Doherty, E.G. Robins, J.G. Knight, C.R. Newman, B. Rhodes, P.A. Champkin, W. Clegg, Selectivity for the methoxycarbonylation of ethylene versus CO/ethylene copolymerization with catalysts based on C4-bridged bidentate phosphines and phospholes, *J. Organomet. Chem.* 640(1-2) (2001) 182-196.
- [8] J. Hydrio, M. Gouygou, F. Dallemer, J.-C. Daran, G.G.A. Balavoine, New chiral phosphole ligands: their coordination behaviour and application in palladium-catalysed asymmetric allylic substitution, *Tetrahedron: Asymmetry* 13(10) (2002) 1097-1102.
- [9] G. Keglevich, T. Kégl, T. Chuluunbaatar, B. Dajka, P. Mátyus, B. Balogh, L. Kollár, Hydroformylation of styrene in the presence of rhodium-2,4,6-trialkylphenyl-phosphole in situ catalytic systems, *J. Mol. Catal. A: Chem.* 200(1-2) (2003) 131-136.
- [10] P.-A. Bouit, A. Escande, R. Szűcs, D. Szieberth, C. Lescop, L. Nyulászi, M. Hissler, R. Réau, Dibenzophosphapentaphenes: Exploiting P Chemistry for Gap Fine-Tuning and Coordination-Driven Assembly of Planar Polycyclic Aromatic Hydrocarbons, *J. Am. Chem. Soc.* 134(15) (2012) 6524-6527.
- [11] H. Chen, W. Delaunay, L. Yu, D. Joly, Z. Wang, J. Li, Z. Wang, C. Lescop, D. Tondelier, B. Geffroy, Z. Duan, M. Hissler, F. Mathey, R. Réau, 2,2'-Biphospholes: Building Blocks for Tuning the HOMO-LUMO Gap of π -Systems Using Covalent Bonding and Metal Coordination, *Angew. Chem. Int. Ed.* 51(1) (2012) 214-217.
- [12] J. Crassous, R. Reau, π -Conjugated phosphole derivatives: synthesis, optoelectronic functions and coordination chemistry, *Dalton Transactions* (48) (2008) 6865-6876.
- [13] F. Mathey, Phosphametalloenes: from discovery to applications, *J. Organomet. Chem.* 646(1-2) (2002) 15-20.
- [14] P.L. Floch, Phosphaalkene, phospholyl and phosphinine ligands: New tools in coordination chemistry and catalysis, *Coord. Chem. Rev.* 250(5-6) (2006) 627-681.

- [15] M. Ogasawara, S. Arae, S. Watanabe, V. Subbarayan, H. Sato, T. Takahashi, Synthesis and Characterization of Benzo[b]phosphaferrocene Derivatives, *Organometallics* 32(17) (2013) 4997-5000.
- [16] H.-C. Su, O. Fadhel, C.-J. Yang, T.-Y. Cho, C. Fave, M. Hissler, C.-C. Wu, R. Réau, Toward Functional π -Conjugated Organophosphorus Materials: Design of Phosphole-Based Oligomers for Electroluminescent Devices, *J. Am. Chem. Soc.* 128(3) (2006) 983-995.
- [17] Y.J. Ahn, R.J. Rubio, T.K. Hollis, F.S. Tham, B. Donnadieu, Slip-Inversion-Slip Mechanism of Phosphametallocene Isomerization. Spectroscopic Characterization of an (η^1 -Phospholyl)titanium Complex. Synthesis and Structures of Chiral Mono(phospholyl)titanium Complexes, *Organometallics* 25(5) (2006) 1079-1083.
- [18] D. Carmichael, F. Mathey, L. Ricard, N. Seeboth, Synthesis of 2-silyl substituted phospharuthenocenes and an elaboration into the first phospharuthenocene-phosphine, *Chem. Commun.* (24) (2002) 2976-2977.
- [19] L. Weber, Phosphorus Heterocycles: From Laboratory Curiosities to Ligands in Highly Efficient Catalysts, *Angew. Chem. Int. Ed.* 41(4) (2002) 563-572.
- [20] F. Nief, F. Mathey, L. Ricard, F. Robert, Coordination chemistry of the new 2,3,4,5-tetramethylphospholyl (C4Me4P) .pi.-ligand. Crystal and molecular structure of (.eta.5-C4Me4P)2ZrCl2.cntdot.1/2C10H8, *Organometallics* 7(4) (1988) 921-926.
- [21] P. Braunstein, Workshop on Clusters and Surfaces Interplay between bridging groups and metal-metal bonds in heterometallic clusters, *Mater. Chem. Phys.* 29(1) (1991) 33-63.
- [22] G. Huttner, J. Schneider, H.-D. Müller, G. Mohr, J. von Seyerl, L. Wohlfahrt, Reversible Opening of a Trinuclear Heterometal Cluster, *Angewandte Chemie International Edition in English* 18(1) (1979) 76-77.
- [23] F. Richter, H. Vahrenkamp, Basic cluster reactions. 1. Reversible unfolding of FeCo2Mo and FeCo2W clusters, *Organometallics* 1(5) (1982) 756-757.
- [24] M.I. Bruce, J.G. Matison, J.R. Rodgers, R.C. Wallis, Cluster chemistry. Synthesis and X-ray structure of [Ru5(CO)14(CNBut)2], containing an unusual open Ru5 cluster, *J. Chem. Soc., Chem. Commun.* (20) (1981) 1070-1071.
- [25] D.H. Farrar, P.F. Jackson, B.F.G. Johnson, J. Lewis, J.N. Nicholls, M. McPartlin, A high-yield synthesis of Ru5C(CO)15 by the carbonylation of Ru6C(CO)17; the X-ray structure analyses of Ru5C(CO)15 and Ru5C(CO)14PPh3, *J. Chem. Soc., Chem. Commun.* (9) (1981) 415-416.
- [26] R.D. Adams, M. Chen, X. Yang, Iridium-Gold Cluster Compounds: Syntheses, Structures, and an Unusual Ligand-Induced Skeletal Rearrangement, *Organometallics* 31(9) (2012) 3588-3598.
- [27] A.J. Arce, Y. De Sanctis, M.C. Goite, R. Machado, Y. Otero, T. Gonzalez, Insertion of a phospholide unit into a metal-metal bond: Synthesis and X-ray structure of [Ru3(CO)9(μ : η^1 : η^5 -PC4H2Me2)(η^1 -C6H5)], *Inorg. Chim. Acta* 392(0) (2012) 241-245.
- [28] A.J. Arce, Y. De Sanctis, J. Manzur, A.J. Deeming, N.I. Powell, Oxidative addition of phenylphosphole: X-ray crystal structures of two ring-opened products [Os3(μ 3-PhPC4H4)(CO)x] where x = 8 or 9, *J. Organomet. Chem.* 408(1) (1991) C18-C21.
- [29] Y. Otero, A. Arce, Y.D. Sanctis, R. Machado, M.C. Goite, T. Gonzalez, A. Briceño, High versatility of 3,4-dimethyl-1-phenyl-phosphole and 3-methyl-1-phenyl-phosphole as ligands in the reaction with [Re2(CO)8(CH3CN)2] and [Ru3(CO)12]: X-ray structures of [Re2(CO)7(η^1 : η^2 : η^2 -PhPC4H2Me2)], [Re2(CO)7(η^1 : η^2 : η^2 -PhPC4H3Me)] and [Ru2(CO)4(PhPC4H3Me)2], *Inorg. Chim. Acta* 404(0) (2013) 77-81.
- [30] Y. Otero, D. Peña, Y. De Sanctis, A. Arce, E. Ocando-Mavarez, R. Machado, T. Gonzalez, Reactivity of triosmium clusters with 3,4-dimethyl-1-phenylphosphole and cyanoethyl-di-tert-butylphosphine ligands: X-ray crystal structures of [Os3(CO)9(μ -OH)(μ -H)(η^1 -PhPC4H2Me2)] and [Os3(CO)11(η^1 -t Bu2PC2H4CN)], *Transition Met. Chem.* 39(2) (2014) 239-246.
- [31] Y. Otero, D. Peña, A. Arce, M. Hissler, R. Réau, Y. De Sanctis, E. Ocando-Mavárez, R. Machado, T. González, Fluxional behaviour of phosphole and phosphine ligands on triosmium clusters, *J. Organomet. Chem.* 799-800 (2015) 45-53.
- [32] C. Fave, M. Hissler, T. Kárpáti, J. Rault-Berthelot, V. Deborde, L. Toupet, L. Nyulászi, R. Réau, Connecting π -Chromophores by σ -P-P Bonds: New Type of Assemblies Exhibiting σ - π -Conjugation, *J. Am. Chem. Soc.* 126(19) (2004) 6058-6063.
- [33] A. Decken, F. Bottomley, B.E. Wilkins, E.D. Gill, Organometallic Complexes of Benzannelated Phospholyls: Synthesis and Characterization of Benzophospholyl and the First iso-Benzophospholyl Metal Complexes, *Organometallics* 23(15) (2004) 3683-3693.
- [34] R.F.W. Bader, *Atoms in molecules—a quantum theory*, Clarendon Press, Oxford, *Atoms in molecules—a quantum theory*, Clarendon Press, Oxford (1990).
- [35] M.I. Bruce, P.J. Low, Expeditious synthesis of Re3(μ -H)3(CO)11(NCMe), *J. Organomet. Chem.* 519(1) (1996) 221-222.

- [36] A. Inc., DMol3 is available as part of Material Studio San Diego, USA, DMol3 is available as part of Material Studio San Diego, USA (2010).
- [37] B. Delley, An all-electron numerical method for solving the local density functional for polyatomic molecules, The Journal of Chemical Physics 92(1) (1990) 508-517.
- [38] S. Aime, R. Gobetto, E. Valls, NMR characterization of four new isomers of $H(\mu-H)Os_3(CO)_{10}(\text{phosphine})$, Inorg. Chim. Acta 275–276(0) (1998) 521-527.
- [39] J.P. Perdew, Y. Wang, Accurate and simple analytic representation of the electron-gas correlation energy, Physical Review B 45(23) (1992) 13244-13249.
- [40] B. Delley, From molecules to solids with the DMol3 approach, The Journal of Chemical Physics 113(18) (2000) 7756-7764.
- [41] D. Vega, D. Almeida, J. Comput. Methods Sci. Eng., J. Comput. Methods Sci. Eng. 14, 1-3 (2014) 131-136.

(η^5 -phospholyl)rhenium complex is obtained from the Re–Re and P–P bonds cleavage.

A phospholyl unit inserted into a M–M bond represents the 2nd example.

A new tetranuclear cluster with two μ -phosphides units is obtained.

A ruthenium tetranuclear cluster with two π -conjugated phospholyl units coordinated as μ -phosphide and a (η^5 -phospholyl)rhenium complex are obtained from M-M and P-P bonds cleavage.