Dysmetabolic iron overload syndrome: a mild but unquestionable iron overload is identified by MRI

Fabrice Lainé, Edouard Bardou-Jacquet, Anita Paisant, Yves Gandon, Yves Deugnier

► To cite this version:

Fabrice Lainé, Edouard Bardou-Jacquet, Anita Paisant, Yves Gandon, Yves Deugnier. Dysmetabolic iron overload syndrome: a mild but unquestionable iron overload is identified by MRI. *Hepatology*, 2017, 65 (6), pp.2119-2120. 10.1002/hep.29032 . hal-01531023

HAL Id: hal-01531023

<https://hal-univ-rennes1.archives-ouvertes.fr/hal-01531023>

Submitted on 1 Jun 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Dysmetabolic iron overload syndrome: a mild but unquestionable iron overload is identified by MRI

Authors and affiliations:

Lainé Fabrice^{1,2}, Edouard Bardou-Jacquet¹⁻³, Paisant Anita^{1,3,4}, Yves Gandon^{3,4}, Yves Deugnier¹⁻³

¹ INSERM CIC 1414, Rennes, France

² Liver Unit, CHU Pontchaillou, Rennes, France

³ University of Rennes1, Rennes, France

⁴ Department of Abdominal Imaging, CHU Pontchaillou, Rennes, France

Dear Editor,

Castiella et al claim that liver iron concentration (LIC) is not increased in patients with hyperferritinemia and metabolic syndrome on the basis of a study¹ where LIC was measured using MRI in 54 patients referred for hyperferritinemia. This study has a major limitation since 5% of patients had viral hepatitis and 35% were excessive drinkers (>40g/day), two conditions that can induce hyperferritinemia without iron overload.

Dysmetabolic Iron Overload Syndrome (DIOS) is a well-defined condition where usual causes of hyperferritinemia (alcohol, viral hepatitis, genetic hemochromatosis...) have been excluded. Hyperferritinemia is associated with proven hepatic iron overload and one or more features of the metabolic syndrome. In the initial description of DIOS, the reality of hepatic

iron excess was demonstrated by histology, determination of biochemical LIC and calculation of iron removed ².

In our study ³, LIC was determined by MRI according to the widely used method of Gandon et al ⁴. The threshold of 50 $\mu\text{mol/g}$ was chosen as inclusion criterion because it was previously demonstrated to ensure hepatic iron excess with good specificity. Moreover, in our randomized study, the number of grams of iron subtracted by phlebotomy, the gold standard for the determination of body iron stores, was calculated in 126 patients. A mean of 12 ± 4 bi-monthly phlebotomies, corresponding to 4.9 ± 1.6 l of blood removed, i.e. 2.45 ± 0.8 g of iron, were necessary to obtain low ferritin levels in these patients. This was in line with results of Jezequel⁵ et al who showed in a prospective matched-controlled study that patients diagnosed as having DIOS on the basis of hyperferritinemia and $\text{LIC} > 50 \mu\text{mol/g}$ at MRI had four times higher total body iron stores ($2.5 \pm 0.7 \text{g}$) than overweight controls with normal serum ferritin ($0.8 \pm 0.3 \text{g}$).

In conclusion, our study included a large sample of strictly selected patients representative of DIOS. These patients had mild but unquestionable iron overload.

1. Castiella A, Zapata E, Zubiaurre L, Iribarren A, Alústiza JM, Otazua P, et al. Liver iron concentration is not raised in patients with dysmetabolic hyperferritinemia. *Ann Hepatol* 2016;15:540-544.
2. Mendler MH, Turlin B, Moirand R, Jouanolle AM, Sapey T, Guyader D et al. Insulin resistance-associated hepatic iron overload. *Gastroenterology* 1999; 117:1155-63.

3. Lainé F, Tuivard M, Loustaud-Ratti V, Bonnet F, Calès P, Bardou-Jacquet E, et al. Metabolic and hepatic effects of bloodletting in dysmetabolic iron overload syndrome: a randomized controlled study in 274 patients. *Hepatology* 2016 Sep29; doi: 10.1002/hep.28856. [Epub ahead of print].
4. Gandon Y, Olivie D, Guyader D, Aube C, Oberti F, Sebille V, et al. Noninvasive assessment of hepatic iron stores by MRI. *Lancet* 2004;363:357-362.
5. Jezequel C, Lainé F, Laviolle B, Kiani A, Bardou-Jacquet E, Deugnier Y. Both hepatic and body iron stores are increased in dysmetabolic iron overload syndrome. A case-control study. *PLoS One* 2015; 10(6): e0128530.