

Mechanically Isolated Stromal Vascular Fraction by Nanofat Emulsification Techniques

Nicolas Bertheuil, Audrey Varin, Raphael Carloni, Paul Girard, Benoit
Chaput

► **To cite this version:**

Nicolas Bertheuil, Audrey Varin, Raphael Carloni, Paul Girard, Benoit Chaput. Mechanically Isolated Stromal Vascular Fraction by Nanofat Emulsification Techniques. *Plastic and Reconstructive Surgery*, Lippincott, Williams & Wilkins, 2017, 140 (3), pp.508e-509e. 10.1097/PRS.0000000000003630 . hal-01579829

HAL Id: hal-01579829

<https://hal-univ-rennes1.archives-ouvertes.fr/hal-01579829>

Submitted on 31 Aug 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Plastic and Reconstructive Surgery Advance Online Article

DOI: 10.1097/PRS.0000000000003630

Mechanically isolated Stromal Vascular Fraction by Nanofat Emulsification techniques.

Nicolas Bertheuil, MD^{1,2,3,4}, Audrey Varin, PhD, Raphael Carloni, MD¹, Paul Girard, MD¹,
Benoit Chaput, MD^{4,5}

1 - Department of Plastic, Reconstructive and Aesthetic Surgery, South Hospital, University of Rennes 1, Rennes, France.

2 - INSERM U917, University of Rennes 1, Rennes, France.

3 - SITI Laboratory, Etablissement Français du Sang Bretagne, Rennes University Hospital, Rennes, France.

4 - STROMA lab, UMR5273 CNRS/UPS/EFS - INSERM U1031, Rangueil Hospital, Toulouse, France.

5 - Department of Plastic, Reconstructive and Aesthetic Surgery, Rangueil Hospital, University Paul Sabatier, Toulouse, France.

Corresponding author:

Dr Nicolas Bertheuil

Department of Plastic, Reconstructive and Aesthetic Surgery, Hospital Sud, Rennes, France.

16 Boulevard de Bulgarie, 35200, Rennes

Phone: 00 33 2 99 26 71 68

Fax: 00 33 2 99 26 67 18

E-mail: nbertheuil@gmail.com

None of the authors has a financial interest in any of the products, devices, or drugs mentioned in this manuscript

Sir,

We read with great interest the manuscript titled “Mechanical Micronization of Lipoaspirates: Squeeze and Emulsification Techniques” by Mashiko *et al* ¹. Foremost, we want to congratulate the authors for this beautiful work that adds significant details to our knowledge on mechanical extraction of stroma vascular fraction (SVF) of adipose tissue without enzymatic digestion. Indeed, the authors demonstrated the possible presence of adipose-derived stromal cells (ASCs) (CD45-/CD31-/CD34+ cells) in squeezed and emulsified fat. The emulsification technique developed by the author’s results in two distinct products: a solid fraction (called REF) and a fluid fraction (called FEF). The analysis of the cell composition is performed essentially to the solid portion which is different from the product called nanofat by Tonnard ². Indeed, Tonnard et al analyzed the liquid phase eg: the SVF obtained after a similar technique without the two steps of centrifugation. The fluid portion is particularly interesting in clinical practice for indications of regenerative surgery (post radiotherapy, burns sequelae...), whereas the solid portion maybe useful for intermediate indications between fat grafting (volumetric correction of defect) and nanofat grafting (regenerative surgery). This new study characterizes a portion not studied until now.

We would like to discuss two points that has not been addressed by the authors.

First, in our opinion, it would be really interesting to determine if the ASCs (CD45-/CD31-/CD34+) isolated by these two free enzymatic extraction methods are “classical” adipose-derived stromal cells. Indeed, ASCs are characterized by different criteria ³ such as: their ability to form colony-forming unit-fibroblasts cells (CFU-Fs), the ability to differentiate into various cell types derived from the mesodermal lineage (bone, cartilage and adipose tissue) and the presence of mesenchymal markers (CD105, CD73, CD90) which were not assayed by the authors

Moreover, we demonstrated that the SVF obtained after emulsification of the fat with the nanofat technique, described by Tonnard contains functional ASCs ⁴. Indeed, ASCs isolated with this technique present all the criteria described below and are functional with immunosuppressive properties characterized by the capability to decrease the proliferation of human T cells. Functional ASCs, through secretion of paracrine factors can modulate inflammation and can favorise angiogenesis, two factors that play an essential role in tissue regeneration ⁵.

In conclusion, all changes in the technique of mechanical extraction of SVF cells can modify cell composition and may alter cell properties. Therefore, SVF cells isolated with new mechanical techniques has to be tested *in vitro* to demonstrate the presence of ASCs cells. Moreover, presence of functional ASCs has to be determined to judge and to validate the use of this new technique in the operating room and to choose the best regenerative mechanical technique.

References:

1. Mashiko T, Wu SH, Feng J, et al. Mechanical Micronization of Lipoaspirates: Squeeze and Emulsification Techniques. *Plast Reconstr Surg*;139(1):79-90.
2. Tonnard PL, Verpaele AM. Discussion: Mechanical Micronization of Lipoaspirates: Squeeze and Emulsification Techniques. *Plast Reconstr Surg*;139(1):91-92.
3. Bourin P, Bunnell BA, Casteilla L, Dominici M, Katz AJ, March KL, Redl H, Rubin JP, Yoshimura K, Gimble JM. Stromal cells from the adipose tissue-derived stromal vascular fraction and culture expanded adipose tissue-derived stromal/stem cells: a joint statement of the International Federation for Adipose Therapeutics and Science (IFATS) and the International Society for Cellular Therapy (ISCT). *Cytotherapy*. 2013 Jun;15(6):641-8.
4. Chaput B, Bertheuil N, Escubes M, Grolleau JL, Garrido I, Laloze J, Espagnol N, Casteilla L, Sensebé L, Varin A. Mechanically Isolated Stromal Vascular Fraction Provides a Valid and Useful Collagenase-Free Alternative Technique: A Comparative Study. *Plast Reconstr Surg*;138(4):807-19.
5. Bertheuil N, Chaput B, Ménard C, Varin A, Garrido I, Grolleau JL, Sensebé L, Watier E, Tarte K. Adipose-derived stromal cells: history, isolation, immunomodulatory properties and clinical perspectives. *Ann Chir Plast Esthet*. 2015 Apr;60(2):94-102.