



**HAL**  
open science

# Spontaneous approaches of divers by free-ranging orcas (Orcinus orca): age- and sex-differences in exploratory behaviours and visual laterality

Stéphanie Chanvallon, Catherine Blois-Heulin, Pierre Robert de Latour,  
Alban Lemasson

## ► To cite this version:

Stéphanie Chanvallon, Catherine Blois-Heulin, Pierre Robert de Latour, Alban Lemasson. Spontaneous approaches of divers by free-ranging orcas (Orcinus orca): age- and sex-differences in exploratory behaviours and visual laterality. *Scientific Reports*, 2017, 7 (1), pp.10922. 10.1038/s41598-017-11488-3. hal-01583555

**HAL Id: hal-01583555**

**<https://univ-rennes.hal.science/hal-01583555>**

Submitted on 7 Sep 2017

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# SCIENTIFIC REPORTS



OPEN

## Spontaneous approaches of divers by free-ranging orcas (*Orcinus orca*): age- and sex-differences in exploratory behaviours and visual laterality

Stéphanie Chanvallon<sup>1</sup>, Catherine Blois-Heulin<sup>1</sup>, Pierre Robert de Latour<sup>2</sup> & Alban Lemasson<sup>1</sup>

Running comparative studies of laterality in mammals is a way to deepen our understanding of the evolution of the brain hemisphere functions. Studies on vision highlighted a possible task-sharing between hemispheres depending on the characteristics of the observers, the nature of the observed stimulus and the context of the observation, a phenomenon that could go beyond the monitoring of conspecifics. Cetaceans are predators that adapted to an aquatic habitat and display a clear crossing of fibers to the side of the brain opposite the eye of origin. Here, we analysed the interactions between humans and cetaceans when free-ranging orcas approach divers. Our study concentrated on the spontaneous exploratory behaviours of divers by orcas depending on their age and sex, and on the possible expression of a visual laterality. The results showed a significant preference for the use of the left eye but exclusively in adult females. Adult males had a more sustained attention than adult females, marked by a higher spatial proximity to divers, slower approaches and longer look durations. Adult females, probably more cautious, explored from the distance and more furtively. Our findings support a possible link between attentional/motivational states and visual laterality in mammals.

Recent studies suggest that brain laterality is much older than we thought until now and certainly not unique to humans<sup>1,2</sup>. We actually find behavioural descriptions of motor and perceptual laterality in nearly all the groups of vertebrates as well as in some invertebrates. Visual laterality has notably been documented in a large number of species (i.e. Fish:<sup>3,4</sup>; Amphibians:<sup>5,6</sup>; Reptiles:<sup>7,8</sup>; Birds:<sup>9,10</sup>; Mammals:<sup>11-15</sup>; Invertebrates:<sup>16,17</sup>), but rarely in large marine mammals because of the difficulty of observing them in the field and their relative rarity in captivity. According to Karenina *et al.*<sup>18</sup>, when individuals need to coordinate their behaviour with others, the uniformity of preferential sides inside a population is advantageous. For many animal species, though a consensus appears around the dominant use of one eye over the other one (and therefore the dominant processing by a given hemisphere), the variations regarding which eye (i.e. left or right) is used preferentially are important and seem to depend on the nature of the observed stimulus (e.g. novelty, valence and complexity<sup>4,19-26</sup>, as well as on the individual characteristics of the observer (e.g. vigilance state and level of stress:<sup>27</sup>; age:<sup>28</sup>) and on the social context of the interaction<sup>29-32</sup>. Laterality at population or group levels is pointed out in different contexts such as foraging for food<sup>23,24</sup>, antipredator<sup>33</sup> or social vigilance<sup>29,34,35</sup>, as well as environmental exploration<sup>13,22</sup>.

Cetaceans, contrasting to what was once believed, because of their often dark natural habitat (i.e. deep water) and developed echolocation<sup>36</sup>, have good vision in the air as underwater<sup>37</sup>. Furthermore, cetaceans have eyes in a semi lateral position with a near total crossing of optical nerves (left eye - right hemisphere)<sup>38</sup>. A few studies actually underline the existence of a possible visual laterality in cetaceans, even though the factors determining the expression of this laterality are still not very well known. For example, in free-ranging belugas (*Delphinapterus leucas*) and orcas (*Orcinus orca*), the juvenile is often positioned on the right side when it swims near its mother, maintaining a visual contact with its left eye<sup>18,39</sup>. This preference for a lateralized swim position was recently

<sup>1</sup>Université de Rennes 1, Ethologie animale et humaine – EthoS, UMR 6552 – CNRS – Université Caen Normandie, Station Biologique, 35380, Paimpont, France. <sup>2</sup>USEA, Orques Sans Frontières, 40280, Bretagne-de-Marsan, France. Correspondence and requests for materials should be addressed to A.L. (email: [alban.lemasson@univ-rennes1.fr](mailto:alban.lemasson@univ-rennes1.fr))

confirmed with a beluga population in managed care<sup>40</sup>. Nevertheless, differences are found between social and non-social monitoring. When hunting, orcas attack their prey by placing them in their right visual field, whereas no lateral bias is found when they jump out of water and fall back on their side<sup>41</sup>. The right hemisphere of cetaceans would thus play a predominant role for social interactions<sup>18, 39, 42</sup>, whereas in response to non-social or non-familiar stimuli, a predominance of the left hemisphere can be found<sup>26</sup>.

In captivity, the left eye was preferentially used by bottlenose dolphins (*Tursiops truncatus*) to observe a human being, familiar or not, approaching the pool<sup>17</sup>. The individual left eye lateralized preferences were later confirmed by studies with captive belugas and pacific white-sided dolphins (*Lagenorhynchus obliquidens*)<sup>43</sup>. In the same experimental conditions, when the human being was replaced by an object, the degree of familiarity with the observed item was important. Indeed, Blois-Heulin *et al.*<sup>22</sup> found that, while non-familiar objects are observed preferentially with the right eye, the left eye is used to look at very familiar (previously manipulated) objects. Both eyes are used indiscriminately for intermediary items, i.e. non-familiar objects which have become to some extent familiar from previous observations but never manipulated. All in all, the mother, the human (assimilated to the one that is providing care and food) and the toys in captivity are preferentially processed by the right hemisphere, whereas prey and new objects are processed by the left hemisphere. Interestingly, the degree of familiarity with the human or object observed also influenced gaze duration in several cetacean species<sup>21, 43–45</sup>. The question thus remains open as to whether or not the observer's attention and motivational state may be a determining factor. More studies are now needed, notably with free-ranging cetaceans, and the cases of human – orca spontaneous encounters is particularly interesting because the former is a potential prey for these animals.

Orcas show numerous and complex capacities of social coordination, which involve both communication and cognition. They indeed show a great diversity of innovated and learned foraging techniques and feeding cooperation strategies<sup>46–48</sup>, individual and matrilineal acoustic signatures with dialectal cultural variations<sup>47–49</sup> and a long period of maternal dependence for the single offspring<sup>50</sup>. Orcas are also curious animals characterized by a highly developed neophilia in captivity<sup>51</sup> and frequent manipulation of various objects in the wild<sup>52</sup>. Recent studies have described cases of play behaviour innovations in resident orcas living in the U.S. west coast<sup>46, 53</sup>.

In several mammal species, innovations, risk-taking and exploratory behaviours vary with the age and sex of the individual (e.g. rats and primates<sup>54–57</sup>). In captive dolphins, youngsters are generally more curious than adults<sup>58</sup>, and 80% of the new behaviours observed in plays come from juveniles<sup>59</sup>. However, Baird *et al.*<sup>60</sup> found no influence of age and sex in diving rates in orcas, but male adults would dive more often during the day than female adults. The question of exploratory behaviour regarding age, sex, as far as we know, has never been studied in free-ranging orcas. Previous studies in cetaceans (e.g. dolphins and belugas) have shown that gaze duration (and possibly swimming speed and distance of approach) plays a key role during exploration of familiar and unfamiliar humans and objects<sup>21, 43–45</sup>. Although the exploratory behaviours of cetaceans have still been poorly described in the literature, particular behaviours like head and body movements, some being associated with vocalisations (notably echolocation), have also been previously described in this context<sup>61, 62</sup>.


How humans are perceived by free-ranging orcas is a current hot topic, notably because whale-watching and swimming-with-whales ecotouristic sites are multiplying. To this day, little research has been conducted scientifically about the nature of human – orca spontaneous encounters in the wild and in the diversity of associated behaviours. Orcas' interest is expressed by behaviours like following boats and many spontaneous interactions with humans (cetologists, fishermen, tourists), including play and even actions of supposed cooperation with humans in their fishing activities<sup>63</sup>. In the wild, orcas are totally free of their movements and voluntarily approach divers. By using underwater videos, made in Norway, we therefore concentrated on gazes and other behaviours in the context of spontaneous approaches of divers by orcas. We studied an area where diving is organised every year. We have analysed the influence of individual's age and sex on the behaviours displayed during these approaches and tested a possible eye-preference during visual monitoring.

## Results

Among all approaches sampled, 24%, 61.3% and 14.7% consisted of adult males, adult females and subadults respectively. Monocular gazes were clearly the majority (82.5%) of all gazes sampled. Nevertheless, the eye preferentially chosen by the different individuals varied with their age and sex. Thus, while adult males (Wilcoxon matched-paired test,  $N = 18$ ,  $Q_{obs} = 40$ ,  $p = 0.97$ ) and subadults ( $N = 11$ ,  $Q_{obs} = 7.5$ ,  $p = 1$ ) did not preferentially use one eye over the other, adult females used their left eye more frequently ( $N = 60$ ,  $Q_{obs} = 191$ ,  $p = 0.04$ ) (Fig. 1).

In most cases (86%), the “approach consisted in a progressive movement towards the divers and more rarely (14%) in a “passing by” movement (i.e. quasi-linear trajectory, constant distance with the divers). Moreover, orcas approached preferentially from the front or below (98%) and in very rare cases from the back (2%). After having excluded the rare approaches from behind, we found that all units were relatively frequent (i.e. from the right side 39%, from the left side 29%, facing 14%, just under 10%, from deep water 8%).

However, certain types of approach were more frequently observed than others in the three age-sex classes of individuals (see all statistical intra-class comparisons in Table 1). First, concerning the social context and the minimum distance of approach, we found that all individuals approached significantly more often alone than accompanied, and more often close than far. However, while adult females approached more often close than very close, no difference was found for adult males and subadults. Second, concerning visual monitoring and swimming speed, while adult females and subadults displayed more short than long gazes, no difference was found for adult males. This has to be related to some extent to the swimming speed that was more frequently slow than fast in adult males but equally slow and fast in both adult females and subadults. Third, concerning head movements, while adult males displayed more horizontal than vertical movements, both types were equally observed in adult females and subadults. Fourth, all individuals approached significantly more often silently than accompanied by whistle calls.



**Figure 1.** Preferential use of the right (grey bars) or left (black bars) eye by orcas according to their sex and age. Wilcoxon test: \* $p < 0.05$ .

Comparisons	Adult females	Adult males	Subadults
Alone versus with another orca(s)	3.05 ± 0.06/1.16 ± 0.02	4.61 ± 0.65/0.88 ± 0.37	4.72 ± 2.04/0.72 ± 0.54
	N = 56 z = 3.32	N = 17 z = 5.51	N = 11 z = 2.93
	<b>p = 0.0009</b>	<b>p = 0.0007</b>	<b>p = 0.003</b>
Close versus Far	3.86 ± 0.07/0.3 ± 0	5.11 ± 0.83/0.2 ± 0.1	5.45 ± 2.67/0.1 ± 0.09
	N = 56 z = 6.23	N = 18 z = 3.62	N = 11 z = 2.93
	<b>p = 0.00001</b>	<b>p = 0.0003</b>	<b>p = 0.003</b>
Very close versus Close	0.96 ± 0.03/2.9 ± 0	3.11 ± 0.57/2.05 ± 0.59	2.91 ± 2.25/2.54 ± 0.55
	N = 54 z = 4.89	N = 15 z = 1.47	N = 10 z = 1.68
	<b>p &lt; 0.0001</b>	<b>p = 0.14</b>	<b>p = 0.09</b>
Short versus Long gaze	2.71 ± 0.04/1.46 ± 0.04	2.16 ± 0.61/3.33 ± 0.56	3.72 ± 1.84/1.63 ± 0.86
	N = 40 z = 3.91	N = 16 z = 1.75	N = 9 z = 2.07
	<b>p = 0.00009</b>	<b>p = 0.08</b>	<b>p = 0.04</b>
Slow versus Fast swimming	4.25 ± 0.07/1.87 ± 0.03	4.22 ± 0.61/1.11 ± 0.43	4.45 ± 1.34/4.49 ± 1.35
	N = 52 z = 1.48	N = 17 z = 3.40	N = 10 z = 0.96
	<b>p = 0.14</b>	<b>p = 0.0006</b>	<b>p = 0.33</b>
Horizontal versus Vertical head movement	0.36 ± 0.01/0.25 ± 0.01	0.5 ± 0.18/0.22 ± 0.10	0.36 ± 0.36/0.45 ± 0.36
	N = 16 z = 0.98	N = 6 z = 2.20	N = 1
	<b>p = 0.33</b>	<b>p = 0.03</b>	
Whistling versus Silence	1.05 ± 0.27/3.13 ± 0.50	1.27 ± 0.40/4.22 ± 0.65	1.83 ± 1.05/8.33 ± 4.41
	N = 55 z = 4.47	N = 16 z = 2	N = 11 z = 3.059
	<b>p = 0.00008</b>	<b>p = 0.0006</b>	<b>p = 0.003</b>


**Table 1.** Approach characteristics of the different age-sex classes. In each cell: First line – mean number of occurrences (+/– standard error) for the two compared behavioural units; Second and third lines – Number of individuals taken into account in the statistical analysis (equal scores excluded), z score and p value of the Wilcoxon test (significances in bold). NA: test not applicable.

Finally, two behaviours were found only in adult females and subadults, and not in adult males: belly presentation (adult females 6.7%, subadults 18% of times), and body rotation (adult females 16.1%, subadults 16.4%).


The above differences were confirmed for some of the measured variables when running comparisons between adult males, adult females and subadults (inter-classes comparisons). Indeed, adult males approached significantly the diver at a shorter distance than adult females (Mann-Whitney test,  $n_1 = 18$   $n_2 = 60$ ,  $z = -3.11$ ,  $p = 0.002$ ), but only a trend was found with subadults ( $n_1 = 18$   $n_2 = 11$ ,  $z = 1.77$ ,  $p = 0.08$ ) (Fig. 2). Also, adult females and subadults did not differ in the proportion of times they approached at a very close distance ( $z = 0.01$ ,  $p = 0.99$ ). Moreover, adult males displayed significantly fewer short gazes than adult females ( $z = 4.05$ ,  $p = 0.0001$ ) and than subadults ( $z = -2.89$ ;  $p = 0.004$ ), whereas adult males and subadults did not differ ( $z = -0.72$ ;  $p = 0.47$ ) (Fig. 3).

The swimming speed was also significantly slower in adult males than in subadults ( $z = 2.29$ ,  $p = 0.007$ ), while adult females' speed did not differ (when applying Bonferroni correction) with those of adult males ( $z = -2.001$ ,  $p = 0.045$ ) and subadults ( $z = 1.35$ ,  $p = 0.18$ ) (Fig. 4).


No influence of age and sex was found for the proportion of solitary approaches (Kruskal-Wallis test:  $H(2.89) = 2.69$ ,  $p = 0.26$ ; males  $88.56 \pm 4.10\%$ , females  $69.91 \pm 5.26\%$  and subadults  $93.39 \pm 3.57\%$ ), horizontal head movements ( $H(2.89) = 3.68$   $p = 0.16$ ; males  $34.72 \pm 10.91\%$ , females  $20.97 \pm 4.98\%$  and subadults



**Figure 2.** Variation of the individual percentage of approaches (mean  $\pm$  s.e.) from a very close distance with the age and sex of the orca. Mann-Whitney test: \*\* $p = 0.002$ .



**Figure 3.** Variation of the individual percentage of short gazes (mean  $\pm$  s.e.) with the age and sex of the orca. Mann-Whitney test: \*\*\* $p = 0.0001$ , \*\* $p = 0.004$ .



**Figure 4.** Variation of the individual percentage of slow swimming (mean  $\pm$  s.e.) with the age and sex of the orca. Mann-Whitney test: \*\* $p = 0.007$ .

4.55  $\pm$  4.55%), whistling approaches ( $H(2.89) = 0.74$ ,  $p = 0.69$ ; males 34.72  $\pm$  5.25%, females 22.51  $\pm$  4.23% and subadults 12.14  $\pm$  4.37%).

In order to better understand the context of whistling, we tested whether certain variables were more often associated than others. Indeed, we found that while there were as many approaches with whistles when the orca approached alone or not (Wilcoxon matched-paired test:  $N = 83$ ,  $Q = 1806$ ,  $p = 0.743$ ), there were significantly more approaches with whistles when the swimming speed was slow than fast ( $N = 83$ ,  $Q = 2478$ ,  $p = 0.0001$ ).

## Discussion

This first investigation of the nature of the exploratory behaviour of free-ranging orcas when approaching human divers shows two important findings. First, adult females prefer to visually monitor humans with their left eye than with their right eye, but no eye preference was found in adult males and subadults. Second, the approach behaviours differed significantly between different individuals; adult males demonstrated stronger attention, as well as potentially greater curiosity (shorter distance, slower swimming and longer gazes) than adult females and subadults, the latter being possibly more cautious.

The visual laterality in favour of the left eye is in accordance with the previous findings on social interactions in wild and captive cetaceans<sup>18, 39</sup> and with captive dolphins when looking at humans<sup>21</sup> and at familiar toys<sup>22</sup>. It



**Figure 5.** Photos of an adult male (above) and an adult female (below) orcas approaching divers with the respective use of a right and a left eye for visual monitoring (acknowledgments to Pierre Robert de Latour for providing the pictures).

therefore confirms the dominance of the right hemisphere in the visual exploration of some salient and familiar items by cetaceans (see also ref. 64) and allows us to support the hypothesis that the human diver is a source of interest for the free-ranging orcas habituated to their presence (for about 18 years). This hypothesis is also supported by the fact that no aggressiveness or sudden withdrawal has been noticed.

However, not all individuals preferred a left-side gazing in our study. Several non-mutually exclusive hypotheses may explain the individual variations observed. First, it is possible that laterality is more pronounced in older individuals than younger ones, due to their experiences with those unusual encounters, which would account for the absence of laterality in subadults. Indeed, the strength of laterality in animals is frequently related to the degree of familiarity with the observed item<sup>22</sup>. This is also supported by the fact that more adult females are in contact with divers than adult males in this area, and thus more experienced with human encounters. Over two consecutive years, the number of approaches by adult females was 4.5 times higher than the number of approaches by adult males. Second, it is possible that the emotional state of male and female adults differed in our study, as this is another factor of variation in humans and animals for both strength and direction of laterality<sup>65–67</sup>. The fact that females generally keep a greater distance than males here and only observe humans briefly could be the sign of cautiousness in females that, while approaching, would remain more vigilant. Third, it is very likely that there is a link between the direction of the laterality and the degree of attention of the observer, as it was found earlier in birds<sup>68</sup>. Males, being more attentive, perhaps integrate more details, thus mobilizing their two hemispheres. The left hemisphere is known to play a key role for detailed visual exploration in various animal species<sup>69–73</sup>. Fourth, the hormonal difference between males and females may also be responsible for the sex-dependent laterality observed here. A study on testosterone-treated chicks found that an effect of testosterone was greater reliance on the frontal visual field<sup>74</sup>, which could be a possible explanation for the absence of lateral bias due to a stronger collaboration between the two hemispheres in male orcas. Fifth, differences in cognitive abilities between males and females could explain the sex-dependent laterality found in our study. In humans, females perform better than males in various memory tasks<sup>75,76</sup>. This female superiority would be explained by a more effective use of the abilities of the right hemisphere for accessing memory of patterns made up of multiple items<sup>2</sup>.

Also in our study, adult males approached less often than adult females; this is probably due simply to their smaller number in the overall population. The fact that males approached divers at a closer distance and for longer exploration sequences is an important point of discussion. One abovementioned explanation is that males may be more curious. Unfortunately, to our knowledge, there is no such “personality type” study done with cetaceans that could confirm this hypothesis. However, sex-specific responses to human presence have been found earlier in cetaceans. Williams *et al.*<sup>77</sup> studied the habituation to human activities by orcas in Johnson Detroits (British Columbia). They found that the animal’s reaction to boats changed in a sex-specific way with time. While females tended to accelerate their swimming speed and increase the angle of their diving routes, males kept their swimming speed constant with less predictable trajectory. This said, the fact that females approached less closely in our study does not mean that they did not “observe”; cetaceans may use echolocation to complete their

Measured variables	Behavioural unit definitions
<b>Distance</b> (minimum distance between the orca and the diving guide reached during a given approach)	The orca approaches the diver from:
	- Very close (less than 5 meters)
	- Close (between 5 and 20 meters)
	- Far (more than 20 meters).
<b>Angle of approach</b>	The orca approaches the diver from:
	- The front:
	* from the right side of the diver (angle: 30 to 120°)
	* from the left side of the diver (angle: 200 to 330°)
	* facing the diver (angle: 0°)
	- Below:
	* moving just under the diver
* arriving vertically from deep water	
- The back.	
<b>Swimming speed</b>	- Slow: the orca moves but the movements of its caudal fin are hardly detectable
	- Fast: at least one caudal fin movement per second.
<b>Gaze laterality</b>	The orca looks at the diver with:
	- its right eye
	- its left eye
	- both eyes (bilateral)
	<i>For a given approach, the eye used may change during the behavioural sequence and each gaze change is counted for the laterality analysis.</i>
<b>Gaze duration</b> (mono- and binocular pooled)	The orca looks at the diver for:
	- less than 6 seconds (Short gaze)
	- more than 6 seconds (Long gaze)
	<i>For a given approach, we measured the total gazing duration.</i>
<b>Head movements</b>	- Horizontal (the orca turns slightly its head towards the diver)
	- Vertical (the orca is facing the diver and moves the head up and down several times in a row).
<b>Whistle calls</b>	- Whistling (a high-pitched modulated frequency is heard)
	- Silence (no species-specific sound is heard)
	<i>It is not possible to confirm whether the caller is the observed orca or a neighbouring one.</i>
<b>Belly presentation</b>	The orca moves the head up presenting its belly to the humans.
<b>Body rotation</b>	The orca rotates its body along the horizontal axis.

**Table 2.** Characteristics of the different types of approach and associated behaviours.

visual exploration<sup>36,78</sup>. It is possible that females prioritized long-distance echolocation over short-distance eye observation. In line with this, vertical head movements, typically associated with echolocation<sup>61</sup>, were particularly frequent in adult females. Scheer *et al.*<sup>61</sup> also described this echolocation behaviour directed to humans at a 20-meter distance in free-ranging pilot whales (*Globicephala macrorhynchus*). As Blois-Heulin *et al.*<sup>22</sup>, we suppose that echolocation, associated to a binocular vision, completes the amount of information collected on objects, facilitating the construction of its representation.

The approaches described here are signs of a certain curiosity, and possibly attempts to initiate interactions with divers. The fact that the vast majority of the approaches come from the front clearly indicates that animals behave to remain visible. Scheer *et al.*<sup>61</sup> with free-ranging pilot whales and Duroyon<sup>62</sup> with captive bottlenose dolphins (*Tursiops truncatus*) have also described behaviours like belly presentation, body rotation, horizontal head movements that were clearly associated with non-agonistic encounters and interaction initiations. To our knowledge, the function of whistling in this context has never been studied, but it is intriguing to find more calls associated with closer and longer approaches. This also shows that orcas do not seek to be silent, whereas they do when hunting mammals<sup>79–81</sup>. We must acknowledge however that we could not identify callers and more solid conclusions require further investigation.

In conclusion, the fact that visual laterality and exploratory behaviours are sex- and age-dependent has interesting perspectives in both basic (understanding of the underlying, notably psychological, mechanisms and evolutionary processes) and applied research fields. Indeed, human-cetacean encounters are increasing all around the world<sup>82,83</sup> and the development of ecotourism over the last 20 years has created a significant pressure on the environment of cetaceans, and this is not limited to cetaceans. Complementary studies are necessary to explore further the different hypotheses raised here, and notably to test to which extent the motivation, the temperament, the degree of attention and the level of experience of animals are determining factors of behavioural variations.

## Materials and Methods

**Study animals, context of the study and video recordings.** The presence of orcas in the Kalkfjord (Norway) is directly linked to the massive presence of specific prey, i.e. atlantic herrings (*Clupea harengus*). As a result, the same population of orcas comes back to this location every year<sup>84</sup>. Individuals are identified based on classic criterion for that species, such as the size and the shape of the dorsal fin and the saddle patch as well as bodies scars<sup>85</sup>. Only individuals clearly identified and observed at least twice during the study have been included in the data set, i.e. 60 adult females, 18 adult males, 11 unsexed subadults.

The underwater videos used for the analyses have been made by Pierre Robert de Latour during ecotouristic dives organised in Norway on winters 2014 and 2015 (November to February). The expeditions have been set up from the base M/S SULA and the dives started from a small barge, within a narrow period of daylight (i.e. 10 am–2 pm). Orcas are first observed at a distance, and then the barge is used to approach the animals following the USEA (Undersea Soft Encounter Alliance) method developed by Pierre Robert de Latour. This method is based on the principle of a limited intrusion into the orcas' space, who can then decide to approach the barge and the divers further or not. The divers (including snorkelers), with a maximum of eight people simultaneously in the water, are gathered just behind the guide, Pierre Robert de Latour. All people are commanded to stay as a group, move slowly and silently.

Video recordings have been made with an underwater camera Sony Ilce A6000 with a Nauticam NA 6000 housing, and equipped with a 24 mm focal equivalent to 24\*36 (shutter speed: 1/25 to 1/100, aperture: f2.8, ISO: automatic limited at 3200, White Balance: 4800k). All in all, 902 video sequences (collected with an *ad libitum* sampling method, i.e. at various times of the diving session and with various durations) have been analysed, that is 540 minutes of film sampled (2014: 323 video sequences lasting 163 minutes in total; 2015: 579 video sequences lasting 377 minutes in total).

Our study is observational and totally non-invasive, based on video scoring. Videos come from the archive of the USEA – Orques Sans Frontieres program (permit #USEA2017-0007).

**Behavioural repertoire and data collection.** A detailed behavioural repertoire has been established to sample the different movements and other exploratory behaviours of orcas when approaching the divers (see photo examples in Fig. 5). Here, an approach is defined as follows<sup>1</sup>: an approach is systematically an animal movement oriented in the direction of the divers<sup>2</sup>; an approach starts from the moment where an orca enters the camera range and ends as soon as it gets out. The behavioural repertoire, composed of 10 measured variables (most of them being divided into mutually exclusive units), is given in Table 2. For each single approach, we used a 1-0 sampling method to score all measured variables.

**Statistical analyses.** In a first step, to study the possible existence of a visual laterality bias during approaches at the population level, we compared, using Wilcoxon tests for matched samples, the number of gazes from the left eye with the number of gazes from the right eye directed to the humans based on all individuals sampled in each age-sex class.

In a second step, we tested whether the different individuals differed in the way they approached humans. On the one hand, using Wilcoxon tests for matched samples, we compared, for each measured variable and each age-sex class of individuals, the number of occurrences of the different behavioural units sampled. On the other hand, we calculated for each individual sampled the percentage of expression of a given unit over all the behavioural units of each measured variable. These individual percentages were then used to compare the behavioural profiles of adult males, adult females and unsexed subadults using Kruskal-Wallis and post-hoc Mann - Whitney tests (applying Bonferroni corrections for multiple comparisons). Significant threshold was set at 0.05 (and 0.0167 in case of Bonferroni correction).

## References

- Rogers, L. J. & Andrew, R. *Comparative vertebrate lateralization* (Cambridge University Press, 2002).
- Rogers, L. J., Vallortigara, G. & Andrew, R. J. *Divided brains: the biology and behaviour of brain asymmetries* (Cambridge University Press, 2013).
- Sovrano, V. A., Bisazza, A. & Vallortigara, G. Lateralization of response to social stimuli in fishes: A comparison between different methods and species. *Physiol. Behav.* **74**, 237–244 (2001).
- Lippolis, G., Joss, J. M. P. & Rogers, L. J. Australian Lungfish (*Neoceratodus forsteri*): a missing link in the evolution of complementary side biases for predator avoidance and prey capture. *Brain Behav. Evol.* **73**, 295–303 (2009).
- Robins, A. & Rogers, L. J. Lateralized prey catching responses in the toad (*Bufo marinus*): analysis of complex visual stimuli. *Anim. Behav.* **68**, 567–575 (2004).
- Lippolis, G., Bisazza, A., Rogers, L. J. & Vallortigara, G. Lateralization of predator avoidance responses in three species of toads. *Laterality* **7**, 163–183 (2002).
- Bonati, B. & Csermely, D. Lateralization in Lizards: Evidence of presence in *several contexts in Behavioral Lateralization in Vertebrates* (eds Csermely, D. & Regolin, L.) 35–38 (Springer, Heidelberg, 2011).
- Deckel, A. W. & Fuqua, L. Effects of serotonergic drugs on lateralized aggression and aggressive displays in *Anolis carolinensis*. *Behav. Brain. Res.* **95**, 227–232 (1998).
- Franklin, W. E. & Lima, S. L. Laterality in avian vigilance; do sparrow have a favourite eye? *Anim. Behav.* **62**, 879–885 (2001).
- Ventolini, N. *et al.* Laterality in the wild: preferential hemifield use during predatory and sexual behaviour in the black winged stilt (*Himantopus himantopus*). *Anim. Behav.* **69**, 1077–1084 (2005).
- Chapelain, A. S. & Blois-Heulin, C. Eye preference in Campbell's monkeys (*Cercopithecus c. campbelli*). *Anim. Cogn.* **12**, 11–19 (2009).
- Hook-Costigan, M. A. & Rogers, L. J. Eye preference in common marmosets (*Callithrix jacchus*): influence of age, stimulus and hand preferences. *Laterality* **3**, 180–191 (1998).
- Bracchini, S. N., Lambeth, S. P., Schapiro, S. J. & Tecumseh Fitch, W. Eye preferences in captive chimpanzees. *Anim. Cogn.* **15**, 971–978 (2012).

14. Des Roches, A. D. B., Richard-Yris, M. A., Henry, S., Ezzaoui, M. & Hausberger, M. Laterality and emotions: visual laterality in the domestic horse (*Equus caballus*) differs with objects' emotional value. *Physiol. Behav.* **94**, 487–490 (2008).
15. Siniscalchi, M., Sasso, R., Pepe, A. M., Vallortigara, G. & Quaranta, A. Dogs turn left to emotional stimuli. *Behav. Brain Res.* **208**, 516–521 (2010).
16. Frasnelli, E., Vallortigara, G. & Rogers, L. J. Response competition associated with right-left antennal asymmetries of new and old olfactory memory traces in honeybees. *Behav. Brain Res.* **209**, 36–41 (2010).
17. Jozet-Alves, C. *et al.* Visual lateralization is task- and age-dependent in cuttlefish (*Sepia officinalis*). *Anim. Behav.* **83**, 1313–1318 (2012).
18. Karenina, K., Giljov, A., Ivkovich, T., Burdin, A. & Malashichev, Y. Lateralization of spatial relationships between wild mother and infant orcas (*Orcinus orca*). *Anim. Behav.* **86**, 1225–1231 (2013).
19. Cantalupo, C., Bisazza, A. & Vallortigara, G. Lateralization of predator evasion response in a teleost fish (*Girardinus falcatus*: Poeciliidae). *Neuropsychologia* **33**, 1637–1646 (1995).
20. Quaranta, A., Siniscalchi, M. & Vallortigara, G. Asymmetric tail wagging responses by dogs to different emotive stimuli. *Curr. Biol.* **17**, 199–201 (2007).
21. Thielges, H., Lemasson, A., Kuczaj, S., Böye, M. & Blois-Heulin, C. Visual laterality in dolphins when looking at (un)familiar humans. *Anim. Cogn.* **14**, 303–308 (2011).
22. Blois-Heulin, C., Crevel, M., Lemasson, A. & Böye, M. Visual laterality in dolphins: importance of the familiarity of stimuli. *BMC Neuroscience* **13**, 9 (2012).
23. Giljov, A. N., Karenina, K. A. & Malashichev, Y. B. An eye for a worm: lateralisation of feeding behaviour in aquatic anamniotes. *Laterality* **14**(3), 273–286 (2009).
24. Rogers, L. J., Zucca, P. & Vallortigara, G. Advantage of having a lateralized brain. *Proc. R. Soc. Lond. B. Biol. Sci.* **271**, S420–S422 (2004).
25. Rogers, L. J. & Kaplan, G. An eye for a predator: lateralization in birds, with particular reference to the Australian magpie in *Behavioral and morphological asymmetries in vertebrates* (eds Malashichev, Y. & Deckel, W.) 47–57 (Austin, Landes Bioscience, 2006).
26. Siniscalchi, M., Dimatteo, S., Pepe, A. M., Sasso, R. & Quaranta, A. Visual lateralisation in wild striped dolphins (*Stenella coeruleoalba*) in response to stimuli with different degrees of familiarity. *PLoS One* **7**(1), e3001 (2012).
27. Hook-Costigan, M. A. & Rogers, L. J. Eye preferences in common marmosets (*Callithrix jacchus*): influence of age, stimulus and hand preferences. *Laterality* **3**, 109–130 (1998).
28. Kruper, D. C., Boyle, B. E. & Patton, R. A. Eyes and hand preferences in rhesus monkeys (*Macaca mulatta*). *Psycho. Sciences* **5**, 277–278 (1966).
29. Baraud, I., Buytet, B., Bec, P. & Blois-Heulin, C. Social laterality and 'transversality' in two species of mangabeys: influence of rank and implication for hemispheric specialization. *Behav. Brain Res.* **198**, 449–458 (2009).
30. Casperd, J. M. & Rim, D. Asymmetries in the visual processing of emotional cues during agonistic interactions by gelada baboons. *Behav. Process.* **37**, 57–65 (1996).
31. Hopkins, W. D. & Bennett, A. J. Handedness and approach-avoiding behaviour in chimpanzees. *J. Exp. Psychol. Anim. Behav. Process.* **20**, 413–418 (1994).
32. Quaresmini, C., Forrester, G. S., Spiezio, C. & Vallortigara, G. Social environment elicits lateralized behaviors in gorillas and chimpanzees. *J. Comp. Psychol.* **128**, 276–284 (2014).
33. Rogers, L. J. Evolution of hemispheric specialisation: advantages and disadvantages. *Brain Lang.* **73**, 236–253 (2000).
34. Casperd, L. M. & Dunbar, R. I. M. Asymmetries in the visual processing of emotional cues during agonistic interactions by gelada baboons. *Behav. Processes* **37**, 57–65 (1996).
35. Austin, N. P. & Rogers, L. J. Lateralization of agonistic and vigilance responses in Przewalski horses (*Equus przewalskii*). *Appl. Anim. Behav. Sci.* **151**, 43–50 (2014).
36. Pack, A. A., Herman, L. M., Hoffmann-Khunt, M. & Branstetter, B. K. The object behind the echo: dolphins (*Tursiops truncatus*) perceive object shape globally through echolocation. *Behav. Process.* **58**, 1–26 (2002).
37. Dawson, W. W. The cetacean eye in *Cetacean Behavior* (eds Herman, L. M.) 53–100 (Wiley, New York, 1980).
38. Marino, L. Evolution of the brain and cognition in cetaceans in *Evolutionary cognitive neuroscience* (eds Platek, S. M., Keenan, J. P. & Shackelford, T. K.) 163–193 (Cambridge MIT Press, 2006).
39. Karenina, K., Giljov, A., Glazov, D. & Malashichev, Y. Social laterality in wild beluga whale infants: comparisons between locations, escort conditions, and ages. *Behav. Ecol. Sociobiol.* **67**, 1195–1204 (2013).
40. Hill, H. M. *et al.* Lateralized swim positions are conserved across environments for beluga (*Delphinapterus leucas*) mother-calf pairs. *Behav. Processes* **138**, 22–28 (2017).
41. Karenina, K., Giljov, A., Ivkovich, T., Burdin, A. & Malashichev, Y. Evidence for the perceptual origin of right-sided feeding biases in cetaceans. *Anim. Cogn.* **19**(1), 239–243 (2015).
42. Sakai, M., Hishii, T., Takeda, S. & Kohshima, S. Laterality of flipper rubbing and behaviour in wild bottlenose dolphins (*Tursiops aduncus*): caused by asymmetry of eye use? *Behav. Brain Res.* **170**(2), 204–210 (2006).
43. Yeater, D. B., Hill, H. M., Baus, N., Farnell, H. & Kuczaj, S. A. Visual laterality in belugas (*Delphinapterus leucas*) and Pacific white-sided dolphins (*Lagenorhynchus obliquidens*) when viewing familiar and unfamiliar humans. *Anim. Cogn.* **17**(6), 1245–1259 (2014).
44. Guarino, S., Yeater, D., Lacy, S., Dees, T. & Hill, H. M. Responses to familiar and unfamiliar objects by belugas (*Delphinapterus leucas*), bottlenose dolphins (*Tursiops truncatus*), and Pacific white-sided dolphins (*Lagenorhynchus obliquidens*). *Anim. Cogn.* (2017).
45. Hill, H. M. *et al.* Responses to familiar and unfamiliar humans by belugas (*Delphinapterus leucas*), bottlenose dolphins (*Tursiops truncatus*), & Pacific white-sided dolphins (*Lagenorhynchus obliquidens*): a replication and extension. *Int. J. Comp. Psychol.* **29**, 1–21 (2016).
46. Baird, R. W. Killer Whales of the world: Natural history and conservation. Stillwater (MN: Voyageur Press, 2002).
47. Rendell, L. & Whitehead, H. Culture in whales and dolphins. *Behav. Brain Sci.* **24**, 309–324 (2001).
48. Yurk, H., Barrett-Lennard, L. G., Ford, J. K. B. & Matkin, C. Cultural transmission within matrilineal lineages: Vocal clans in resident killer whales in southern Alaska. *Anim. Behav.* **63**, 1103–1119 (2002).
49. Ford, J. K. B. Acoustic behavior of resident killer whales (*Orcinus orca*) off Vancouver Island, British Columbia. *Canadian Journal of Zoology* **67**, 727–745 (1989).
50. Whitehead, H. & Mann, J. Female reproductive strategies of cetaceans in *Cetacean societies, field studies of dolphins and whales* (eds Mann, J., Connor, R. C., Tyack, P. L. & Whitehead, H.) 219–246 (Chicago, The University of Chicago Press, 2000).
51. Nakahra, F. & Takemura, A. A survey on the behaviour of captive odontocetes in Japan. *Aquatic Mammals* **23**(3), 135–143 (1997).
52. Paulos, R., Trone, M. & Kuczaj, S. A. Play in wild in captive cetaceans. *Int. J. Comp. Psychology* **23**, 701–722 (2010).
53. Whitehead, H., Rendell, L. E., Osborne, R. W. & Würsing, B. Culture and conservation of non-humans with reference to whales and dolphins Review and new directions. *Biol. Conserv.* **120**, 427–437 (2004).
54. Lynn, D. A. & Brown, G. R. The Ontogeny of Exploratory Behavior in Male and Female Adolescent Rats (*Rattus norvegicus*). *Dev Psychobiol.* **51**(6), 513–520 (2009).
55. Reader, S. M. & Laland, K. N. Primate Innovation: sex, age and social rank differences. *Int. J. Primatol.* **22**(5) (2001).
56. Fairbanks, L. A. Risk-Taking by Juvenile Vervet Monkeys. *Behaviour* **124**(1/2), 57–72 (1993).
57. Ouattara, K., Lemasson, A. & Zuberbühler, K. Anti-predator strategies of free-ranging Campbell's monkeys. *Behaviour* **146**, 1687–1708 (2009).

58. Kuczaj, S. A. & Walker, R. T. How do dolphins solve problems? In *Comparative Cognition: Experimental Explorations of Animal Intelligence* (eds Wasserman, E.A. & Zentall, T.R.) 580–601 (New-York: Oxford University, 2006).
59. Kuczaj, S. A., Makecha, R., Trone, M., Paulos, R. D. & Ramos, J. A. Role of peers in cultural innovation and cultural transmission: Evidence from the play of dolphin calves. *International J. Comp. Psychol.* **19**, 223–240 (2006).
60. Baird, R. W., Bradley Hanson, M. & Lawrence, M. D. Factors influencing the diving behaviour of fish-eating killer whales: sex differences and diel and interannual variation in diving rates. *Can. J. Zool.* **83**, 257–267 (2005).
61. Scheer, M., Hofmann, B. & Behr, I.P. Ethogram of selected behaviors initiated by free-ranging short-finned pilot whales (*Globicephala macrorhynchus*) and directed to human swimmers during open water encounters. *Anthrozoös* **17**(3) (2004).
62. Duroyon, C. Problématique de la constitution d'un répertoire comportemental chez une espèce de mammifères marins, le dauphin souffleur (*Tursiops truncatus*). Thèse vétérinaire. Ecole Nationale Vétérinaire d'Alfort (2009).
63. Patterson, E. M. & Mann, J. Cetacean Innovation in *Animal creativity and innovation* (eds. Kaufman, A.B. & Kaufman, J.) 73–120 (Academic Press, 2015).
64. Siniscalchi, M., Dimatteo, S., Pepe, A. M., Sasso, R. & Quaranta, A. Visual lateralization in wild striped dolphins (*Stenella coeruleoalba*) in response to stimuli with different degrees of familiarity. *PLoS ONE* **7**(1), e30001 (2012).
65. Erhan, H., Borod, J. C., Tenke, C. E. & Bruder, G. E. Identification of emotion in a dichotic listening task: event-related brain potential and behavioral findings. *Brain Cogn.* **37**, 286–307 (1998).
66. Borod, J. C. Interhemispheric and intrahemispheric control of emotion: a focus on unilateral brain damage. *J. Consult. Clinic. Psychol.* **60**, 339–348 (1992).
67. Andrew, R. J. & Rogers, L. J. The nature of lateralization in tetrapods in *Comparative vertebrate lateralization* (eds Rogers, L. J. & Andrew, R. J.) 94–125 (Cambridge University Press, 2002).
68. George, I., Vernier, B., Richard, J.-P., Hausberger, M. & Cousillas, H. Hemispheric Specialization in the Primary Auditory Area of Awake and Anesthetized Starlings (*Sturnus vulgaris*). *Behavioral Neuroscience* **118**(3), 597–610 (2004).
69. Fagard, J. & Streri, J. Développement des asymétries chez le bébé, in *Droitiers et gauchers, des asymétries dans tous les sens* (ed. Fagard, J.) 49–70 (Solal Press, 2006).
70. MacNeilage, P. F., Rogers, L. J. & Vallortigana, G. Origins of the left and right brain. *Scientific American* **301**, 60–67 (2009).
71. Tommasi, L. & Vallortigana, G. Encoding of geometric and landmark information in the left and right hemispheres of the avian brain. *Behav. Neurosci.* **115**, 602–613 (2001).
72. Tommasi, L. & Vallortigana, G. Hemispheric processing of landmark and geometric information in male and female domestic chicks (*Gallus gallus*). *Behav. Brain Res.* **155**, 85–96 (2004).
73. Vallortigana, G. & Rogers, L. J. Survival with an asymmetrical brain: advantages and disadvantages of cerebral lateralization. *Behav. Process.* **33**, 41–57 (2005).
74. Rogers, L. J. & Andrew, R. J. Frontal and lateral visual field use after treatment with testosterone. *Anim. Behav.* **38**, 394–405 (1989).
75. James, T. W. & Kimura, D. Sex differences in remembering the locations of objects in an array: location-shift versus location-exchanges. *Evol. Hum. Behav.* **18**, 155–163 (1997).
76. Herlitz, A., Nilsson, L.-G. & Backman, L. Gender differences in episodic memory. *Mem. Cognit.* **25**, 801–811 (1997).
77. Williams, R., Trites, A. W. & Bain, D. E. Behavioral responses of killer whales (*Orcinus orca*) to whale-watching boats: opportunistic observations and experimental approaches. *J. Zool. Lond.* **256**, 255–270 (2002).
78. Barrett-Lennard, L. G., Ford, J. K. B. & Heise, K. A. The mixed blessing of echolocation: differences in sonar use by fish-eating and mammal-eating killer whales. *Anim. Behav.* **51**, 553–565 (1996).
79. Morton, A. B. A quantitative comparison of the behaviour of resident and transient forms of the killer whale off the central British Columbia coast. *Report of the International Whaling Commission* (special issue **12**), 245–248 (1990).
80. Simon, M., McGregor, P. K. & Ugarte, F. The relationship between the acoustic behaviour and surface activity of killer whales (*Orcinus orca*) that feed on herring (*Clupea harengus*). *Acta Ethol.* **10**, 47–53 (2007).
81. Simon, M., Wahlberg, M. & Miller, L. A. Echolocation clicks from killer whales (*Orcinus orca*) feeding on herring (*Clupea harengus*) (L.). *J. Acoust. Soc. Am.* **121**, 749–752 (2007).
82. Hoyt, E. The Worldwide Value and Extent of Whale Watching (Bath, UK: Whale and Dolphin Conservation Society), (1995).
83. Hoyt, E. Whale Watching in *Worldwide Tourism Numbers, Expenditures, and Expanding Socioeconomic Benefits* (Crowborough, UK: International Fund for Animal Welfare, 2001).
84. Vester, H. & Hammerschmidt, K. First record of killer whales (*Orcinus Orca*) feeding on Atlantic salmon (*Salmon salar*) in northern Norway suggest a multi-prey feeding type. *Marine Biodiversity Records* **6**, 1–5 (2012).
85. Ford, J. K. B., Elis, G. & Balcomb, K. *Killer Whales* (UBC Press, 1994).

## Acknowledgements

We thank the USEA – Orques Sans Frontières program for providing the videos used in this study. We are very grateful to Veronique Biquand for her assistance with the statistics and to Hadley Couraud for correcting our English.

## Author Contributions

P.R.L. collected the videos. S.C. scored the videos. S.C., C.B.H. and A.L. analysed the data and wrote the paper.

## Additional Information

**Competing Interests:** The authors declare that they have no competing interests.

**Publisher's note:** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2017