Promises and limitations of nanoparticles in the era of cell therapy: Example with CD19-targeting chimeric antigen receptor (CAR)-modified T cells

Helene Jakobczyk, Flavien Sciortino, Soizic Chevance, Fabienne Gauffre,
Marie-Bérengère Troadec

► To cite this version:

Helene Jakobczyk, Flavien Sciortino, Soizic Chevance, Fabienne Gauffre, Marie-Bérengère Troadec. Promises and limitations of nanoparticles in the era of cell therapy: Example with CD19-targeting chimeric antigen receptor (CAR)-modified T cells. *International Journal of Pharmaceutics*, 2017, 532 (2), pp.813-824. 10.1016/j.ijpharm.2017.07.075 . hal-01617055

HAL Id: hal-01617055

<https://hal-univ-rennes1.archives-ouvertes.fr/hal-01617055>

Submitted on 24 Oct 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

TITLE:

**PROMISES AND LIMITATIONS OF NANOPARTICLES IN THE ERA OF CELL
THERAPY: EXAMPLE WITH CD19-TARGETING CHIMERIC ANTIGEN
RECEPTOR (CAR)-MODIFIED T CELLS**

Authors:

Hélène Jakobczyk ^{a,b}, Flavien Sciortino ^c, Soizic Chevance ^c, Fabienne Gauffre ^c,
Marie-Bérengère Troadec ^{a,b}

Affiliations :

^a Institut de Génétique et Développement de Rennes, UMR 6290 CNRS, Université
de Rennes 1, Rennes, France

^b SFR Biosit UMS CNRS 3480/US INSERM 018, Rennes, France

^c Institut des Sciences Chimiques de Rennes, UMR 6226 CNRS, Université de
Rennes 1, Rennes, France

corresponding author :

Marie-Bérengère Troadec

marie-berengere.troadec@univ-rennes1.fr

ABSTRACT

A number of nanoparticles has been developed by chemists for biomedical applications to meet imaging and targeting needs. In parallel, adoptive T therapy with chimeric antigen receptor engineered T cells (CAR T cells) has recently held great promise in B-cell malignancy treatments thanks to the development of anti-CD19 CAR T cells. Indeed, CD19 is a reliable B cell marker and a validated target protein for therapy. In this perspective article, we propose to discuss the advantages, limits and challenges of nanoparticles and CAR T cells, focusing on CD19 targeting objects: anti-CD19 nanoparticles and anti-CD19 CAR T cells, because those genetically-modified cells are the most widely developed in clinical setting. In the first part, we will introduce B cell malignancies and the CD19 surface marker. Then we will present the positioning of nanomedicine in the topic of B cell malignancy, before exposing CAR T technology. Finally, we will discuss the complementary approaches between nanoparticles and CAR T cells.

KEY WORDS

Nanoparticles, CD19, chimeric antigen receptor, T cell, B cell, cell therapy

46 OUTLINES

47	1 CD19, A B CELL RESTRICTED SURFACE PROTEIN AND A RELIABLE MARKER OF B CELL	
48	MALIGNANCIES	5
49	1.1 THE FUNCTIONS OF B LYMPHOCYTES	6
50	1.2 B CELL MALIGNANCIES	6
51	1.3 THE SURFACE PROTEIN CD19: A VALIDATED TARGET PROTEIN FOR THERAPY	7
52	1.3.1 CD19 structure and function	7
53	1.3.2 Internalization of CD19 after binding to anti-CD19 antibody	8
54	1.3.3 Cells that express CD19	8
55	1.3.4 CD19 as a target for therapy	9
56	2 NANOMEDICINE IN THE TOPIC OF B CELL MALIGNANCY	10
57	2.1 NON TARGETING NANOPARTICLES FOR THERAPY AND IMAGING OF B CELL MALIGNANCY	10
58	2.2 CD19-TARGETING NANOPARTICLES	12
59	2.2.1 Imaging with anti-CD19 nanoparticles	13
60	2.2.2 Therapy with anti-CD19 nanoparticles	14
61	3 CD19-TARGETED CHIMERIC ANTIGEN RECEPTOR (CAR) T CELLS IMMUNOTHERAPY	17
62	3.1 IMMUNOTHERAPIES: ANTIBODY-BASED AND ADOPTIVE CELLULAR THERAPIES	18
63	3.1.1 The concept of CAR T cell: retargeting a cytolytic immune cell by genetic-modification to	
64	eliminate a tumor cell	18
65	3.1.2 The main biological challenges for an effective antibody-based adoptive cellular therapy	18
66	3.1.3 The choice of a relevant target antigen: CD19 gene therapy	19
67	3.1.4 The role of CAR: conferring T cell the ability to persist and expand <i>in vivo</i> and to exert cytolytic	
68	activity	19
69	3.2 CURRENT CLINICAL OUTCOMES, BENEFITS AND LIMITATIONS OF CD19 CAR T THERAPY	21
70	3.2.1 Clinical outcomes	21
71	3.2.2 Advantages	21
72	3.2.3 Limitations	22
73	4 PERSPECTIVES: HOW NANOPARTICLES AND CAR T CELL THERAPY COULD BE	
74	COMPLEMENTARY?	26
75	4.1 MULTIMODALITY	26
76	4.2 TO TRACK MALIGNANT AND CAR T CELLS	27
77	4.3 TO IMPROVE TUMOR ACCESSIBILITY	28
78	4.4 TO MANAGE TOXICITIES OF CAR T CELLS AND MODULATE THE EXPANSION OF CAR T CELLS	30
79	5 CONCLUSION	30
80		

INTRODUCTION

A hematological malignant cell is defined as a hematopoietic cell blocked at an early stage of differentiation and undergoing an uncontrolled clonal proliferation. So far, tremendous improvement in cancer treatment has been obtained thanks to the identification of therapeutic drugs, better molecular understanding of the onset and progression of malignancy, more sensitive detection of tumor cells, more effective follow-up of the disease, better management of adverse effects, optimization of protocol design... Many challenges are still to be undertaken. From the time a patient arrives to be diagnosed to the moment he is cured, physicians and medical staff encounter at least the following issues: the early identification of the tumor, the imaging of malignant cells (where are localized the malignant cells? Is that the primary tumor or a metastasis?), the delivery of therapeutic drugs and avoidance of adverse effects on non-malignant cells (sometimes minimizing the risk of generation of a secondary cancer), and finally the identification of residual cells that could ultimately be at the origin of refractory cancer or relapse.

The chimeric antigen receptor (CAR) T cell therapy is a revolutionary approach of targeted immunotherapy to treat cancer. In CAR T cell therapy, the therapeutic effector is a genetically modified cell. CAR T cell therapy may not yet be poised to overtake chemotherapy as the standard of care, however, it is looking as a promising treatment for certain patients with no other feasible therapeutic option, such as in relapsed or refractory leukemia. An alternative research approach for the treatment of cancer is offered by nanoparticles, which have been proposed as carriers for drug encapsulation in the 60's. Since then, a variety of organic and inorganic nanoparticles, with sizes ranging from *circa* 5 nm to 200 nm, have been designed for

a wide range of applications including targeted drug delivery and imaging, thus boosting the activity of nanomedicine, with some remarkable results particularly in the field of cancer diagnosis and therapy.

In this perspective article, we will propose to discuss the challenges of nanoparticles and CAR T cells in the context of hematological malignancies. We will focus on CD19 targeting objects: anti-CD19 nanoparticles and anti-CD19 CAR T cells because those genetically modified cells are the most widely developed in clinical setting.

In the first part, we will introduce B cell malignancies and their CD19 surface marker, then we will present the positioning of nanomedicine in the topic of B cell malignancy, before exposing CAR T technology. Finally, we will discuss the complementary approaches between nanoparticles and CAR T cells. From the biological point of view, anti-CD19-grafted nanoparticles and anti-CD19 CAR T cells target the same B cell lineage. From the therapeutic perspective, nanoparticles and CAR T cells approaches share common objectives: the optimization of therapeutic effect on target cells and the minimization of adverse effects. However, the mechanisms of action are different (see the graphical abstract). It seems reasonable to conceive that nanoparticles could play a significant role for the potentiation of, and the cooperation with CAR T cell therapy in the future.

1 CD19, A B CELL RESTRICTED SURFACE PROTEIN AND A RELIABLE MARKER OF B CELL MALIGNANCIES

1.1 THE FUNCTIONS OF B LYMPHOCYTES

B cells (also named B lymphocytes) achieve multiple functions that explain their central role in the immune system (Figure 1). Their main role is the production of antibodies to identify and neutralize pathogens. The binding of a B lymphocyte to an antigen triggers an initial step of multiplication and differentiation either into plasma cell which secretes antibodies or into memory B cell. Besides their role in humoral immunity, B cells are involved in cytokine production (e.g. IFN γ , IL6, IL10), antigen presentation to T cells, wound healing, cytokine balance for the differentiation between T lymphocytes (Th1 and Th2 cells), but also in the transplant rejection (review in (LeBien and Tedder, 2008)).

B cells undergo differentiation, from hematopoietic stem cells to plasma cells or memory B cells, through a series of stages characterized by the orderly rearrangement and expression of immunoglobulins genes including CD19 (Figure 1). The development of B cells is also distinguished into different stages by the sequential expression of different transcription factors that induce immunoglobulin gene recombination and the expression of specific surface phenotypes. The onset of B cell lineage occurs in the bone marrow until the immature stage, then mature B cells move into the periphery (*i.e.* out of the bone marrow) (Zhu and Emerson, 2002).

1.2 B CELL MALIGNANCIES

B cell malignancies are hematological cancer characterized by uncontrolled proliferation of B lymphocytes blocked along their differentiation process. B cell malignancies are classified as leukemia (which develops in the bone marrow and disseminates into the body), lymphoma (a cancer of the lymphatic system characterized by the development of a cancer cells in lymph nodes) and myeloma

(cancer of mature B lymphocytes in the bone marrow) (review in (Wang et al., 2012)). B cell malignancies represent 4% of all cancers in adults and 40% of all cancers in children. The clinical outcomes of these cancers under standard chemotherapy depend on the type of B cell malignancies. For instance, children with B-Acute Lymphoblastic Leukemia (ALL) have an overall good prognosis, but some of them are refractory to chemotherapy or develop multiple relapses and have a poor prognosis (review in (Park et al., 2016)). Relapsed or refractory B cell ALL in adults are associated with a poor prognosis (review in (Geyer and Brentjens, 2016)).

1.3 THE SURFACE PROTEIN CD19: A VALIDATED TARGET PROTEIN FOR THERAPY

1.3.1 CD19 structure and function

CD19 is a 95 kDa transmembrane glycoprotein of the immunoglobulin superfamily composed of an extracellular domain, a single transmembrane domain, and a cytoplasmic domain (Stamenkovic and Seed, 1988). CD19 belongs to the CD19 complex on the surface of B cells with CD21 and CD81 proteins (Figure 2). CD19 activation induces two downstream pathways. The first cascade of activation is dependent on the B Cell Receptor (BCR). The BCR is composed of a membrane immunoglobulin and a signaling subunit composed of a heterodimer of immunoglobulin alpha and beta. The BCR plays a role as antigen receptor and CD19 is a co-receptor for BCR signal transduction (review in (Wang et al., 2012)). The second pathway depending on CD19 is independent of the BCR: the CD19 complex is able to bind activated complement fragment C3d and modulates BCR signaling (review in (Wang et al., 2012)).

1.3.2 Internalization of CD19 after binding to anti-CD19 antibody

CD19 proteins on the surface of each B lineage leukemia/lymphoma cells are rapidly internalized upon ligation with anti-CD19 antibodies or immunoconjugates (Uckun et al., 1988; Yan et al., 2005), and are ultimately taken up by lysosomes (Carter, 2006 ; Gerber et al., 2009; Hong et al., 2015).

1.3.3 Cells that express CD19

CD19 is a B cell-specific protein expressed early in B cell ontogeny (Stamenkovic and Seed, 1988) (**Figure 1**). CD19 transcripts are restricted to members of the B cell lineage and are not expressed in other hematological lineages including normal myeloid, erythroid, megakaryocytic, or multilineage bone marrow progenitor cells (Uckun et al., 1988). CD19 protein is found on the surface of B cells from the proB cell stage until plasma cell differentiation of the B lineage (Tedder et al., 1994). Several hundred thousand CD19 proteins can be found on the surface of each B-lineage leukemia/lymphoma (Uckun et al., 1988)(review in (Li et al., 2017)). All resting B cells display CD19 antigens, and CD19 expression persists upon activation, but is lost upon further differentiation to immunoglobulin-secreting plasma cells (Stamenkovic and Seed, 1988). CD19 is also more abundant in pre-B cell lines and less abundant in plasmacytomas (Stamenkovic and Seed, 1988). Almost all early B cell malignancies show CD19 expression at normal to high levels: 80% of ALL, 88% of B cell lymphomas and 100% of B cell leukemias (review in (Wang et al., 2012)). However its expression decreases in myeloma cases (review in (Wang et al., 2012)).

1.3.4 CD19 as a target for therapy

Twenty years ago, CD19 was already proposed as a « suitable target for immunotoxin-mediated treatment of aggressive forms of B cell lymphomas and leukemia that responds poorly to conventional chemotherapy» (Uckun et al., 1988). Currently, CD19 antibody-based therapy has become reality to treat B cells malignancy. In the 2010's, various strategies harnessing the potential of targeting B cells restricted to CD19 antigen were in development: antibody-drug conjugate, Fc-engineered human CD19 antibody with antibody-dependent cell-mediated cytotoxicity, chimeric antigen receptor, etc. (Hammer, 2012). The most advanced anti-CD19 therapy is the Blinatumomab (BLINCYTO®, Amgen) (review in (Hammer, 2012)) (Goebeler and Bargou, 2016), a bispecific CD19-directed CD3 T cell engager (BiTE) antibody construct. Blinatumomab binds specifically to CD19 expressed on the surface of cells of B-lineage origin, and to CD3 expressed on the surface of T cells. It brings both cells in contact so that the activated T cells can kill the B cells. Blinatumomab is approved by the US Food-and-Drug-Administration (FDA) and the European Commission (EC) for the treatment of Philadelphia chromosome-negative relapsed or refractory B-ALL, in adults (USA and EC) as well as in children (USA only). Additionally, anti-CD19 antibodies are also in development for radioimmunotherapy in preclinical studies. ¹³¹I-labeled anti-CD19 antibody has been largely explored for conventional ¹³¹I radioimmunotherapy because antigen rapidly internalizes upon binding of antibody – resulting in catabolism and release of ¹³¹I (Scheinberg and Strand, 1983). Moreover, ⁹⁰Y-particle-labeled anti-CD19 antibody has shown an efficacy comparable to ⁹⁰Y-labeled anti-CD20 antibody in radioimmunotherapy of mice with xenografts of human B lymphoma cell lines (Ma et al., 2002).

226

227 **2 NANOMEDICINE IN THE TOPIC OF B CELL MALIGNANCY**

228

229 A number of nanoparticles has been proposed by chemists for cancer diagnostics
230 and therapeutics, as summarized **Table 1**. Organic nanoparticles, such as
231 liposomes, oil-in-water emulsions or polymeric particles, are mainly used as carriers,
232 whereas nanoparticles, such as superparamagnetic iron oxide nanocrystals or
233 quantum dots, show interesting intrinsic properties for imaging and therapy.

234

235 2.1 **NON TARGETING NANOPARTICLES FOR THERAPY AND IMAGING OF B CELL**

236 **MALIGNANCY**

237 Some anticancer encapsulation nanosystems have made their way to the market
238 (Pattni et al., 2015). Liposomal formulations encapsulating drugs, such as
239 doxorubicin, are commercialized under the name of Myocet, Doxil, Lipodox and
240 Caelyx. Related to hematological malignancy, a phase III clinical trial is open for a
241 liposome combinational delivery of two cytotoxic drugs (cytarabine and daunorubicin)
242 for high risk acute myeloid leukemia (clinicaltrials.gov identifier NCT01696084) (Shi
243 et al., 2017). With the ultimate goal of achieving both spatial and temporal control of
244 drug delivery, nanocarriers have evolved from the mere "sustained" release to
245 "triggered" release (**Figure 3**). Indeed, in cancer, abnormal local conditions, such as
246 pH, enzymatic activity or concentration in reactive oxygen species, can trigger the
247 delivery of the drug. In addition to these endogenous signals, nanocarriers can also
248 release their load on the effect of applied light, ultrasounds or a magnetic field
249 (Bhattacharya et al., 2016; Kamaly et al., 2016).

250 In the topic of B cell malignancy, only few nanoparticles-based therapies are in
251 development (Stephenson and Singh, 2017) (Shi et al., 2017). Among all the recent
252 clinical-stage nanomedicines (Shi et al., 2017), a phase II clinical trial is open to
253 evaluate a liposome, carrying a DNA oligonucleotide against the anti-apoptotic
254 protein BCL-2, in relapsed or refractory B cell lymphomas (clinicaltrials.gov identifiers
255 NCT01733238 and NCT02226965). Similar approaches of gene/RNAi delivery by
256 silica-based nanoparticles to target B-cell lymphoma were described in mouse model
257 (Martucci et al., 2016). Additionally, between 2011 and 2014, a phase I/II clinical trial
258 was opened to evaluate the safety and tolerability of a poly(ethylenimine)-based
259 transfecting polyplex carrying siRNA against eIF5A and a plasmid expressing a pro-
260 apoptotic mutant of eIF5A under the control of a B cell specific promoter. This
261 therapeutic agent was evaluated in relapsed or refractory B cell malignancies
262 (clinicaltrials.gov identifier NCT01435720). Finally, an immunostimulant lipoplex
263 composed of liposome and plasmid DNA (Chang et al., 2009) is in a phase I clinical
264 trial in relapsed or refractory leukemia (clinicaltrials.gov identifier NCT00860522).

265 Tumors are currently diagnosed using various imaging modalities such as
266 radiography, computed tomography (CT), positron emission tomography (PET) and
267 magnetic resonance imaging (MRI) (Salem et al., 2014)(Navarro et al., 2017).
268 However, the diagnosis of hematological malignancies can be challenging due to the
269 diversity of imaging appearances and clinical behavior of these diseases (Navarro et
270 al., 2017). Multimodal imaging approaches have been proposed to overcome these
271 limitations, since they offer the ability to image with different resolutions and over
272 different temporal and spatial scales. Cistaro *et al.* demonstrated the high potential of
273 combined PET (using ^{18}F -fluorodeoxyglucose) and MRI (using paramagnetic contrast
274 agent) in the evaluation of pediatric patients with ALL (Cistaro et al., 2017). By their

work, they highlighted the real need of developing hybrid PET/MRI instruments and dual contrasts agents.

In line with that idea, a variety of nanoparticles has been designed to combine several imaging modes, multiple therapies, (e.g. photothermal therapy and conventional chemotherapy) or imaging and therapeutic functions (theranostics) and therefore holds great prospects in cancer treatment (Riley and Day, 2017). Among others, our group has recently reported on a vesicular platform, with a shell of inorganic nanoparticles named Hybridosomes® (Sciortino et al., 2016). The large number of nanoparticles forming the shell is a clear advantage for imaging applications, since an enhanced contrast is observed. Initially designed for MRI, these Hybridosomes® can not only be prepared from iron oxide superparamagnetic nanoparticles but also from any types and combinations of inorganic particles with imaging or therapeutic properties. Therefore, those multimodal nano-objects are suitable tools for multimodal imaging as well as theranostics. The feasibility of a theranostic approach has been demonstrated in acute myeloid leukemia patients where *in vivo* molecular imaging of CXCR4, a crucial protein involved in the retention of hematopoietic stem cells within the hematopoietic niche, has been achieved by means of positron emission tomography (Herhaus et al., 2016). However, as far as we know, there is still no open clinical trial using those combined strategies in B cell malignancies.

2.2 CD19-TARGETING NANOPARTICLES

The efficiency of imaging and treatment can be greatly improved by targeting specifically the malignant cells. As mentioned above, CD19 is currently the antigen of

choice used to target B cells. Recently, CD19-targeting nanoparticles were designed for nanomedicine by grafting anti-CD19 antibody or its derivatives (Fab, F(ab)₂...) to the nanoparticles (Figure 4). As an example, Cheng *et al.* produced liposomal doxorubicin targeted *via* anti-CD19 monoclonal antibody fragments: either the single-chain variable fragment (scFv), or the variable fragment (Fab), or the monoclonal antibody (mAb) (Figure 4). The authors compared the efficacy of the three targeted constructs and concluded that the scFv single-chain variable fragment would be more suitable for development of immunotherapy for the following reasons: i) it contained less foreign peptides, ii) the production was easier, and iii) the cost of production was more economical thanks to the expression in bacterial systems (Cheng and Allen, 2008). Typically, four types of chemical functions from the antibody or its derivatives (-NH₂, -COOH, -SH, -carbohydrates) can be used for covalent grafting to the nanoparticle. The use of spacers such as PEG derivatives lowers the risk of antibody inactivation (Chen *et al.*, 2016; Manjappa *et al.*, 2011)(Nguyen *et al.*, 2010)(Hong *et al.*, 2015). Alternative strategies were also proposed, as the noncovalent streptavidin/biotin conjugation (Procko *et al.*, 2014) (Dong *et al.*, 2014).

2.2.1 Imaging with anti-CD19 nanoparticles

Few anti-CD19 grafted nanoparticles for *in vitro* imaging have been published so far. Nguyen *et al.* designed pegylated SERS (Surface Enhanced Raman Scattering) gold nanoparticles conjugated to human anti-CD19 antibody that showed specific *in vitro* targeting towards chronic lymphocytic leukemia (CLL) (Nguyen *et al.*, 2010; Walker *et al.*, 2012). The functional SERS nanoparticles were composed of a gold core onto which a reporter dye was adsorbed. The signals were detected by dark-field

microscopy and Raman spectrometry and showed no interference with conventional fluorescent stains used in histology. Ramos B cells labeling through anti-CD19 mediator was demonstrated by Dong *et al.* by grafting an anti-CD19 antibody onto Ag@SiO₂ core-shell nanoparticles (Dong et al., 2014). In this study, the authors monitored the metal-enhanced fluorescence of a reporter (rhodamine B) adsorbed on the surface of the nanoparticles. However, to the best of our knowledge, *in vivo* imaging using anti-CD19-grafted-nanoparticles has not been reported yet.

2.2.2 Therapy with anti-CD19 nanoparticles

2.2.2.1 Chemotherapy: drug delivery

Nanoparticles decorated with anti-CD19 have already been reported as effective carriers for drug delivery on *in vitro* models and preclinical studies (Table 2). Doxorubicin, an inhibitor of topoisomerase involved in DNA synthesis, is frequently the drug of choice for proof-of-concept, as the cytotoxic effect of this drug is well demonstrated on B cells. A doxorubicin loaded immunoliposome targeting B lymphocytes showed a 6-fold more cytotoxic *in vitro* activity on B cells than non-targeted liposomes (Lopes de Menezes et al., 1998). Similar results were observed *in vivo* with an improved survival of mice injected with anti-CD19-doxorubicin-liposomes compared to non-targeted liposomes or free doxorubicin treatments (Lopes de Menezes et al., 1998). Doxorubicin was also encapsulated into block-copolymer nanoparticles grafted with anti-CD19. A clathrin-dependent internalization pathway was identified, suggesting that the physiological internalization pathway of CD19 was conserved. In comparison to the administration of free doxorubicin, both improved *in vitro* apoptosis of CD19 positive cells and better survival of treated mice were demonstrated (Krishnan et al., 2015). *In vivo*, mice xenografted with B cells and

exposed to anti-CD19-liposomes containing doxorubicin or vincristine demonstrated a higher cell cytotoxicity and showed a longer survival time than mice exposed to free drug (Sapra and Allen, 2004). Those anti-CD19-liposomes showed *in vitro* a greater binding, a more effective internalization and an equivalent cytotoxicity on B cells compared to anti-CD20-liposomes (Sapra and Allen, 2004).

Other inhibitors of B-cells than doxorubicin or vincristine have also been evaluated and incorporated into nanoparticles. As an example, the C61 molecule (1,4-bis (9-O-dihydroquinidiny) phthalazine/hydroquinidine 1,4-phthalazinediyl diether) was identified as a potent inhibitor of the cytoplasmic protein SYK (spleen tyrosine kinase), an important regulator of B cell apoptosis (Table 2). Myers *et al.* demonstrated that a liposomal nanoparticle formulation entrapping C61 and decorated with anti-CD19 caused *in vitro* the apoptosis of pre-B ALL cells, twice more than the non-decorated liposomes (Myers *et al.*, 2014). Immunocompromised NOD/SCID mice were then xenografted with pre-B ALL cells, and injected with C61-liposomes decorated with anti-CD19. Tumor cell viability decreased and mice did not develop leukemic splenomegaly, thus showing a better therapeutic efficacy than irradiation with 2Gy γ -rays. In addition, the combination of C61 loaded anti-CD19-liposomal nanoparticles, with exposure to low dose of radiations, caused the abrogation of B leukemia in engrafted mice (Myers *et al.*, 2014).

In addition, multifunctional immunoliposomes grafted with several antibodies were shown to exhibit higher selectivity, greater binding affinity, and enhanced apoptosis induction of B-CLL cells (Woyach *et al.*, 2014). Yu *et al.* also proposed a dual ligand conjugation on immunoliposomes (Yu *et al.*, 2013). The authors first evaluated the level of expression of CD19, CD20 and CD37 antigens in several B cell lines and

primary B-CLL cells, and found comparable level for CD19 and CD37. They also calculated the internalization rate of the three antibodies in lymphoma cells (Raji cells) and confirmed the choice of anti-CD37 as the primary ligand for specific targeting of B cells. Then they measured the binding efficacy of single or mixtures of anti-CD19, anti-CD20 and anti-CD37 on B-CLL cells isolated from patients. Greater binding efficacies occurred with dual combinations of anti-CD19 and anti-CD20, with anti-CD37 antibody. The antibody ratio was finally optimized to improve this synergetic effect.

Note that the combination of several specific antibodies is also a promising strategy to overcome the variability in the expression of target antigens among patients. In this context, hydroxychloroquine, an anti-malaria and anti-rheumatic drug, has been encapsulated in order to overcome pharmacokinetic obstacles and to deliver a larger amount of this apoptotic drug into B-CLL cells from patients. As an example, Mansilla *et al.* encapsulated hydroxychloroquine in PEG-PLGA nanoparticles mono-functionalized by anti-CD19 antibody or bi-functionalized by anti-CD19 and anti-CD20 antibodies (Mansilla *et al.*, 2010). The authors showed a significant induction of apoptosis of B-CLL cells with mono- or bi-functionalized nanoparticles compared to non-functionalized nanoparticles.

2.2.2.2 Nanoparticle-based immunotherapy

An innovative strategy consists in using nanoparticles exposing antibodies in order to stimulate the production of lymphocytes, or even to bridge malignant cells to killer T cells (see the graphical abstract). Schütz *et al.* designed nanoparticles termed antigen-specific T cells redirectors (ATR). The ATR nanoparticles were conjugated to

two antibodies, an anti-TCR and an anti-CD19. The ATR nanoparticles provided a physical proximity between T cells and tumor cells, and redirected T cells to kill tumor cells (Schütz et al., 2016). *In vivo* assays on mice xenografted with lymphoma cells and injected with ATR nanoparticles showed smaller tumors and an improved survival compared to control mice.

3 CD19-TARGETED CHIMERIC ANTIGEN RECEPTOR (CAR) T CELLS IMMUNOTHERAPY

An alternative to nanoparticles for targeting tumor cells is to take advantage of other cells. For years, most of hematological neoplasms have been treated by hematopoietic stem cell transplantations. The transplanted allogeneic hematopoietic stem cells kill residual malignant cells by a graft-versus-tumor effect. This cell therapy approach, used to fight leukemia, lymphoma or myeloma, leads to either remission or immune control of the malignancy; however, some patients relapse. On the other hand, many therapeutic approaches tend to modulate the immune response to eliminate tumor cells. Immunotherapy has marked the past years by generating extraordinary advances in clinical applications for cancer treatment.

Cell immunotherapy harnesses the power of both cell therapy and immunotherapy, and is at the origin of tremendous clinical progresses in the past decade (Ramachandran et al., 2017). For the purpose of the review, we will focus on CD19 antibody-based cell immunotherapies that target B cell neoplasms.

3.1 IMMUNOTHERAPIES: ANTIBODY-BASED AND ADOPTIVE CELLULAR THERAPIES

3.1.1 The concept of CAR T cell: retargeting a cytolytic immune cell by genetic-modification to eliminate a tumor cell

T lymphocytes are cells that play a central role in cell-mediated immunity. Different subsets of T cells achieve cytolytic, regulatory or memory roles. Genetically retargeting T cells against tumor surface antigens to trigger cytotoxic mechanisms against malignant cells is one of the principles of adoptive cell therapy. More precisely, the engineering of T cells to express a chimeric antigen receptor (CAR) is the most common gene-modifying strategy that is being investigated. CARs are synthetic receptors that direct the genetically engineered T cells against tumor surface antigens, for instance CD19 antigen. Adoptive cell therapy using gene-modified T cells has emerged as an exciting therapeutic approach for the treatment of cancer (Porter et al., 2011; Kochenderfer et al., 2012 ; Brentjens et al., 2013).

3.1.2 The main biological challenges for an effective antibody-based adoptive cellular therapy

Conceptually, many challenges should be faced to achieve an *in vivo* therapeutic efficacy. The first one is that CAR T cells must be able to persist *in vivo*, and then undergo cellular expansion (Grupp et al., 2013). They will also have to infiltrate tumor tissues (in case of solid tumors), then to engage their target antigen expressed on tumor cells, and finally, to exert their cytolytic, proliferative, and cytokine secretory activities within the tumor microenvironment to eliminate malignant cells (review in (Beatty and O'Hara, 2016)).

Adoptive T cell therapy with chimeric antigen receptor engineered T cells (CAR T

cells) has shown substantial clinical results against B cell malignancies (Porter et al., 2011; Kochenderfer et al., 2012 ; Brentjens et al., 2013). The fact that CAR T cell therapy approach has proven to be of some effectiveness across a range of hematological malignancies (Gill and June, 2015) may be partly explained by the choice of a relevant target antigen (for instance CD19) and by the fact that those malignancies reside in the natural sites that adoptively transferred T cells naturally invade (review in (Newick et al., 2016)).

3.1.3 The choice of a relevant target antigen: CD19 gene therapy

As mentioned previously, CD19 is a reliable target antigen for antibody-based therapy (review in (Hammer, 2012)(Li et al., 2017)). More than half of all CAR-modified T cell studies in hematological malignancies have targeted CD19 antigen (review in (Beatty and O'Hara, 2016)). CD19-specific CAR T cells have demonstrated potent activity in B cell ALL and lymphomas including CLL and non-Hodgkin lymphoma (Porter et al., 2011 ; Grupp et al., 2013; Maude et al., 2014 ; Davila et al., 2014 ; Lee et al., 2015 ; Brudno et al., 2016 ; review in Beatty and O'Hara, 2016).

3.1.4 The role of CAR: conferring T cell the ability to persist and expand *in vivo* and to exert cytolytic activity

3.1.4.1 Design of CAR

The chimeric antigen receptor (CAR) is composed by two main modules: (i) an extracellular component that recognizes a cell surface protein (e.g. CD19) (this extracellular moiety is a single-chain variable fragment (scFv) derived from an antibody) linked to (ii) an intracellular component consisting in T cell signaling

domains of the T cell receptor (e.g. CD3 ζ) including co-stimulatory domains (e.g. CD28, or 4-1BB) involved in T cell activation (Figure 5) (review in (Beatty and O'Hara, 2016) and (Geyer and Brentjens, 2016)). The extracellular component is responsible for redirecting T cell specifically to the human tumor antigen whereas the intracellular component sustains T cell activation, supporting cell expansion and cytokine release resulting in cytolytic activity.

Intense work is done to optimize each module: the extracellular component which acts as the target-binding domain of the CAR, the hinge region connecting extracellular and intracellular component (Hudecek et al., 2013), and the intracellular component for an effective T cell proliferation and differentiation to mature effector T cells. The successive generations of CD19 CAR T differ in the number and origin of the intracellular co-stimulatory domains (Figure 5) (e.g. 4-1BB or CD28) (Savoldo et al., 2011 ; Porter et al., 2011 ; Maude et al., 2014 ; Park et al., 2016).

3.1.4.2 Mechanism of action of CAR T cells

The binding of the anti-CD19 scFV to CD19 antigen of tumor cell surface (the resulting complex is named the immune synapse) sends a signal through the CAR to the effector T cell. This signal results in the activation of the T cell and in the release of soluble molecules, perforin, granzyme and pro-apoptotic ligands, that kill the tumor cells. Additionally, activated T cells secrete proinflammatory cytokines (e.g. interferon IFN- γ , and IL-2), amplifying the immune response (Davenport et al., 2015) (Geyer and Brentjens, 2016), and leading to the expansion of CAR T cells. The range of *in vivo* expansion of CAR T cells has been reported between 100- to 10 000- fold (Grupp et al., 2013).

3.2 CURRENT CLINICAL OUTCOMES, BENEFITS AND LIMITATIONS OF CD19 CAR T THERAPY

3.2.1 Clinical outcomes

Many patients go into remission with standard chemotherapy for B cell malignancies. However, children and adults with relapsed or refractory B cell ALL have a poor prognosis. Substantial clinical efficacy has been demonstrated with a therapy based on CAR-modified T cells targeted to CD19. Approximately 70% of patients underwent complete or at least partial response to treatment with chimeric antigen receptor CAR-modified T cells targeted to CD19 (Porter et al., 2011; Kochenderfer et al., 2012 ; Brentjens et al., 2013 ; Grupp et al., 2013 ; Maude et al., 2014 ; Davila et al., 2014 ; Lee et al., 2015). Results are less impressive with CLL or with B cell non-Hodgkin lymphoma but still subsets of patients show significant benefits (review in (Geyer and Brentjens, 2016). Clinical trials are ongoing for multiple myeloma.

3.2.2 Advantages

In vivo expansion and persistence of CAR T cells is a clear determinant of clinical benefit (Grupp et al., 2013 ; Porter et al., 2015 ; Beatty and O'Hara, 2016). In addition, the natural trafficking of CAR T cells within the blood, lymph nodes, and bone marrow where they encounter malignant cells also favors the efficacy of the therapy (Beatty and O'Hara, 2016). Furthermore, it appears that the accessibility to malignant cells is less hindered by the tumor microenvironment in those tissues compared to solid tumors (Geyer and Brentjens, 2016 ; Newick et al., 2016).

3.2.3 Limitations

3.2.3.1 Genetic modification of autologous T cells

First, each patient is infused with his own T cells. This specificity limits any large-scale manufacturing process and anticipated stocks. Then, autologous T cells are subjected to genetic modifications by retrovirus, lentivirus or non-viral gene transfer followed by *in vitro* stimulation. Currently, the complicated and individualized production of autologous CAR T cells may be one, among others, of the bottlenecks that reduce accessibility to this personalized therapy to many people. Some strategies using universal T cells (*i.e.* that do not come from the patient) are also in development. Suboptimal expression of the CAR at the surface of CAR T cells may also limit the benefit of CAR T cell therapies. Recently, Eyquem *et al.* have proposed that directing a CD19-specific CAR to the T cell receptor α constant (*TRAC*) locus not only results in uniform CAR expression in human peripheral blood T cells, but also enhances T cell potency, with edited cells vastly outperforming conventionally generated CAR T cells in a mouse model of ALL (Eyquem *et al.*, 2017).

3.2.3.2 The need of lymphodepletion for the patient

The purpose of chemotherapy, whose objective is to achieve lymphodepletion prior to CAR T cells infusion, is to create a more favorable environment for CAR T cells. Most studies corroborated the notion that host lymphopenia (*i.e.* a low number of lymphocytes in the blood) facilitates the expansion of adoptively transferred T cells. Whether lymphodepletion might further enhance the activity of CAR T cells in this setting remains unclear (Brudno *et al.*, 2016 ; Turtle *et al.*, 2016). To date, induction

of lymphodepletion prior to infusion of CAR T cells continues to be often incorporated in clinical trials using CAR T cells.

3.2.3.3 *Toxicity for the patient*

The medical community will have to overcome clinical challenges related to CD19-targeted CAR T cells (Geyer and Brentjens, 2016; Park et al., 2016). Major side-effects, particularly cytokine release syndrome, neurological toxicities, and B cell aplasia have been reported in all clinical trials using CD19-targeted CAR T cells. The cytokine release syndrome is a severe inflammatory response syndrome that appears within the hours to days following CAR T cell infusion. Clinical features include fevers, muscle pain, malaise, and, in more severe cases, hypoxia, hypotension, and occasionally renal dysfunction and coagulopathy. The cytokine release syndrome is characterized by elevation of pro-inflammatory cytokines (e.g. IL-6) and T cell activation and expansion. Tumor burden is positively correlated with the risks of severe cytokine release syndrome and neurotoxicity (Brentjens et al., 2013) (Turtle et al., 2016). The cytokine release syndrome can be life-threatening and requires intensive supportive care. Mitigating strategies to reduce cytokine release syndrome frequency and severity comprise anti-IL-6 receptor antibody, steroids, and possibly a protocol-specified algorithm to potentially start pre-emptive treatments (Maude et al., 2014; Lee et al., 2014 ;Turtle et al., 2016; Ruella et al., 2017).

Reversible neurologic toxicity has been observed after CAR T cell infusion, including delirium, seizure-like activity, confusion, word-finding difficulty, or aphasia.

Finally, CD19-targeted CAR T cells therapy shows “on-target, off-tumor” toxicity that generates B cell aplasia (Porter et al., 2011 ;Grupp et al., 2013; Maude et al., 2014).

Limiting B cell aplasia for CD19-targeted CAR T cells has been successfully managed with intravenous immunoglobulin replacement therapy (Frey and Porter, 2016). Novel approaches to limit B cell aplasia are under investigation as the use of antigen-specific inhibitory CAR to protect normal B cells (Fedorov et al., 2013).

3.2.3.4 CD19-antigen escape

Loss of expression of the CD19-target antigen resulting in an antigen escape (e.g. CD19-negative relapse) may limit the benefit of CD19 CAR T cells therapy (Grupp et al., 2013). Tumor antigen escape has emerged as a main challenge for the long-term disease control (review in (Wang et al., 2017; Velasquez and Gottschalk, 2017)). Studies are going on to understand the mechanism of loss of CD19 expression and overcome this difficulty. Braig *et al.* reported emergence of CD19-relapses due to CD19 mRNA splice variants (Braig et al., 2017). Zah *et al.* proposed a design of bispecific CARs that triggered robust cytotoxicity against target cells expressing either CD19 or CD20 and controlled both wild-type B cell lymphoma and CD19 mutants with equal *in vivo* efficacy (Zah et al., 2016).

3.2.3.5 Infused dose, composition, and control of expansion and function of CAR T cells

So far, the different clinical trials have not led to the identification of a clear correlation between higher CAR T cell infused dose and greater efficacy or CAR T cell persistence (Porter et al., 2011 ; Grupp et al., 2013; Maude et al., 2014 ; Davila et al., 2014 ; Lee et al., 2015 ; Brudno et al., 2016) (review in (Park et al., 2016; Geyer and Brentjens, 2016)). Importantly, the efficacy of CAR T cells relies on their activation in response to CD19 antigen and expansion *in vivo*, making the magnitude of their reactivity unpredictable (Grupp et al., 2013). For instance, anti-CD19 CAR T

cells have been shown to proliferate in excess of 100,000-fold in some patients, ultimately accounting for over 50% of circulating lymphocytes. The lack of control over CAR T cells activation and expansion *in vivo* is a limit to predict the therapeutic response.

Multiple parameters provide clues to explain this unpredictability. The composition of the infused therapeutic agent is source of variability. So far, CAR T cells are generated from autologous T cells, making the received therapeutic agent different for each patient (Sommermeyer et al., 2016) (Turtle et al., 2016). In preclinical studies, where mice were injected with a same pool of CAR T cells, a better correlation between the infused dose and the xenografted mouse survival was observed (Sommermeyer et al., 2016). More precisely, the variability of CAR T cells encompasses extrinsic parameters, from the efficacy of genetic modification to the expression of the CAR at the surface of CAR T cells, but also intrinsic interindividual parameters including composition of CD4+ and CD8+ T cells. In CAR T therapy, CD4+ CAR T cells are responsible for cytokine production whereas CD8+ CAR T cells trigger direct antitumor effects. The ratio of CD4+/CD8+ CAR T cell subsets may be of importance in the balance between efficacy and toxicity (Park et al., 2016). In most reported trials, patients received CAR T products comprising random compositions of CD4+ and CD8+ T cells. In contrast, Sommermeyer *et al.* and Turtle *et al.* showed that CAR T cell products generated from defined T cell subsets (1:1 ratio of CD4+ and CD8+ CAR T cells) can provide uniform potency compared with products derived from unselected T cells and induce complete remission without a high rate of toxicity in patients with a high tumor burden (Sommermeyer et al., 2016 ; Turtle et al., 2016). Approaches to limit expansion and activation are also underway. Rodgers *et al.* propose a method to control CAR T cells using peptide-engrafted

antibody-based molecular switches that act as a bridge between the target cell and CAR T cells (Rodgers et al., 2016).

Interindividual variation in response to the treatment can also be attributed to difference in lymphodepletion between each patient, or to difference in immunological clearance that will impact the persistence of the infused and expanded CAR T cells.

Altogether, the optimal dose and composition of the CAR T cell product remain under development in order to achieve a better predictability in response to the therapeutic agent and to balance toxicity and efficacy.

4 PERSPECTIVES: HOW NANOPARTICLES AND CAR T CELL THERAPY COULD BE COMPLEMENTARY?

4.1 MULTIMODALITY

The efficacy of CAR T cell therapy relies on the multimodality of the therapeutic response. CAR T cells target tumor cells, trigger cytolytic activity, and ensure their own expansion. We can envision that the future of nanomedicine will benefit from the same feature: the multimodality. It is clear that nano-objects, and among them Hybridsomes® (Sciortino et al., 2016), can address many of the challenging issues of hematological cancer diagnosis and therapy. In particular, nanoparticles could play a significant role for the potentiation of, and the cooperation with CAR T cell therapy. Their complementarity (in terms of function, distribution and time of administration) can be envisioned to fulfill at least three objectives: (i) to track malignant cells and

CAR T cells to monitor their biodistribution and expansion, (ii) to increase tumor accessibility, and (iii) to manage CAR T cell toxicity and modulate the expansion of CAR T cells.

4.2 TO TRACK MALIGNANT AND CAR T CELLS

Since the proof-of-concept of CAR T cells has been validated, current developments include the control of cell expansion or avoidance of CD19 escape. There is a need for noninvasive tracking of the transfused T cells in patients to determine their biodistribution, viability, and functionality (review in (Liu and Li, 2014)). Several strategies based on nanoparticle contrast agents have been proposed using either *ex vivo* preloaded nanoparticles on CAR T cells, or *in vivo* administration of nanoparticles after CAR T cell infusion. For instance, in mouse model, CAR T biodistribution has been monitored through radiolabeled-nanoparticles or contrast-agent-nanoparticles loaded into CAR T cells prior to cell infusion (Bhatnagar et al., 2013;Bhatnagar et al., 2014).

Furthermore, detecting the localization of tumor cells is of particular importance in the case of hematological cancer, since hematological malignant cells are intrinsically disseminating. In addition, *in situ* imaging alternatives to the invasive sampling of bone marrow are desirable for diagnosis and for residual disease follow-up. By proposing efficient targeting contrast agents, nanomedicine can greatly improve the diagnosis, and beyond, the determination of localization of tumor cells (Kobayashi et al., 2005).

4.3 TO IMPROVE TUMOR ACCESSIBILITY

A recent statistical review of the literature revealed that less than 1% of the injected nanoparticles systemically reaches the malignant cells in solid tumors, compromising their translation into clinical use (Wilhelm et al., 2016). This figure is due both to nanoparticle uptake by the immune system, and to their poor mobility into the tumor microenvironment. Although hematological malignancies differ from other solid tumors, some limitations of the CAR T therapy due to limited access to specific accumulation sites may be observed as well. According to cancer type, hematological malignant cells originate from the bone marrow (e.g. leukemia, myeloma) or lymph node (e.g. lymphoma), and infiltrate blood stream and solid tissues. The bone marrow niche is a very complex environment essentially composed of a dense network of small arterioles and sinusoids, and of various cell types within an extracellular matrix (Wu et al., 2008) (Morrison and Scadden, 2014) (Gattazzo et al., 2014)(Schepers et al., 2015). Leukemic stem cells, as well as hematopoietic stem cells, are dependent on those cells and extracellular components for their emergence, homing and survival. Disruption of those interactions participates in the efficacy of the therapy.

The combination of the specific properties of CAR T cells and nanoparticles seems promising to enhance the efficacy of treatments. Indeed, CAR T cells will guarantee longer circulation time in the blood stream and specific recognition of B cells, whereas nanoparticles can bring advantageous features such as degradation of the extracellular matrix, disruption of cell-cell interactions, or thermal stimulation. An advance in this direction was already reported in the literature. In mouse studies, Kennedy *et al.* used T cell as chaperones for gold nanoparticle delivery to enhance the efficacy of nanoparticle-based photothermal therapies and imaging applications

688 by increasing accumulation at tumor site (Kennedy et al., 2011). Another innovative
689 strategy, inspired by motile and invasive cells, would be the active enzymatic
690 degradation of the tumor matrix by protease that can be associated with the
691 nanotherapeutic system. For instance, iron oxide nanoparticles coated with
692 collagenase were magnetically driven through *in vitro* extracellular matrix, at a rate
693 similar to invasive cells (Kuhn et al., 2006). Other proteolytic surfaces include
694 bromelain, an enzymatic complex belonging to the papain family and extracted from
695 pineapple which contains a mixture of 9 proteases with distinct pH and enzymatic
696 activities (Parodi et al., 2014). Local heating triggered by external sources can also
697 be used to alter the tumor environment and enhance accessibility to malignant cells,
698 based on gold nanoparticles (Gormley et al., 2013; Smith et al., 2015).

699 An alternative strategy would be the pretreatment with therapeutic nanoparticles prior
700 to CAR-T infusion. In this line, nanoparticles targeting the bone marrow niche could
701 also be utilized to specifically deliver high doses of lymphodepleting agents prior to
702 CAR T infusion. Similarly, pre-treatment with drugs, specifically targeting the
703 interaction of leukemic stem cells with their bone marrow niches, may be useful to
704 mobilize those cells and render them more accessible to CAR T cells in the marrow
705 or the blood stream. Among others, inhibitors of the adhesion molecule E-selectin, or
706 inhibitors of the chemoattractant stromal-cell-derived factor 1 (SDF-1) could be
707 proposed because leukemic stem cells are dependent on those molecules for their
708 homing (Sipkins et al., 2005)(Krause and Scadden, 2015)(Schepers et al., 2015).

709 Identification of additional specific factors in B cell malignancies could be of interest
710 for mobilizing B cells and enhancing CAR T cell therapy, as exemplified by the role of
711 CD44, or various selectins and their ligands in chronic myeloid leukemia or acute
712 myeloid leukemia (Krause et al., 2006)(Jin et al., 2006)(Krause et al., 2013).

713

714 4.4 **TO MANAGE TOXICITIES OF CAR T CELLS AND MODULATE THE EXPANSION OF**
715 **CAR T CELLS**

716 Major toxicity such as severe cytokine release syndrome is intrinsically related to
717 CAR T efficacy, and current developments aim at controlling it. Current strategies to
718 allow preferential removal of CAR T cells include genetic “safety switch” or drug
719 sensitivity (review in (Ranganathan and Foster, 2016)). In this perspective,
720 nanoparticles could be specifically designed to target CAR T cells, making possible a
721 selective apoptosis of those cells or a selective removal of those cells. In this line, an
722 innovative strategy related to hematological diseases is the magnetic sorting of sick
723 cells, after attachment of a magnetic particle. In some cases, such as malaria, the
724 intrinsic magnetic properties of infected cells even allow magnetic sorting of
725 unlabeled cells (Zborowski and Chalmers, 2011). Nanoparticles targeting tumor cells
726 or CAR T cells could be used to lower the tumor burden (lymphodepletion) before
727 treatment or alternatively remove CAR T, after treatment or in case of excessive
728 expansion of CAR T cells.

729

730

731 **5 CONCLUSION**

732 Nanomedicine and cell therapy are two fields that have grown in parallel. Yet, those
733 approaches aim ultimately at common goals, to achieve long remission and ideally
734 the cure of the patients. In this review, based on the example of developing tools to
735 target B cell malignancy (mostly anti-CD19 nano-objects and anti-CD19 CAR T

cells), we have discussed their specificity, limitations and potential complementarity. It appears that even if CART T cell therapy has revolutionized management of patients presenting poor prognosis B cell malignancy, improvements are needed, especially to predict the therapeutic response, to control the intensity and persistence of the treatment, to increase tumor accessibility of the therapeutic agent to leukemic stem cell niches, and to visualize residual leukemic clones, and thus prevent relapses. Therefore, therapeutic developments could benefit from nanoparticles advantages -mainly their multimodality combining imaging and loading capacity, their tendency to accumulate at tumor sites for solid tumors and their relative easiness to be produced- to fill those requirements.

TABLES

Table 1: Main chemical and physical properties of the different types of nanoparticles used in nanomedicine and their principal applications. Note that the given size corresponds to the primary nano-object. In the case of small nanoparticles (NP) such as dendrimers or quantum dots (QD), surface modification with PEG or other macromolecules result in larger dimension.

NP type	Size (nm)	Organic/Inorganic	Principal application
Liposome	30-500	organic	encapsulation
Polymer NP	10-200	organic	encapsulation
Polymersome	50-1000	organic	encapsulation
Dendrimer	< 10	organic	encapsulation / imaging
Solid Lipid NP (and emulsion based particles)	> 100	organic	encapsulation
Silica NP	all range	inorganic	encapsulation / imaging
Quantum dot	5-20	inorganic	imaging
SPION	5-100	inorganic	imaging
Au NP	5-100	inorganic	imaging / therapy
Hybridosome®	80-120	organic/inorganic	imaging / encapsulation / therapy

Table 2: Nanoparticles (NP) grafted with anti-CD19 antibody and their applications in nanomedicine.

Abbreviation: Ag Silver; Au: Gold; Chol: Choline; DOTAP: 1,2-dioleoyl-3-trimethylammoniumpropane; DOPE: dioleoylphosphatidylethanolamine; DSPE: Distearoylphosphatidylethanolamine; EggPC: Egg yolk phosphatidylcholine; HD37-CCH: Hybridomas HD37-c-myc-Cys-His5 scFv; HSPC: hydrogenated soy phosphatidylcholine; LNP: liposomal nanoparticle ; MHC-Ig: Major Histocompatibility Complex-Immunoglobulin; NHS: N-hydroxysuccinimide; PEG: Polyethylene glycol; PLGA: poly(lactic-co-glycolic acid); SERS: Surface Enhanced Raman Scattering; SiO₂: Silicon dioxide ; SYK: Spleen Tyrosine Kinase; TCR: T cell receptor

NP type	Composition	Targeting agent	Size (nm)	Application	Reference
Liposome	PEG-DSPE	anti-CD19	100-120	doxorubicin carrier 140-160 µg/µmol of phospholipid	Lopes de Menezes et al., 1998
Liposome	HSPC/Chol/ mPEG-DSPE	anti-CD19	90-110	doxorubicin carrier	Sapra and Allen, 2004
Liposome	SM/Chol/ mPEG-DSPE	anti-CD19	110-130	vincristin carrier	Sapra and Allen, 2004
Liposome	mPEG ₂₀₀₀ -DSPE	anti-CD19 hd37-cch fragment	80-120	doxorubicin carrier	Cheng and Allen, 2008
Liposome	EggPC/Chol/ PEG ₂₀₀₀ -DSPE	anti-CD19 + anti-CD37 / anti-CD19 + anti-CD20 + anti-CD37	100	FTY720 carrier	Yu et al., 2013
Liposome	DSPE-PEG ₃₄₀₀ -NHS	mouse anti-CD19	~135	C61 carrier 9.4 mg/mL	Myers et al., 2014
Polymer NP	PEG-PLGA	anti-CD19 / anti-CD19 + anti-CD20	~300	hydroxychloroquine carrier 165 µg/mg of polymer	Mansilla et al., 2010
Polymer NP	EG ₁₁₃ -CL ₁₅₂ -TSU ₂₅	anti-CD19	~ 60	doxorubicin carrier 72.1±6.4 µg/mg of polymer	Krishnan et al., 2015
Inorganic	Au@PEG	human anti-CD19	60-80	SERS cell imaging MGITC = Raman tag	Nguyen et al., 2010
Inorganic	Ag@SiO ₂	anti-CD19	100-140	Fluorescence cell imaging	Dong et al., 2014
Inorganic	Iron oxide@dextran	pep-MHC-Ig dimer or anti-TCR-specific with anti-human CD19	~50	Targeting Redirect T cells against tumor cells	Schütz et al., 2016

FIGURE LEGENDS

Figure 1: B cell development and differentiation

B cell development begins in bone marrow and progresses through pre pro B cell, pro B cell, small pre B cell, large pre B cell and immature pre B cell. B cell locates within the circulatory system from mature B cell stage. The CD19 protein is expressed from pro B cell stage.

Figure 2: CD19 signaling complex and activation pathways

(A) Schematic representation of the CD19 signaling complex. The CD19 complex is composed of CD21, CD81 and CD19 transmembrane proteins. CD19 possesses an intracellular tail with multiple tyrosine-kinase residues involved in signal transduction.

(B) The first pathway of CD19 activation is dependent on the B cell receptor (BCR): it is a co-receptor for BCR signal transduction. The second pathway is independent of the BCR: the CD19 complex is able to bind activated complement fragment C3d and modulates BCR signaling (Figure adapted from (Wang et al., 2012)).

Figure 3: The two main modes of controlled release from carrier nanoparticles

Sustained release can be operated by biodegradable carriers, most often polymeric, which are progressively eroded, or by porous (silica, polymer...) particles. Trigger-activated particles deliver their load at once, upon activation by an endogenous or exogenous trigger.

Figure 4: Natural and engineered antibody formats, and functional groups available for covalent labeling or bioconjugation

(A) Schematic representation of full monoclonal antibody (mAb) of 150 kDa and its scFv derivative of 55 kDa. Functional groups present on the antibodies and available for covalent labeling or bioconjugation are schematically represented (amine groups, carboxylate groups, thiol groups and carbohydrate residues). Fab: variable region; Fc region: constant region; VL: Variable Light chain; VH: Variable Heavy chain; CL: Constant Light chain; CH: Constant Heavy chain.

(B) Comparison between mAb and its derivatives in terms of size, pharmacokinetics, valency/specificity and strengths/weaknesses.

Figure 5: Chimeric antigen receptor (CAR)

Chimeric antigen receptor (CAR) of second generation is composed of a targeting element (here the single chain variable fragment (scFv) of anti-CD19), a transmembrane domain, a co-stimulatory domain and a signaling domain.

LIST OF ABBREVIATIONS

ALL:	acute lymphoblastic leukemia
Ag:	Silver
Au:	Gold
BCR:	B cell receptor
B-ALL:	B cell acute lymphoblastic leukemia
CAR:	chimeric antigen receptor,
CL:	Constant Light chain;
CH:	Constant Heavy chain
Chol:	Choline
CLL :	chronic lymphocytic leukemia
DOTAP:	1,2-dioleoyl-3-trimethylammoniumpropane
DOPE:	dioleoylphosphatidylethanolamine
DSPE:	Distearoylphosphatidylethanolamine
EC :	European Commission
EDC:	(1-ethyl-3-(3- dimethyl-aminopropyl)carbodiimide hydrochloride
EggPC:	Egg yolk phosphatidylcholine
EPR :	Enhanced Permeation and Retention
Fab:	variable region
Fc region:	constant region
FDA :	US Food-and-Drug-Administration
IFN γ :	interferon gamma
IL6:	interleukin 6
HD37-CCH:	Hybridomas HD37-c-myc-Cys-His5 scFv
HSPC:	hydrogenated soy phosphatidylcholine
LNP:	liposomal nanoparticle
mAb:	monoclonal antibody
MGITC:	Malachite Green Isothiocyanate
MHC-Ig:	Major Histocompatibility Complex-Immunoglobulin
MPS:	mononuclear phagocyte system
MRI :	Magnetic Resonance Imaging
MRI/CT :	magnetic resonance imaging/ computerized tomography
MRI/PET :	magnetic resonance imaging/ positron emission tomography
NHS:	N-hydroxysuccinimide
NP:	nanoparticle
PEG:	Polyethylene glycol
PLGA:	poly(lactic-co-glycolic acid)
PVP:	polyvinylpyrrolidone
QD:	quantum dots
RES :	reticuloendothelial system
SERS:	Surface Enhanced Raman Scattering
SiO ₂ :	Silicon dioxide
siRNA:	small interference RNA
SMCC:	N-succinimidyl 4-(N maleimidomethyl)cyclohexane-1-carboxylate
SPDP:	N-succinimidyl 3-(2-pyridylthio)propionate
SPECT:	single, photon emission computed tomography
scFv:	single-chain variable fragment
SYK:	Spleen Tyrosine Kinase
TCR:	T cell receptor

858 TEM: transmission electron microscopy
859 UCNPs : up-converting nanoparticles
860 VL: Variable Light chain
861 VH: Variable Heavy chain
862

ACKNOWLEDGEMENTS/FUNDING

This work is supported by SFR Biosit UMS CNRS 3480 - INSERM 018 (Call Interdisciplinary Project, SC, FG, MBT), Université de Rennes 1 (Call Scientific Challenge, SC, FG, MBT), French Ministry of Research (HJ, FS), the Sociétés d'Accélération du Transfert de Technologies Ouest Valorisation (SC, FG, FS), Ligue régionale contre le cancer (comity 22, 35, 56, 79, 41) (MBT), the Société française de lutte contre les cancers et les leucémies de l'enfant et de l'adolescent and the Fédération Enfants et Santé (MBT), a private donator Mrs. M-Dominique Blanc (MBT), the CNRS (MBT, FG), and the People Programme (Marie Curie Actions) of the European Union's Seventh Framework Programme (FP7/2007-2013) under REA grant agreement n°291851 (MBT).

BIBLIOGRAPHY

- Beatty, G.L., and O'Hara, M. (2016). Chimeric antigen receptor-modified T cells for the treatment of solid tumors: Defining the challenges and next steps. *Pharmacol. Ther.* 166, 30–39.
- Bhatnagar, P., Li, Z., Choi, Y., Guo, J., Li, F., Lee, D.Y., Figliola, M., Huls, H., Lee, D.A., Zal, T., et al. (2013). Imaging of Genetically Engineered T Cells by PET using Gold Nanoparticle Complexed to Copper-64. *Integr. Biol. Quant. Biosci. Nano Macro* 5, 231.
- Bhatnagar, P., Alauddin, M., Bankson, J.A., Kirui, D., Seifi, P., Huls, H., Lee, D.A., Babakhani, A., Ferrari, M., Li, K.C., et al. (2014). Tumor Lysing Genetically Engineered T Cells Loaded with Multi-Modal Imaging Agents. *Sci. Rep.* 4.
- Bhattacharya, S., Ganivada, M.N., Dinda, H., Das Sarma, J., and Shunmugam, R. (2016). Biodegradable Copolymer for Stimuli-Responsive Sustained Release of Doxorubicin. *ACS Omega* 1, 108–117.
- Braig, F., Brandt, A., Goebeler, M., Tony, H.-P., Kurze, A.-K., Nollau, P., Bumm, T., Böttcher, S., Bargou, R.C., and Binder, M. (2017). Resistance to anti-CD19/CD3 BiTE in acute lymphoblastic leukemia may be mediated by disrupted CD19 membrane trafficking. *Blood* 129, 100–104.
- Brentjens, R., Davila, M.L., Riviere, I., Park, J., Wang, X., Cowell, L.G., Bartido, S., Stefanski, J., Taylor, C., Olszewska, M., et al. (2013). CD19-targeted T cells rapidly induce molecular remissions in adults with chemotherapy-refractory acute lymphoblastic leukemia. *Sci. Transl. Med.* 5, 177ra38.
- Brudno, J.N., Somerville, R.P.T., Shi, V., Rose, J.J., Halverson, D.C., Fowler, D.H., Gea-Banacloche, J.C., Pavletic, S.Z., Hickstein, D.D., Lu, T.L., et al. (2016). Allogeneic T Cells That Express an Anti-CD19 Chimeric Antigen Receptor Induce Remissions of B-Cell Malignancies That Progress After Allogeneic Hematopoietic Stem-Cell Transplantation Without Causing Graft-Versus-Host Disease. *J. Clin. Oncol.* 34, 1112.
- Carter, P.J. (2006). Potent antibody therapeutics by design. *Nat. Rev. Immunol.* 6, 343–357.
- Chang, S., Warner, J., Liang, L., and Fairman, J. (2009). A novel vaccine adjuvant for recombinant flu antigens. *Biol. J. Int. Assoc. Biol. Stand.* 37, 141–147.
- Chen, G., Roy, I., Yang, C., and Prasad, P.N. (2016). Nanochemistry and nanomedicine for nanoparticle-based diagnostics and therapy. *Chem. Rev.* 116, 2826–2885.
- Cheng, W.W.K., and Allen, T.M. (2008). Targeted delivery of anti-CD19 liposomal doxorubicin in B-cell lymphoma: A comparison of whole monoclonal antibody, Fab' fragments and single chain Fv. *J. Controlled Release* 126, 50–58.
- Cistaro, A., Delfa, V.L., Rosa, G.D., Cogoni, M., and Quartuccio, N. (2017). MRI and 18F-FDG-PET/CT in a rare case of early (precursor) B-lymphoblastic leukaemia with bone involvement as initial manifestation. *Nucl. Med. Rev.* 20, 57–59.
- Davenport, A.J., Jenkins, M.R., Cross, R.S., Yong, C.S., Prince, H.M., Ritchie, D.S., Trapani, J.A., Kershaw, M.H., Darcy, P.K., and Neeson, P.J. (2015). CAR-T Cells Inflict Sequential Killing of Multiple Tumor Target Cells. *Cancer Immunol. Res.* 3, 483–494.
- Davila, M.L., Riviere, I., Wang, X., Bartido, S., Park, J., Curran, K., Chung, S.S., Stefanski, J., Borquez-Ojeda, O., Olszewska, M., et al. (2014). Efficacy and Toxicity Management of 19-28z CAR T Cell Therapy in B Cell Acute Lymphoblastic Leukemia. *Sci. Transl. Med.* 6, 224ra25–224ra25.

923 Dong, M., Tian, Y., and Pappas, D. (2014). Facile functionalization of Ag@SiO₂ core-shell
 924 metal enhanced fluorescence nanoparticles for cell labeling. *Anal. Methods* 6, 1598–
 925 1602.
 926 Eyquem, J., Mansilla-Soto, J., Giavridis, T., van der Stegen, S.J.C., Hamieh, M., Cunanan,
 927 K.M., Odak, A., Gönen, M., and Sadelain, M. (2017). Targeting a CAR to the TRAC locus
 928 with CRISPR/Cas9 enhances tumour rejection. *Nature* 543, 113–117.
 929 Frey, N.V., and Porter, D.L. (2016). The Promise of Chimeric Antigen Receptor T-Cell
 930 Therapy | Cancer Network.
 931 Gattazzo, F., Urciuolo, A., and Bonaldo, P. (2014). Extracellular matrix: A dynamic
 932 microenvironment for stem cell niche. *Biochim. Biophys. Acta* 1840, 2506.
 933 Gerber, H.-P., Kung-Sutherland, M., Stone, I., Morris-Tilden, C., Miyamoto, J., McCormick,
 934 R., Alley, S.C., Okeley, N., Hayes, B., Hernandez-Ilizaliturri, F.J., et al. (2009). Potent
 935 antitumor activity of the anti-CD19 auristatin antibody drug conjugate hBU12-vcMMAE
 936 against rituximab-sensitive and -resistant lymphomas. *Blood* 113, 4352–4361.
 937 Geyer, M.B., and Brentjens, R.J. (2016). Review: Current clinical applications of chimeric
 938 antigen receptor (CAR) modified T cells. *Cytotherapy* 18, 1393–1409.
 939 Gill, S., and June, C.H. (2015). Going viral: chimeric antigen receptor T-cell therapy for
 940 hematological malignancies. *Immunol. Rev.* 263, 68–89.
 941 Goebeler, M.-E., and Bargou, R. (2016). Blinatumomab: a CD19/CD3 bispecific T cell
 942 engager (BiTE) with unique anti-tumor efficacy. *Leuk. Lymphoma*.
 943 Gormley, A.J., Larson, N., Banisadr, A., Robinson, R., Frazier, N., Ray, A., and Ghandehari,
 944 H. (2013). Plasmonic photothermal therapy increases the tumor mass penetration of
 945 HPMA copolymers. *J. Controlled Release* 166, 130–138.
 946 Greish, K. (2007). Enhanced permeability and retention of macromolecular drugs in
 947 solid tumors: a royal gate for targeted anticancer nanomedicines. *J. Drug Target.* 15,
 948 457–464.
 949 Grupp, S.A., Kalos, M., Barrett, D., Aplenc, R., Porter, D.L., Rheingold, S.R., Teachey, D.T.,
 950 Chew, A., Hauck, B., Wright, J.F., et al. (2013). Chimeric Antigen Receptor–Modified T
 951 Cells for Acute Lymphoid Leukemia.
 952 Hammer, O. (2012). CD19 as an attractive target for antibody-based therapy. *MAbs* 4,
 953 571–577.
 954 Herhaus, P., Habringer, S., Philipp-Abbrederis, K., Vag, T., Gerngross, C., Schottelius, M.,
 955 Slotta-Huspenina, J., Steiger, K., Altmann, T., Weißer, T., et al. (2016). Targeted positron
 956 emission tomography imaging of CXCR4 expression in patients with acute myeloid
 957 leukemia. *Haematologica* 101, 932–940.
 958 Hong, E.E., Erickson, H., Lutz, R.J., Whiteman, K.R., Jones, G., Kovtun, Y., Blanc, V., and
 959 Lambert, J.M. (2015). Design of Coltuximab Ravtansine, a CD19-Targeting Antibody–
 960 Drug Conjugate (ADC) for the Treatment of B-Cell Malignancies: Structure–Activity
 961 Relationships and Preclinical Evaluation.
 962 Hudecek, M., Lupo-Stanghellini, M.-T., Kosasih, P.L., Sommermeyer, D., Jensen, M.C.,
 963 Rader, C., and Riddell, S.R. (2013). Receptor Affinity and Extracellular Domain
 964 Modifications Affect Tumor Recognition by ROR1-Specific Chimeric Antigen Receptor T
 965 Cells. *Clin. Cancer Res.* 19, 3153–3164.
 966 Jin, L., Hope, K.J., Zhai, Q., Smadja-Joffe, F., and Dick, J.E. (2006). Targeting of CD44
 967 eradicates human acute myeloid leukemic stem cells. *Nat. Med.* 12, 1167–1174.
 968 Kamaly, N., Yameen, B., Wu, J., and Farokhzad, O.C. (2016). Degradable controlled-
 969 release polymers and polymeric nanoparticles: mechanisms of controlling drug release.
 970 *Chem. Rev.* 116, 2602–2663.
 971 Kennedy, L.C., Bear, A.S., Young, J.K., Lewinski, N.A., Kim, J., Foster, A.E., and Drezek, R.A.

972 (2011). T cells enhance gold nanoparticle delivery to tumors in vivo. *Nanoscale Res. Lett.*
 973 6, 283.
 974 Kochenderfer, J.N., Dudley, M.E., Feldman, S.A., Wilson, W.H., Spaner, D.E., Maric, I.,
 975 Stetler-Stevenson, M., Phan, G.Q., Hughes, M.S., Sherry, R.M., et al. (2012). B-cell
 976 depletion and remissions of malignancy along with cytokine-associated toxicity in a
 977 clinical trial of anti-CD19 chimeric-antigen-receptor-transduced T cells. *Blood* 119,
 978 2709–2720.
 979 Krause, D.S., and Scadden, D.T. (2015). A hostel for the hostile: the bone marrow niche in
 980 hematologic neoplasms. *Haematologica* 100, 1376–1387.
 981 Krause, D.S., Lazarides, K., Andrian, U.H. von, and Etten, R.A.V. (2006). Requirement for
 982 CD44 in homing and engraftment of BCR-ABL[ndash]expressing leukemic stem cells.
 983 *Nat. Med.* 12, 1175–1180.
 984 Krause, D.S., Fulzele, K., Catic, A., Sun, C.C., Dombkowski, D., Hurley, M.P., Lezeau, S.,
 985 Attar, E., Wu, J.Y., Lin, H.Y., et al. (2013). Differential regulation of myeloid leukemias by
 986 the bone marrow microenvironment. *Nat. Med.* 19, 1513–1517.
 987 Krishnan, V., Xu, X., Kelly, D., Snook, A., Waldman, S.A., Mason, R.W., Jia, X., and
 988 Rajasekaran, A.K. (2015). CD19-Targeted Nanodelivery of Doxorubicin Enhances
 989 Therapeutic Efficacy in B-Cell Acute Lymphoblastic Leukemia. *Mol. Pharm.* 12, 2101–
 990 2111.
 991 Kuhn, S.J., Finch, S.K., Hallahan, D.E., and Giorgio, T.D. (2006). Proteolytic Surface
 992 Functionalization Enhances *in Vitro* Magnetic Nanoparticle Mobility through
 993 Extracellular Matrix. *Nano Lett.* 6, 306–312.
 994 LeBien, T.W., and Tedder, T.F. (2008). B lymphocytes: how they develop and function.
 995 *Blood* 112, 1570–1580.
 996 Lee, D.W., Gardner, R., Porter, D.L., Louis, C.U., Ahmed, N., Jensen, M., Grupp, S.A., and
 997 Mackall, C.L. (2014). Current concepts in the diagnosis and management of cytokine
 998 release syndrome. *Blood* 124, 188–195.
 999 Lee, D.W., Kochenderfer, J.N., Stetler-Stevenson, M., Cui, Y.K., Delbrook, C., Feldman, S.A.,
 1000 Fry, T.J., Orentas, R., Sabatino, M., Shah, N.N., et al. (2015). T cells expressing CD19
 1001 chimeric antigen receptors for acute lymphoblastic leukaemia in children and young
 1002 adults: a phase 1 dose-escalation trial. *The Lancet* 385, 517–528.
 1003 Li, X., Ding, Y., Zi, M., Sun, L., Zhang, W., Chen, S., and Xu, Y. (2017). CD19, from bench to
 1004 bedside. *Immunol. Lett.* 183, 86–95.
 1005 Liu, Z., and Li, Z. (2014). Molecular Imaging in Tracking Tumor-Specific Cytotoxic T
 1006 Lymphocytes (CTLs). *Theranostics* 4, 990–1001.
 1007 Lopes de Menezes, D.E., Pilarski, L.M., and Allen, T.M. (1998). In vitro and in vivo
 1008 targeting of immunoliposomal doxorubicin to human B-cell lymphoma. *Cancer Res.* 58,
 1009 3320–3330.
 1010 Ma, D., McDevitt, M.R., Barendswaard, E., Lai, L., Curcio, M.J., Pellegrini, V., Brechbiel,
 1011 M.W., and Scheinberg, D.A. (2002). Radioimmunotherapy for model B cell malignancies
 1012 using 90Y-labeled anti-CD19 and anti-CD20 monoclonal antibodies. *Leukemia* 16, 60–66.
 1013 Maeda, H., Wu, J., Sawa, T., Matsumura, Y., and Hori, K. (2000). Tumor vascular
 1014 permeability and the EPR effect in macromolecular therapeutics: a review. *J. Control.*
 1015 *Release Off. J. Control. Release Soc.* 65, 271–284.
 1016 Manjappa, A.S., Chaudhari, K.R., Venkataraju, M.P., Dantuluri, P., Nanda, B., Sidda, C.,
 1017 Sawant, K.K., and Ramachandra Murthy, R.S. (2011). Antibody derivatization and
 1018 conjugation strategies: Application in preparation of stealth immunoliposome to target
 1019 chemotherapeutics to tumor. *J. Controlled Release* 150, 2–22.
 1020 Mansilla, E., Marin, G.H., Nuñez, L., Drago, H., Sturla, F., Mertz, C., Rivera, L., Ichim, T.,

Riordan, N., and Raimondi, C. (2010). The Lysosomotropic Agent, Hydroxychloroquine, Delivered in a Biodegradable Nanoparticle System, Overcomes Drug Resistance of B-Chronic Lymphocytic Leukemia Cells In Vitro. *Cancer Biother. Radiopharm.* 25, 97–103.
 Martucci, N.M., Migliaccio, N., Ruggiero, I., Albano, F., Calì, G., Romano, S., Terracciano, M., Rea, I., Arcari, P., and Lamberti, A. (2016). Nanoparticle-based strategy for personalized B-cell lymphoma therapy. *Int. J. Nanomedicine* 11, 6089.
 Maude, S.L., Frey, N., Shaw, P.A., Aplenc, R., Barrett, D.M., Bunin, N.J., Chew, A., Gonzalez, V.E., Zheng, Z., Lacey, S.F., et al. (2014). Chimeric Antigen Receptor T Cells for Sustained Remissions in Leukemia. *N. Engl. J. Med.* 371, 1507–1517.
 Moghimi, S., Hunter, A., and Murray, J. (2001). Long-Circulating and Target-Specific Nanoparticles: Theory to Practice. *Pharmacol. Rev.* 53, 283–318.
 Morrison, S.J., and Scadden, D.T. (2014). The bone marrow niche for haematopoietic stem cells. *Nature* 505, 327–334.
 Myers, D.E., Yiv, S., Qazi, S., Ma, H., Cely, I., Shahidzadeh, A., Arellano, M., Finestone, E., Gaynon, P.S., Termuhlen, A., et al. (2014). CD19-antigen specific nanoscale liposomal formulation of a SYK P-site inhibitor causes apoptotic destruction of human B-precursor leukemia cells. *Integr. Biol.* 6, 766.
 Navarro, S.M., Matcuk, G.R., Patel, D.B., Skalski, M., White, E.A., Tomasian, A., and Schein, A.J. (2017). Musculoskeletal Imaging Findings of Hematologic Malignancies. *Radiogr. Rev. Publ. Radiol. Soc. N. Am. Inc* 37, 881–900.
 Newick, K., Moon, E., and Albelda, S.M. (2016). Chimeric antigen receptor T-cell therapy for solid tumors. *Mol. Ther. Oncolytics* 3, 16006.
 Nguyen, C.T., Nguyen, J.T., Rutledge, S., Zhang, J., Wang, C., and Walker, G.C. (2010). Detection of chronic lymphocytic leukemia cell surface markers using surface enhanced Raman scattering gold nanoparticles. *Cancer Lett.* 292, 91–97.
 Park, J.H., Geyer, M.B., and Brentjens, R.J. (2016). CD19-targeted CAR T-cell therapeutics for hematologic malignancies: interpreting clinical outcomes to date. *Blood* 127, 3312–3320.
 Parodi, A., Haddix, S.G., Taghipour, N., Scaria, S., Taraballi, F., Cevenini, A., Yazdi, I.K., Corbo, C., Palomba, R., Khaled, S.Z., et al. (2014). Bromelain Surface Modification Increases the Diffusion of Silica Nanoparticles in the Tumor Extracellular Matrix. *ACS Nano* 8, 9874–9883.
 Pattni, B.S., Chupin, V.V., and Torchilin, V.P. (2015). New Developments in Liposomal Drug Delivery. *Chem. Rev.* 115, 10938–10966.
 Perica, K., Medero, A.D.L., Durai, M., Chiu, Y.L., Bieler, J.G., Sibener, L., Niemöller, M., Assenmacher, M., Richter, A., Edidin, M., et al. (2014). Nanoscale Artificial Antigen Presenting Cells for T Cell Immunotherapy. *Nanomedicine Nanotechnol. Biol. Med.* 10, 119.
 Porter, D.L., Levine, B.L., Kalos, M., Bagg, A., and June, C.H. (2011). Chimeric Antigen Receptor–Modified T Cells in Chronic Lymphoid Leukemia.
 Procko, E., Berguig, G.Y., Shen, B.W., Song, Y., Frayo, S., Convertine, A.J., Margineantu, D., Booth, G., Correia, B.E., Cheng, Y., et al. (2014). A computationally designed inhibitor of an Epstein-Barr viral Bcl-2 protein induces apoptosis in infected cells. *Cell* 157, 1644.
 Ramachandran, M., Dimberg, A., and Essand, M. (2017). The cancer-immunity cycle as rational design for synthetic cancer drugs: Novel DC vaccines and CAR T-cells. *Semin. Cancer Biol.*
 Ranganathan, R., and Foster, M.C. (2016). The Limitations and Promise of Immunotherapy With Chimeric Antigen–Modified T Cells | Cancer Network.
 Riley, R.S., and Day, E.S. (2017). Gold nanoparticle-mediated photothermal therapy:

applications and opportunities for multimodal cancer treatment. Wiley Interdiscip. Rev. Nanomed. Nanobiotechnol. n/a-n/a.

Rodgers, D.T., Mazagova, M., Hampton, E.N., Cao, Y., Ramadoss, N.S., Hardy, I.R., Schulman, A., Du, J., Wang, F., Singer, O., et al. (2016). Switch-mediated activation and retargeting of CAR-T cells for B-cell malignancies. *Proc. Natl. Acad. Sci.* *113*, E459–E468.

Ruella, M., Kenderian, S.S., Shestova, O., Klichinsky, M., Melenhorst, J.J., Wasik, M.A., Lacey, S.F., June, C.H., and Gill, S. (2017). Kinase inhibitor ibrutinib to prevent cytokine-release syndrome after anti-CD19 chimeric antigen receptor T cells for B-cell neoplasms. *Leukemia* *31*, 246–248.

Salem, U., Menias, C.O., Shaaban, A., Bhosale, P.R., Youssef, A., and Elsayes, K.M. (2014). Hematopoietic tumors of the female genital system: imaging features with pathologic correlation. *Abdom. Imaging* *39*, 922–934.

Sapra, P., and Allen, T.M. (2004). Improved outcome when B-cell lymphoma is treated with combinations of immunoliposomal anticancer drugs targeted to both the CD19 and CD20 epitopes. *Clin. Cancer Res. Off. J. Am. Assoc. Cancer Res.* *10*, 2530–2537.

Savoldo, B., Ramos, C.A., Liu, E., Mims, M.P., Keating, M.J., Carrum, G., Kamble, R.T., Bollard, C.M., Gee, A.P., Mei, Z., et al. (2011). CD28 costimulation improves expansion and persistence of chimeric antigen receptor–modified T cells in lymphoma patients. *J. Clin. Invest.* *121*, 1822.

Scheinberg, D.A., and Strand, M. (1983). Kinetic and Catabolic Considerations of Monoclonal Antibody Targeting in Erythroleukemic Mice. *Cancer Res.* *43*, 265–272.

Schepers, K., Campbell, T.B., and Passegué, E. (2015). Normal and Leukemic Stem Cell Niches: Insights and Therapeutic Opportunities. *Cell Stem Cell* *16*, 254.

Schütz, C., Varela, J.C., Perica, K., Haupt, C., Oelke, M., and Schneck, J.P. (2016). Antigen-specific T cell Redirectors: a nanoparticle based approach for redirecting T cells. *Oncotarget*.

Sciortino, F., Casterou, G., Eliat, P.-A., Troadec, M.-B., Gaillard, C., Chevance, S., Kahn, M.L., and Gauffre, F. (2016a). Simple Engineering of Polymer–Nanoparticle Hybrid Nanocapsules. *ChemNanoMat* *2*, 796–799.

Sciortino, F., Casterou, G., Eliat, P.-A., Troadec, M.-B., Gaillard, C., Chevance, S., Kahn, M.L., and Gauffre, F. (2016b). Simple Engineering of Polymer–Nanoparticle Hybrid Nanocapsules. *ChemNanoMat* *2*, 796–799.

Shi, J., Kantoff, P.W., Wooster, R., and Farokhzad, O.C. (2017). Cancer nanomedicine: progress, challenges and opportunities. *Nat. Rev. Cancer* *17*, 20–37.

Sipkins, D.A., Wei, X., Wu, J.W., Runnels, J.M., Côté, D., Means, T.K., Luster, A.D., Scadden, D.T., and Lin, C.P. (2005). In vivo imaging of specialized bone marrow endothelial microdomains for tumour engraftment. *Nature* *435*, 969–973.

Smith, B.E., Roder, P.B., Zhou, X., and Pauzauskie, P.J. (2015). Nanoscale materials for hyperthermal theranostics. *Nanoscale* *7*, 7115–7126.

Sommermeier, D., Hudecek, M., Kosasih, P.L., Gogishvili, T., Maloney, D.G., Turtle, C.J., and Riddell, S.R. (2016). Chimeric antigen receptor-modified T cells derived from defined CD8+ and CD4+ subsets confer superior antitumor reactivity in vivo. *Leukemia* *30*, 492–500.

Stamenkovic, I., and Seed, B. (1988). CD19, the earliest differentiation antigen of the B cell lineage, bears three extracellular immunoglobulin-like domains and an Epstein-Barr virus-related cytoplasmic tail. *J. Exp. Med.* *168*, 1205–1210.

Stephenson, R., and Singh, A. (2017). Drug discovery and therapeutic delivery for the treatment of B and T cell tumors. *Adv. Drug Deliv. Rev.*

Tedder, T.F., Zhou, L.J., and Engel, P. (1994). The CD19/CD21 signal transduction

complex of B lymphocytes. *Immunol. Today* *15*, 437–442.

Turtle, C.J., Hanafi, L.-A., Berger, C., Gooley, T.A., Cherian, S., Hudecek, M., Sommermeyer, D., Melville, K., Pender, B., Budiarto, T.M., et al. (2016). CD19 CAR-T cells of defined CD4+:CD8+ composition in adult B cell ALL patients. *J. Clin. Invest.* *126*, 2123–2138.

Uckun, F.M., Jaszcz, W., Ambrus, J.L., Fauci, A.S., Gajl-Peczalska, K., Song, C.W., Wick, M.R., Myers, D.E., Waddick, K., and Ledbetter, J.A. (1988a). Detailed studies on expression and function of CD19 surface determinant by using B43 monoclonal antibody and the clinical potential of anti-CD19 immunotoxins. *Blood* *71*, 13–29.

Uckun, F.M., Jaszcz, W., Ambrus, J.L., Fauci, A.S., Gajl-Peczalska, K., Song, C.W., Wick, M.R., Myers, D.E., Waddick, K., and Ledbetter, J.A. (1988b). Detailed studies on expression and function of CD19 surface determinant by using B43 monoclonal antibody and the clinical potential of anti-CD19 immunotoxins. *Blood* *71*, 13–29.

Uckun, F.M., Qazi, S., and Cheng, J. (2015a). Targeting leukemic stem cells with multifunctional bioactive polypeptide nanoparticles. *Future Oncol.* *11*, 1149–1152.

Uckun, F.M., Ma, H., Cheng, J., Myers, D.E., and Qazi, S. (2015b). CD22ΔE12 as a molecular target for RNAi therapy. *Br. J. Haematol.* *169*, 401–414.

Velasquez, M.P., and Gottschalk, S. (2017). Targeting CD19: the good, the bad, and CD81. *Blood* *129*, 9–10.

Walker, G.C., Maclaughlin, C.M., and Ip, S. (2012). Lipid Encapsulation of Surface Enhanced Raman Scattering (sers) Nanoparticles.

Wang, K., Wei, G., and Liu, D. (2012). CD19: a biomarker for B cell development, lymphoma diagnosis and therapy. *Exp. Hematol. Oncol.* *1*, 1.

Wang, Z., Wu, Z., Liu, Y., and Han, W. (2017). New development in CAR-T cell therapy. *J. Hematol. Oncol.* *J Hematol Oncol* *10*.

Wilhelm, S., Tavares, A.J., Dai, Q., Ohta, S., Audet, J., Dvorak, H.F., and Chan, W.C.W. (2016). Analysis of nanoparticle delivery to tumours. *Nat. Rev. Mater.* *1*, 16014.

Wong, J.K.L., Mohseni, R., Hamidieh, A.A., MacLaren, R.E., Habib, N., and Seifalian, A.M. (2017). Will Nanotechnology Bring New Hope for Gene Delivery? *Trends Biotechnol.*

Woyach, J.A., Awan, F., Flinn, I.W., Berdeja, J.G., Wiley, E., Mansoor, S., Huang, Y., Lozanski, G., Foster, P.A., and Byrd, J.C. (2014). A phase 1 trial of the Fc-engineered CD19 antibody XmAb5574 (MOR00208) demonstrates safety and preliminary efficacy in relapsed CLL. *Blood* *124*, 3553.

Yan, J., Wolff, M.J., Unternaehrer, J., Mellman, I., and Mamula, M.J. (2005). Targeting antigen to CD19 on B cells efficiently activates T cells. *Int. Immunol.* *17*, 869–877.

Yin, H., Liao, L., and Fang, J. (2014). Enhanced Permeability and Retention (EPR) Effect Based Tumor Targeting: The Concept, Application and Prospect. *2*, 1010.

Yu, B., Mao, Y., Yuan, Y., Yue, C., Wang, X., Mo, X., Jarjoura, D., Paulaitis, M.E., Lee, R.J., Byrd, J.C., et al. (2013). Targeted drug delivery and cross-linking induced apoptosis with anti-CD37 based dual-ligand immunoliposomes in B chronic lymphocytic leukemia cells. *Biomaterials* *34*, 6185–6193.

Zah, E., Lin, M.-Y., Silva-Benedict, A., Jensen, M.C., and Chen, Y.Y. (2016). T Cells Expressing CD19/CD20 Bispecific Chimeric Antigen Receptors Prevent Antigen Escape by Malignant B Cells. *Cancer Immunol. Res.* *4*, 498–508.

Zborowski, M., and Chalmers, J.J. (2011). Rare Cell Separation and Analysis by Magnetic Sorting. *Anal. Chem.* *83*, 8050–8056.

Zhu, J., and Emerson, S.G. (2002). Hematopoietic cytokines, transcription factors and lineage commitment.

FIGURE 1

A CD19 signaling complex

B CD19 activation pathways

FIGURE 2

FIGURE 3

A

B

Name	Size	Pharmacokinetics	Valency/ specificity	Strengths & Weaknesses
mAb	~ 150 kDa	long systemic clearance <i>half-time</i> ~ few days to weeks	Monospecific <i>bivalent</i>	High specificity/sensitivity towards their target Costly to produce Highly immunogenic
FAb	55 kDa	short systemic clearance <i>half-time</i> ~ 10 h	Monospecific <i>monovalent</i>	Rapid tumor targeting Less immunogenic than mAb Improved tumor penetration compared to mAb Low avidity (monovalency) Renal toxicity (high renal uptake)
F(ab)₂	110 kDa	short systemic clearance <i>half-time</i> ~ 10 h	Bispecific <i>bivalent</i>	Approved by FDA Better avidity for the target than mAb Renal toxicity (high renal uptake)
F(ab)₃	165 kDa	very short systemic clearance <i>half-time</i> ~ 4-5 h	Trispecific <i>trivalent</i>	High avidity for the target (multivalency) Improved tumor penetration compared to the mAb Renal toxicity (high uptake)
scFv	28 kDa	ultra-short systemic clearance <i>half-time</i> > 1 h	Monospecific <i>monovalent</i>	Improved tumor penetration compared to mAb Less immunogenic than full mAb Low functional avidity (monovalency) Renal and hepatic toxicity (high uptake)
Minibody	75-105 kDa	intermediate systemic clearance	Monospecific <i>bivalent</i>	Faster tumor addressing Better therapeutic efficacy than mAb High tumor uptake Renal toxicity (high uptake)
Multimers of scFv			Monospecific	High avidity for the target (multivalency)
Dibody	50 kDa	intermediate systemic clearance	<i>bivalent</i>	Higher tumor uptake than mAb
Tribody	75 kDa		<i>trivalent</i>	Poor tumor penetration
Tetrbody	100 kDa		<i>tetravalent</i>	Renal toxicity (high uptake)

FIGURE 4

Anti-CD19 CAR second generation

FIGURE 5