Studies on chill coma recovery in the ladybird, *Harmonia axyridis* Ontogenetic profile, effect of repeated cold exposures, and capacity to predict winter survival

Michal Knapp, Philippe Vernon, D Renault

► To cite this version:

Michal Knapp, Philippe Vernon, D Renault. Studies on chill coma recovery in the ladybird, *Harmonia axyridis* Ontogenetic profile, effect of repeated cold exposures, and capacity to predict winter survival. *Journal of Thermal Biology*, 2018, 74, pp.275-280. 10.1016/j.jtherbio.2018.04.013 . hal-01807843

HAL Id: hal-01807843

<https://hal-univ-rennes1.archives-ouvertes.fr/hal-01807843>

Submitted on 28 Aug 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Studies on chill coma recovery in the ladybird, *Harmonia axyridis*: ontogenetic profile,
effect of repeated cold exposures, and capacity to predict winter survival**

MICHAL KNAPP^{1*}, PHILIPPE VERNON², DAVID RENAULT^{3,4}

¹ Department of Ecology, Faculty of Environmental Sciences, Czech University of Life
Sciences Prague, Kamýcká 129, Prague - Suchbát, 165 00, Czech Republic

² University of Rennes, CNRS, ECOBIO, UMR 6553, Station Biologique de Paimpont, 35380
Paimpont, France

³ University of Rennes 1, UMR 6553 EcoBio CNRS, 263 Avenue du Gal Leclerc, 35000
Rennes Cedex, France

⁴ Institut Universitaire de France, 1 Rue Descartes, 75231 Paris Cedex 05

* Correspondence:

Michal Knapp, Department of Ecology, Faculty of Environmental Sciences, Czech University
of Life Sciences Prague, Kamýcká 129, Praha – Suchbát, 165 00, Czech Republic. Tel. +420
22438 3853 e-mail: knapp@fzp.czu.cz

21 **Abstract**

22 The harlequin ladybird, *Harmonia axyridis*, is one of the most successful invasive insect
23 species worldwide. We investigated whether (i) chill coma recovery time (CCRt) changes
24 during the ontogenetic development of this species, (ii) CCRt varies in response to repeated
25 cold shocks, and (iii) CCRt could be a good predictor of winter survival ability in adults.
26 CCRt decreased during larval development, the lowest CCRt values were observed in teneral
27 adults (one and four days old), and significantly higher values were observed for older adults
28 (16 and 32 days old). Repeated cold shocks (two hours at -3 °C), interrupted by short (30
29 minutes) warm periods (22 °C) resulted in decreased CCRt after the second cold shock,
30 probably depicting an acclimation response, but then CCRt increased with additional cold
31 shocks, likely revealing the accumulation of chill injuries. The CCRt of pre-overwintering
32 individuals was not correlated with their winter survival. This indicates that CCRt is not a
33 reliable measure of cold tolerance in *H. axyridis*. However, this result could be partially
34 affected by the experimental setup – the use of laboratory-reared individuals who experienced
35 standardized conditions and thus the variability in CCRt of tested animals could be much
36 lower than the variability present in nature. The substantial variation observed over the
37 ontogenetic development of *H. axyridis* poses important methodological implications for
38 future studies, as animals of the same stage/age should be compared with each other. The
39 observed U-shaped response to repeated cold shocks indicates that the expectation of linearity
40 between cold exposure and insect response is an oversimplification of real situations.

41 **Highlights:**

- 42 - chill coma recovery time (CCRt) varies significantly during ladybird ontogeny
- 43 - repeated cold shocks result in a U-shaped CCRt response
- 44 - pre-overwintering CCRt was not related to winter survival in this species
- 45 - the relevance of CCRt as a measure of cold tolerance needs to be evaluated in other
- 46 species

47

48 **Key Words:**

49 Ageing, chill injury, cold tolerance, invasive species, ontogenetic stage, overwintering

1. Introduction

Because of their ectothermic nature and small body sizes, insects are particularly prone to rapid changes in their body temperature and have had to develop a wealth of behavioural and physiological responses to survive at low temperatures (Teets and Denlinger, 2013). As a result, thermal tolerance is a crucial physiological trait that is significantly related to the distribution of insect species on Earth (Araujo et al., 2013). A range of techniques are available for assessing the cold tolerance of insects; while some of them directly measure mortality (e.g., estimation of survival capacities, measurements of lower lethal limits), other techniques are only proxies of insect tolerance to the deleterious effects of low temperatures (e.g., supercooling point, chill coma onset and recovery; see Terblanche et al., 2011 and Sinclair et al., 2015 for a review).

Chill coma onset and recovery have been described as reliable metrics of thermal plasticity and tolerance in insects (Sinclair et al., 2015). Chill coma is initiated when neuromuscular function is disrupted as a result of the organism's inability to maintain proper ion balance due to depolarization of the resting membrane potential (MacMillan and Sinclair, 2011; Overgaard and MacMillan, 2017). Entering into chill coma has substantial ecological consequences for an individual, as immobile insects are not able to acquire food or escape from predators. While the measurement of the onset of chill coma can be challenging in immature insects, the recovery from chill coma, i.e., the ability of a specimen to exhibit a pre-determined behaviour, can be easily measured in both juveniles and adults (see Sinclair et al., 2015 for a discussion of this point).

Chill coma recovery time (CCRt) is expected to differ greatly among and within insect species and particularly among populations of different geographical origin (Andersen et al., 2015; Bowler and Terblanche, 2008; Chown, 2001). Within-population CCRt variations are

also frequently observed, as developmental plasticity (i.e., the environmental conditions experienced during preimaginal development partly determine adult stress tolerance) or individual acclimation (i.e., exposure to mild stress enhances individual stress tolerance; Colinet and Hoffmann, 2012; Terblanche and Chown, 2006) can affect this measure. Moreover, thermal tolerance can substantially vary over the course of an insect's life, but ontogenetic variations in the responses to low temperatures have been poorly examined compared to heat resistance (see Bowler and Terblanche, 2008). The existing studies have focused on preimaginal development or adult ageing (Colinet et al., 2013; David et al., 1998; Halle et al., 2015; Pujol-Lereis et al., 2014), while studies that included measures on both preimaginal and adult stages are commonly focused on a few developmental stages and age classes (e.g., Arias et al., 2011, but see Jensen et al., 2007 on *Drosophila melanogaster*). Differences between individuals in their cold tolerance are expected to have consequences for their overwintering success (Denlinger and Lee, 2010), but this prediction has only rarely been verified experimentally (e.g., Alford et al., 2014; Kostal et al., 2011).

The majority of studies investigating insect cold hardiness focus on the effects of a single exposure to low temperatures (Marshall and Sinclair, 2012). However, in the field, insects can be exposed to knockdown temperatures on multiple occasions over days, and these repeated exposures could affect insect performance in the following ways: 1) cold hardiness will progressively increase across successive exposures, depicting thermal acclimation; 2) cold hardiness will progressively decrease across successive exposures, revealing an accumulation of chill injuries; 3) cold hardiness will exhibit a complex pattern of a U-shaped cold hardiness response curve as a result of acclimation of the insect during the first sequence of the exposure to the cold conditions, followed by an accumulation phase of chill injuries (Colinet et al., 2016; Marshall and Sinclair, 2012, 2015).

In this study, we present the results of a pilot study on the thermal biology of the harlequin ladybird, *Harmonia axyridis* (Pallas, 1773). This species is currently considered one of the most invasive insects worldwide (Brown et al., 2011; Lombaert et al., 2011), with probable specific abilities to colonize cold environments (Barahona-Segovia et al. 2016). *H. axyridis* overwinters as an adult and creates large winter aggregations, frequently in human settlements (Berkvens et al., 2010). Previous studies measured supercooling points and lower lethal temperatures in adult *H. axyridis* originating from European and US invasive populations (Berkvens et al., 2010; Koch et al., 2004) and from Japanese native populations (Watanabe, 2002). To our knowledge, patterns of CCRt variation have not been investigated in this species thus far. We addressed three different questions using a single metric – CCRt: 1) we examined the influence of ontogeny on CCRt and hypothesized that CCRt would decrease from the 1st to 4th larval instar because of the increasing probability of exposure to low temperatures in nature (late larval instars can experience colder autumn weather in Central Europe); 2) we investigated the effect of repeated cold exposures on CCRt in adult ladybirds and expected CCRt prolongation because of the accumulation of chill injuries during exposures to cold and insufficient damage repair during very short warm periods among subsequent cold exposures; 3) finally, we were interested in the predictive power of pre-overwintering CCRt for winter survival and expected that individuals achieving lower pre-overwintering CCRt would suffer from lower winter mortality. The ecophysiological meaning of CCRt clearly differs for summer active (questions 1 and 2) and winter diapausing insects (question 3). For summer active insects we employed CCRt as a measure that allowed us to investigate physiological changes during the ontogenetic / physiological responses to repeated cold shocks. For winter diapausing insects we tested the ecological relevance of CCRt as a proxy of insect cold tolerance.

2. Materials and Methods

2.1. Insect collection and rearing

The parental individuals used in study were collected August 2015 from shrubs and lime trees in the university campus of the Czech University of Life Sciences Prague, Czech Republic (GPS: 50°8' N, 14°21' E; 300 m a.s.l.). After transportation to the laboratory, the beetles were sexed, and parental pairs were formed. Each couple was placed in a separate Petri dish (9 cm in diameter) containing crumpled filter paper strips, which provided a suitable substrate for egg laying. New egg clutches were placed into Petri dishes and exposed to a standardized laboratory condition (16L:8D photoperiod, 70 % humidity, 26 °C). A temperature of 26 °C is the optimal rearing temperature for *H. axyridis* (Hodek et al., 2012; Knapp and Nédved, 2013). Newly hatched larvae were fed *ad libitum* with eggs of *Ephestia kuehniella* (Zeller, 1879) (Lepidoptera: Pyralidae) and provided with water in cotton wool.

2.2. Chill coma induction and chill coma recovery time measurements

For all beetles investigated in the following experiments (sections 2.3., 2.4. and 2.5.), chill coma was induced and chill coma recovery time (CCRT) was measured in a standardized way. Pools of six individuals were placed individually into clean Petri dishes and transferred to -3 °C for two hours. A freezing climatic chamber made on order by AVIKO-PRAHA a.s. company was employed for this purpose. The temperature inside the climatic chamber was monitored with a datalogger Tinytag PLUS2 (Supporting Information file Fig. A.1). After 2 hours at -3 °C, the Petri dishes were returned to room temperature (22 ± 1 °C) in an air-conditioned laboratory and the beetles were turned onto their dorsum. The duration necessary for observing the first movement of an appendage (most often a leg or a palpus) was measured. As ladybirds can often “play dead” (remain motionless despite being already able

to move), individuals were stimulated by gentle teasing with an entomological pin every five seconds.

2.3. Experiment 1: Effects of ontogeny on chill coma recovery time

Three fecund parental pairs were selected at random, and their offspring were used in the laboratory experiment investigating the ontogenetic development of cold resistance in *H. axyridis*. Egg clutches, hatched larvae and 2nd generation adults were housed under the same rearing conditions as the parental pairs (16L:8D, 26 °C, 70 % humidity, food and water *ad libitum*). Following egg laying, egg clutches were randomly assigned to one of the following eight treatments to assess the effects of ontogenetic stages on CCRt: 1st larval instar, 2nd larval instar, 3rd larval instar, 4th larval instar, newly hatched adult (one-day old), four-day-old adult, 16-day-old adult and 32-day-old adult. CCRt was measured only once for each individual (not repeatedly in the course of the experiment). Ladybirds are long-lived insects, with adult longevity exceeding a year; therefore, senescence is expected to take place far later than at the age of 32 days (Hodek et al., 2012). The first and second instar larvae were reared together (a complete clutch) in one Petri dish. When the larvae moulted to the third instar, they were assigned to new Petri dishes in groups of four to five individuals. The CCRt of six larvae per parental pair was measured per treatment, and the CCRt of five individuals per sex per parental pair per treatment was recorded for adult treatments.

2.4. Experiment 2: Effects of repeated cold shock on the chill coma recovery time of adults

Twelve males and twelve females aged 14 to 20 days obtained from the same parental pairs as those individuals from the previous experiment were reared at a constant 26 °C, photoperiod

16L:8D, 70 % humidity and with food and water *ad libitum*. One day before the CCRt measurements, the beetles were individually transferred to Petri dishes and were provided with water only (this short food deprivation period was used to reduce the possible confusing effects of variable stomach contents across repeated measurements). Repeated exposures to knockdown temperatures were achieved as follows: each ladybird was transferred to -3.0 °C for two hours, then returned to room temperature (22 ± 1 °C) for 30 minutes before being transferred again to -3.0 °C. This cyclic procedure was repeated four times (i.e., four exposures of two hours at -3.0 °C), and the CCRt was repeatedly measured for each individual after each exposure to the knockdown temperature (four CCRt measurements for each individual).

2.5. Experiment 3: Chill coma recovery time and winter survival of adults

Five males and five females per parental pair were selected at random from 10 parental pairs (i.e., 100 ladybirds in total). Ladybird offspring were reared at 26 °C (photoperiod 16L:8D, 70 % humidity) and supplied with food (*Ephestia kuehniella* eggs) and water *ad libitum*. On October 7th, when the ladybirds were 15-25-day-old adults, the beetles were transferred to an 18 °C and 12L:12D photoperiod regime to initiate winter diapause. The insects were maintained individually in Petri dishes and supplied with water but without food (as they naturally stop feeding during winter diapause). On October 14th, the CCRt was measured (as described above) for each individual. One day later, the thermal regime was changed to fluctuating temperatures and a short photoperiod [8L(12 °C):16D(6 °C)] to mimic outdoor conditions in late autumn. Six days later, the overwintering phase started and the beetles, still maintained in the climatic chamber, were exposed to the mean temperature regime experienced in outdoor shelters in Prague (data were computed as the long-term mean

temperature 1 cm below the soil surface; hourly means from the meteorological station situated in the Crop Research Institute, Praha-Ruzyně were used). The beetles were maintained in clear Petri dishes containing a small watered piece of cotton wool to limit desiccation and placed into the climatic chamber set to the overwintering regime. The climatic chamber was adjusted each hour with a new temperature value (the temperatures applied are attached as Supporting Information file Table A.1). The mean temperature during the overwintering phase was 0.7 °C, and the minimum temperature reached was -4.1 °C.

In the course of the overwintering experiment, the beetles were checked at monthly intervals (November 16th, December 15th, January 14th, February 17th and March 17th, 2016 – when the overwintering experiment was terminated). During all inspections, the Petri dishes with beetles were moved to outdoor conditions to minimize unwanted warming of the experimental beetles, the survival of each individual was recorded, and a new piece of watered cotton wool was inserted into each Petri dish (the old one was removed). Live diapausing ladybirds are commonly tightly attached to the substrate, whereas dead individuals have dropped off and are lying freely on their dorsum. Petri dishes with dead individuals were not removed from the experiment as there was a little risk of false death records (an animal seeming to be dead but is still alive); however, no false records were revealed.

2.6. Statistical analyses

To analyse ontogenetic changes in the CCRt, a linear mixed-effects model (LME) was run using the “lme” function implemented in the “nlme” package (Pinheiro et al., 2013) for R (R Development Core Team, 2016). Parental pair identity was used as a random effect in our analysis. Ontogenetic stage was used as a fixed effect in our model. Tukey’s HSD test was employed to test for significant differences between particular ontogenetic stages using the

“glht” function as implemented in the “multcomp” package (Hothorn et al., 2008). Raw data serving as inputs for the abovementioned analysis are available in the Supporting Information file (Table A.2).

To analyse the effects of repeated cold exposures on the CCRt, a linear mixed-effects model (LME) was conducted using the “lme” function implemented in the “nlme” package (Pinheiro et al., 2013) for R (R Development Core Team, 2016). Beetle identity was used as a random effect in our analysis. The order of cold exposure (from 1 to 4; a continuous variable) was used as a fixed effect in our model, while a quadratic term was also included in the model for the case where the response would not be purely linear. The significance of the independent variables (order, order²) was tested using F-tests (“anova.lme” function). To test for the real necessity of both terms (order, order²) in the final model, the most parsimonious model based on AIC-values was also selected (Akaike, 1974). Raw data are available in the Supporting Information file (Table A.3).

To analyse the effects of CCRt on *H. axyridis* survival during winter, the Cox proportional-hazards model (Cox-PH) was run using the “coxph” function implemented in the “survival” package (Therneau, 2015) for R (R Development Core Team, 2016). Winter survival in months (one to five months) was used as the response variable, and CCRt was used as an independent variable in the model. Beetles that survived until the end of the experiment (March) were excluded in the model, as their future longevity was unknown. Raw data serving as inputs for the above analysis are available in the Supporting Information file (Table A.4).

The effects of sex on the CCRt were analysed in the preliminary analyses performed for all experiments. As there were no significant effects of sex in any model, this term was not included in the final models.

242

243 3. Results

244 There were significant changes in chill coma recovery time (CCRt) during the ontogenetic
245 development of the harlequin ladybird, *Harmonia axyridis* (LME: $F = 19.92$, $P < 0.001$). The
246 CCRt decreased by ca. 22 % in the course of early larval development from the 1st to 2nd
247 larval instars and reached the lowest value in teneral adults (between one and four days old
248 adults). The CCRt of older adults (16 or 32 days old adults) was increased and did not differ
249 from the CCRt of the 2nd, 3rd and 4th larval instars (Fig. 1).

250 Repeated two-hour exposures to sub-zero temperatures interrupted by 30-minute
251 recovery periods at 22 °C resulted in increasing the CCRt from the second cold exposure
252 onwards in *H. axyridis*. However, the CCRt decreased from the first to the second cold
253 exposure; thus, alongside the significant linear term (LME: $F = 23.92$, $P < 0.001$), there was
254 also a significant quadratic term in the final statistical model ($F = 11.24$, $P = 0.001$). The most
255 parsimonious model included both terms (order and order²) according to the AIC (Table 1),
256 demonstrating that a U-shaped response to repeated cold exposures was measured in *H.*
257 *axyridis* (Fig. 2).

258 No significant relationship was observed between the CCRt of the pre-overwintering
259 individuals and their ability to survive during the winter (Cox-PH: $z = -1.50$, $P = 0.13$). The
260 observed trend was even opposite that of the hypothesized one: the four individuals who had
261 the lowest CCRt died during the first months of the winter, whereas the three individuals with
262 the highest CCRt survived until the last month of the winter (two of them even survived the
263 whole winter; Fig. 3).

264

4. Discussion

In this work, we used CCRt to examine different aspects of ladybird thermal physiology: we 1) investigated the physiological changes during ontogeny, 2) examined the physiological responses of summer active ladybirds to repeated cold shocks, and 3) investigated the relevancy of the CCRt as a proxy for winter cold hardiness. Significant variations of the CCRt were measured across the ontogeny of *Harmonia axyridis*, with the lowest CCRt values observed in teneral adults. Repeated cold shocks altered the CCRt of *H. axyridis*, and a U-shaped response was measured, with reduced CCRt during the first phase, followed by a CCRt increase when the number of cold shocks was increased. Finally, we found no relationship between the CCRt of pre-overwintering individuals and their subsequent ability to survive during the winter months.

Changes in the degree of heat tolerance during ontogeny are widespread in insects, but adjustments to cold resistance during ontogeny have been less examined (e.g., Bowler and Terblanche, 2008; Klockmann et al., 2017; Knapp and Ndved, 2013; Zhao et al., 2017). Relevant experimental data that reported ontogenic variations of insect cold resistance exist mostly for insect model species (e.g., Arias et al., 2011; Colinet et al., 2013; Jensen et al., 2007; Pujol-Lereis et al., 2014; for the review of older studies see Bowler and Terblanche, 2008). In univoltine holometabolous species, differences in coping with cold exposures among stages could reflect adaptive responses, as the development stages can experience different thermal conditions, with the overwintering stage being most often the most cold-resistant one (Denlinger and Lee, 2010). Nevertheless, adjustments of insect cold hardiness are not only dictated by environmental temperatures, as differences have also been reported during the ontogeny of multivoltine species over the course of the growth season (e.g., Jensen et al., 2007). One of possible explanations is that levels of heat-shock proteins (HSPs), which are partially responsible for the actual cold resistance and are also involved in various stress

responses unrelated to cold challenge, vary substantially in the course of insect ontogeny (Colinet et al., 2013), including in the harlequin ladybird (Wang et al., 2017).

There is no unified ontogenetic profile of cold resistance observed for a wide range of insect species. In our study, CCRt was significantly higher in the 1st larval instar compared with later larval instars. Interestingly, Jensen et al. (2007) observed a distinct pattern of lower lethal temperature causing 50 % mortality (LT₅₀) in *Drosophila melanogaster*, which slightly increased from the 2nd to the 3rd larval instar. We measured the lowest CCRt value in teneral ladybird adults, which is consistent with the finding of Jensen et al., (2007) for *D. melanogaster*, but opposite the pattern reported for *Tenebrio molitor* (Arias et al., 2011). Differences in the changes of ontogenetic patterns of CCRts across species may result from the distinct life histories of the studied species, particularly in terms of overwintering strategies. Importantly, the contrasting cold resistance patterns among studies can also be caused by the various proxies that are employed, as these measures are not necessarily correlated (Andersen et al., 2015).

The higher cold resistance of teneral adults (few hours/days old) of *D. melanogaster* reported by Colinet et al. (2013) was also found in our study for *H. axyridis*. The cold resistance of adult ladybirds subsequently decreased with increasing age, a pattern also observed for *D. melanogaster* and *Ceratitis capitata* (David et al., 1998; Pujol-Lereis et al., 2014). Decreased cold hardiness with age in adult insects was thought to be related to senescence-associated processes (Bowler and Terblanche, 2008; Halle et al., 2015). However, this hypothesis is unlikely for our 32-day-old *H. axyridis* whose longevity can easily reach one year; senescence is thus expected to take place later in their lives (Hodek et al., 2012; note also that the CCRt slightly decreased from the age of 16 days to 32 days in our study). Moreover, increasing cold resistance in the course of senescence was observed for the sub-Antarctic fly *Anatalanta aptera* (Lalouette et al., 2010). Altogether, our results have important

methodological implications for future studies on insect cold hardiness, as the exact age of experimental animals should be precisely stated, and animals of the same stage/age should be compared with each other in studies investigating issues other than the effects of ontogeny.

Low temperatures cause chill injuries to chill-susceptible insects, and the extent of the injuries depends on the severity of the treatment, i.e., the severity of the temperatures experienced and the duration of exposure (Marshall and Sinclair, 2015; Overgaard and MacMillan, 2017). Despite the fact that thermal fluctuations are ubiquitous in nature, the majority of ecophysiological studies are focused on exposures of insects to a single period at low temperature (Marshall and Sinclair, 2012; see the literature focusing on fluctuating thermal regimes and repeated cold exposures, e.g., Colinet et al., 2016; Marshall and Sinclair, 2015). In this study, we observed the changes in CCRt values in *H. axyridis* subjected to repeated cold shocks and found that values decreased in the first stage followed by a continuous increase from the second exposure onwards. The observed U-shaped response curve depicted the interplay between acclimation (rapid cold-hardening response; Lee et al., 1987) elicited by the first cold exposure and the accumulation of chill injuries (e.g., the creation of reactive oxygen species; Lalouette et al., 2011) inflicted by each successive cold shock. The short warm periods may not have been sufficient for restoring the aerobic metabolism of the insects and completely repairing the physiological injuries caused by exposures to low temperatures (Colinet et al., 2016). A few hours may have been needed for a complete recovery of the injuries accumulated by cold-shocked ladybirds. For example, five cold exposures separated by recovery periods of 24 hours lowered CCRt values in the red flour beetle (Scharf et al., 2018). The physiological responses of insects to repeated cold stress are still not known in detail and it is possible that rapid switching between stress and injury repair states itself poses some physiological costs (Marshall and Sinclair, 2015).

We failed to report any significant relationship between the CCRt of pre-overwintering specimens of *H. axyridis* and their subsequent overwintering success. This finding is surprising as between- and within-species variations in CCRt commonly exist in insects (Chown, 2001). This variation seems to be adaptive as species' CCRt generally correlates to their distribution ranges (winter conditions experienced; Andersen et al., 2015; Chown, 2001), and overwintering individuals often have a significantly lower CCRt compared with their relatives sampled during the breeding season (Chown, 2001). Our data even showed slight support for the opposite pattern, i.e., individuals with the lowest CCRt died in early winter while those with the highest CCRt survived. A possible explanation is a higher metabolic rate in animals with lower CCRt (Williams et al., 2016). Animals with higher energy demands can die during overwintering due to the depletion of their energy reserves.

A possible shortcoming of our study is that overwintering success was investigated for laboratory-reared insects that experienced standardized environmental conditions. Moreover, their genetic variability may have been limited (all were offspring of 10 parental pairs). This experimental setting may have reduced the differences in CCRt values among individuals compared with the variation existing for individuals in the field. As laboratory rearing could affect ladybird overwintering performance (Berkvens et al., 2010), we encourage a repetition of this experiment using pre-overwintering individuals collected directly in the field. As winter survival in insects is a result of complex interactions between individual physiological traits and environmental conditions, future research investigating the effects of individual cold resistance (represented by various measures/proxies) on winter survival under various overwintering conditions for various insect species is needed to disentangle the effects of individual traits on winter survival in insects. If additional studies confirm that CCRt is not the most reliable predictor of insect overwintering success, chill coma temperatures or lower

lethal temperatures, both good predictors of the latitudinal distribution of *Drosophila* species, (Andersen et al., 2015) might be used as better indicators of overwintering success.

5. Conclusions

This study provides the first measurements of chill coma recovery time (CCRt) in the invasive harlequin ladybird, *Harmonia axyridis*. We showed substantial changes in CCRt in the course of the ontogenetic development of *H. axyridis*. The significant ontogenetic signal in *H. axyridis* cold hardiness poses important methodological implications for future studies, as animals of the same stage/age should be compared with each other. We also showed that repeated exposures to cold shocks interrupted by short warm periods caused a U-shaped cold resistance response. Individual CCRt was a bad predictor of *H. axyridis* overwintering success. Together with the limited evidence in the existing literature, our results indicated that CCRt is not a reliable measure of insect overwinter success. Future studies investigating the relationship between prewintering CCRt and following winter survival for various insect species are needed to extend the generality of this pattern.

Acknowledgements

We are grateful to Michal Řeřicha and Radek Svoboda for their help with the laboratory rearing of ladybirds and realization of the overwintering experiment and to Oldřich Nedvěd for sharing his knowledge of *H. axyridis* thermal biology. We are also grateful to six reviewers who significantly improved the language quality of the text and provided us with insightful suggestions on a previous version of the manuscript.

387 **Funding:** This study was supported by grant no. 42900/1312/3166 awarded by the Internal
388 Grant Agency of the Czech University of Life Sciences Prague.

389

390 **Appendices:**

391 The following additional supporting information may be found in the online version of this
392 article.

393 **Table A.1 – Temperature profile experienced by ladybirds during overwintering in**
394 **computer-instructed freezing climatic chambers.**

395 **Table A.2 – Raw data analysed in experiment 1 (Ontogeny).**

396 **Table A.3 – Raw data analysed in experiment 2 (Repeated cold shock).**

397 **Table A.4 – Raw data analysed in experiment 3 (Winter survival).**

398 **Fig. A.1 – Temperature record from a datalogger placed between the Petri dishes with**
399 **ladybirds during chill coma induction.**

Tables

Table 1. Effect of repeated cold shocks on chill coma recovery time – model selection.

The most parsimonious linear mixed effect model (the first line) was selected on the basis of the Akaike information criterion (AIC). The term “order” indicates the linear effects of the cold shock order on chill coma recovery time; the term “order²” indicates the quadratic (non-linear) effect of the cold shock order on the chill coma recovery time. A difference in the AIC-value (ΔAIC) > 2 indicates a significant difference between models, i.e., the model with lower AIC-value is significantly more parsimonious.

Model structure			
Independent variable	Random effect	AIC-value	ΔAIC
~ order + order ²	individual beetle	826.303	0.000
~ order ²	individual beetle	836.124	9.821
~ order	individual beetle	837.654	11.352

Figures

Fig. 1. Effect of ontogenetic stage on cold resistance of *Harmonia axyridis*. The mean chill coma recovery time (CCRt) values + SEM based on 18 larvae measured per instar (grey bars) and 30 adults (15 males and 15 females) measured per adult age class (black bars) are shown. Significant differences (Tukey post hoc test: $P < 0.05$) in CCRt among particular ontogenetic stages are indicated by unshared letters above the bars.

Fig. 2. Effect of repeated cold shocks on chill coma recovery time in adult *Harmonia*

***axyridis*.** Mean chill coma recovery time (CCRt) values + SEM are shown. Each ladybird (24 in total) was exposed to four successive cold shocks (two hours at -3 °C) interrupted by short warm periods (30 minutes at 22 °C). The CCRt was measured for each individual after each cold shock.

Fig. 3. Relationships between individual pre-overwintering chill coma recovery time and subsequent winter survival. Dates on the vertical axis indicate control dates, i.e., the dates on which a given beetle was seen alive for the last time. Survival until March 17th indicates successfully completed overwintering. Short vertical lines indicate the mean pre-overwintering chill coma recovery time (CCRt) achieved by individuals who were seen alive for the last time during a given control.

- 433 Akaike, H., 1974. New look at statistical-model identification. IEEE Trans. Autom. Control. AC19, 716-
434 723.
- 435 Alford, L., Andrade, T.O., Georges, R., Burel, F., van Baaren, J., 2014. Could Behaviour and Not
436 Physiological Thermal Tolerance Determine Winter Survival of Aphids in Cereal Fields? Plos One 9,
437 e114982.
- 438 Andersen, J.L., Manenti, T., Sorensen, J.G., MacMillan, H.A., Loeschcke, V., Overgaard, J., 2015. How
439 to assess *Drosophila* cold tolerance: chill coma temperature and lower lethal temperature are the
440 best predictors of cold distribution limits. Funct. Ecol. 29, 55-65.
- 441 Araujo, M.B., Ferri-Yanez, F., Bozinovic, F., Marquet, P.A., Valladares, F., Chown, S.L., 2013. Heat
442 freezes niche evolution. Ecol. Lett. 16, 1206-1219.
- 443 Arias, M.B., Poupin, M.J., Lardies, M.A., 2011. Plasticity of life-cycle, physiological thermal traits and
444 Hsp70 gene expression in an insect along the ontogeny: Effect of temperature variability. J. Therm.
445 Biol. 36, 355-362.
- 446 Barahona-Segovia, R.M., Grez, A.A., Bozinovic, F., 2016. Testing the hypothesis of greater
447 eurythermality in invasive than in native ladybird species: from physiological performance to life-
448 history strategies. Ecol. Entomol. 41, 182-191.
- 449 Berkvens, N., Bale, J.S., Berkvens, D., Tirry, L., De Clercq, P., 2010. Cold tolerance of the harlequin
450 ladybird *Harmonia axyridis* in Europe. J. Insect Physiol. 56, 438-444.
- 451 Bowler, K., Terblanche, J.S., 2008. Insect thermal tolerance: what is the role of ontogeny, ageing and
452 senescence? Biol. Rev. 83, 339-355.
- 453 Brown, P.M.J., Thomas, C.E., Lombaert, E., Jeffries, D.L., Estoup, A., Handley, L.-J.L., 2011. The global
454 spread of *Harmonia axyridis* (Coleoptera: Coccinellidae): distribution, dispersal and routes of
455 invasion. Biocontrol 56, 623-641.
- 456 Chown, S.L., 2001. Physiological variation in insects: hierarchical levels and implications. J. Insect
457 Physiol. 47, 649-660.
- 458 Colinet, H., Hoffmann, A.A., 2012. Comparing phenotypic effects and molecular correlates of
459 developmental, gradual and rapid cold acclimation responses in *Drosophila melanogaster*. Funct.
460 Ecol. 26, 84-93.
- 461 Colinet, H., Renault, D., Javal, M., Berkova, P., Simek, P., Kostal, V., 2016. Uncovering the benefits of
462 fluctuating thermal regimes on cold tolerance of *Drosophila* flies by combined metabolomic and
463 lipidomic approach. BBA - Mol. Cell Biol. L. 1861, 1736-1745.
- 464 Colinet, H., Siaussat, D., Bozzolan, F., Bowler, K., 2013. Rapid decline of cold tolerance at young age is
465 associated with expression of stress genes in *Drosophila melanogaster*. J. Exp. Biol. 216, 253-259.
- 466 David, R.J., Gibert, P., Pla, E., Petavy, G., Karan, D., Moreteau, B., 1998. Cold stress tolerance in
467 *Drosophila*: Analysis of chill coma recovery in *D. melanogaster*. J. Therm. Biol. 23, 291-299.
- 468 Denlinger, D.L., Lee, R.E., 2010. Low temperature biology of insects. Cambridge University Press,
469 Cambridge, UK.
- 470 Halle, S., Nowizki, A., Scharf, I., 2015. The consequences of parental age for development, body mass
471 and resistance to stress in the red flour beetle. Biol. J. Linn. Soc. 115, 305-314.
- 472 Hodek, I., Honěk, A., van Emden, H.F., 2012. Ecology and Behaviour of the Ladybird Beetles
473 (Coccinellidae). Willey-Blackwell, Chichester, UK.
- 474 Hothorn, T., Bretz, F., Westfall, P., 2008. Simultaneous inference in general parametric models.
475 Biomet. J. 50, 346-363.
- 476 Jensen, D., Overgaard, J., Sorensen, J.G., 2007. The influence of developmental stage on cold shock
477 resistance and ability to cold-harden in *Drosophila melanogaster*. J. Insect Physiol. 53, 179-186.
- 478 Klockmann, M., Gunter, F., Fischer, K., 2017. Heat resistance throughout ontogeny: body size
479 constrains thermal tolerance. Glob. Change Biol. 23, 686-696.
- 480 Knapp, M., Nēdved, O., 2013. Gender and Timing during Ontogeny Matter: Effects of a Temporary
481 High Temperature on Survival, Body Size and Colouration in *Harmonia axyridis*. Plos One 8, e74984.

482 Koch, R.L., Carrillo, M.A., Venette, R.C., Cannon, C.A., Hutchison, W.D., 2004. Cold hardiness of the
 483 multicolored Asian lady beetle (Coleoptera : Coccinellidae). *Environ. Entomol.* 33, 815-822.
 484 Kostal, V., Dolezal, P., Rozsypal, J., Moravcova, M., Zahradnickova, H., Simek, P., 2011. Physiological
 485 and biochemical analysis of overwintering and cold tolerance in two Central European populations of
 486 the spruce bark beetle, *Ips typographus*. *J. Insect Physiol.* 57, 1136-1146.
 487 Lalouette, L., Vernon, P., Amat, H., Renault, D., 2010. Ageing and thermal performance in the sub-
 488 Antarctic wingless fly *Anatalanta aptera* (Diptera: Sphaeroceridae): older is better. *Biol. Lett.* 6, 346-
 489 349.
 490 Lalouette, L., Williams, C.M., Hervant, F., Sinclair, B.J., Renault, D., 2011. Metabolic rate and oxidative
 491 stress in insects exposed to low temperature thermal fluctuations. *Comp. Biochem. Physiol., Part A*
 492 *Mol. Integr. Physiol.* 158, 229-234.
 493 Lee, R.E., Chen, C.P., Denlinger, D.L., 1987. A rapid cold-hardening process in insects. *Science* 238,
 494 1415-1417.
 495 Lombaert, E., Guillemaud, T., Thomas, C.E., Handley, L.J.L., Li, J., Wang, S., Pang, H., Goryacheva, I.,
 496 Zakharov, I.A., Jousset, E., Poland, R.L., Migeon, A., van Lenteren, J., De Clercq, P., Berkvens, N.,
 497 Jones, W., Estoup, A., 2011. Inferring the origin of populations introduced from a genetically
 498 structured native range by approximate Bayesian computation: case study of the invasive ladybird
 499 *Harmonia axyridis*. *Mol. Ecol.* 20, 4654-4670.
 500 MacMillan, H.A., Sinclair, B.J., 2011. Mechanisms underlying insect chill-coma. *J. Insect Physiol.* 57,
 501 12-20.
 502 MacMillan, H.A., Williams, C.M., Staples, J.F., Sinclair, B.J., 2012. Reestablishment of ion homeostasis
 503 during chill-coma recovery in the cricket *Gryllus pennsylvanicus*. *PNAS* 109, 20750-20755.
 504 Marshall, K.E., Sinclair, B.J., 2012. The impacts of repeated cold exposure on insects. *J. Exp. Biol.* 215,
 505 1607-1613.
 506 Marshall, K.E., Sinclair, B.J., 2015. The relative importance of number, duration and intensity of cold
 507 stress events in determining survival and energetics of an overwintering insect. *Funct. Ecol.* 29, 357-
 508 366.
 509 Overgaard, J., MacMillan, H.A., 2017. The Integrative Physiology of Insect Chill Tolerance. *Ann. Rev.*
 510 *Physiol.*, 79, 187-208.
 511 Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., R Development Core Team, 2013. *nlme: linear and*
 512 *nonlinear mixed effects models*. R package version 3.1-107.
 513 Pujol-Lereis, L.M., Rabossi, A., Quesada-Allue, L.A., 2014. Analysis of survival, gene expression and
 514 behavior following chill-coma in the medfly *Ceratitis capitata*: Effects of population heterogeneity
 515 and age. *J. Insect Physiol.* 71, 156-163.
 516 R Development Core Team, 2016. *R: A language and environment for statistical computing*. R
 517 Foundation for Statistical Computing, Vienna, Austria. URL <http://www.R-project.org>.
 518 Scharf I., Wertheimer, K.O., Xin, J.L., Gilad, T., Goldenberg, I., Subach, A., 2018. Context-dependent
 519 effects of cold stress on behavioral, physiological, and life-history traits of the red flour beetle. *Insect*
 520 *Sci.* in press.
 521 Sinclair, B.J., Alvarado, L.E.C., Ferguson, L.V., 2015. An invitation to measure insect cold tolerance:
 522 Methods, approaches, and workflow. *J. Therm. Biol.* 53, 180-197.
 523 Teets, N.M., Denlinger, D.L., 2013. Physiological mechanisms of seasonal and rapid cold-hardening in
 524 insects. *Physiol. Entomol.* 38, 105-116. Terblanche, J.S., Hoffmann, A.A., Mitchell, K.A., Rako, L., le
 525 Roux, P.C., Chown, S.L., 2011. Ecologically relevant measures of tolerance to potentially lethal
 526 temperatures. *J. Exp. Biol.* 214, 3713-3725.
 527 Terblanche, J.S., Chown, S.L., 2006. The relative contributions of developmental plasticity and adult
 528 acclimation to physiological variation in the tsetse fly, *Glossina pallidipes* (Diptera, Glossinidae). *J.*
 529 *Exp. Biol.* 209, 1064-1073.
 530 Therneau, T., 2015. *A Package for Survival Analysis in S*. version 2.38. [https://CRAN.R-](https://CRAN.R-project.org/package=survival)
 531 [project.org/package=survival](https://CRAN.R-project.org/package=survival).

532 Wang, H.J., Shi, Z.K., Shen, Q.D., Xu, C.D., Wang, B., Meng, Z.J., Wang, S.G., Tang, B., Wang, S., 2017.
533 Molecular Cloning and Induced Expression of Six Small Heat Shock Proteins Mediating Cold-Hardiness
534 in *Harmonia axyridis* (Coleoptera: Coccinellidae). Front. Physiol. 8: 60.
535 Watanabe, M., 2002. Cold tolerance and myo-inositol accumulation in overwintering adults of a lady
536 beetle, *Harmonia axyridis* (Coleoptera : Coccinellidae). Eur. J. Entomol. 99, 5-9.
537 Williams, C.M., Szejner-Sigal, A., Morgan, T.J., Edison, A.S., Allison, D.B., Hahn, D.A., 2016. Adaptation
538 to low temperature exposure increases metabolic rates independently of growth rates. Integr. Comp.
539 Biol. 56, 62-72.
540 Zhao, F., Hoffmann, A.A., Xing, K., Ma, C.S., 2017. Life stages of an aphid living under similar thermal
541 conditions differ in thermal performance. J. Insect Physiol. 99, 1-7.

542