



HAL
open science

Imaging in Systemic Sclerosis make "US" great again?

Alain Lescoat, Cristian Caimmi, Éric Hachulla, Patrick Jegou, Guillaume Coiffier

► **To cite this version:**

Alain Lescoat, Cristian Caimmi, Éric Hachulla, Patrick Jegou, Guillaume Coiffier. Imaging in Systemic Sclerosis make "US" great again?. *Arthritis Care & Research = Arthritis Care and Research*, 2019, 71 (5), pp.694-695. 10.1002/acr.23690 . hal-01834091

HAL Id: hal-01834091

<https://hal-univ-rennes1.archives-ouvertes.fr/hal-01834091>

Submitted on 10 Sep 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Imaging in Systemic Sclerosis: *make “US” great again?*

Alain Lescoat^{1,2} MD, Cristian Caimmi³ MD, Éric Hachulla⁴ MD PhD, Patrick Jégo^{1,2} MD PhD, Guillaume Coiffier^{5,6} MD.

1-Univ Rennes, CHU Rennes, Inserm, EHESP, Irset (Institut de recherche en santé, environnement et travail) – UMR_S 1085, F-35000 Rennes, France

2-Department of Internal Medicine, Rennes University Hospital, 35203, Rennes, France.

3-Department of Rheumatology, University of Verona, Verona, Italy

4-Department of Internal Medicine, CHU Lille, Lille Nord-de-France University Hospital, Lille, France.

5-Department of Rheumatology, Rennes University Hospital, 35203, Rennes, France

6- INSERM UMR 991, 35043 Rennes, France; Faculté de médecine, université Rennes 1, 35043 Rennes, France.

Corresponding author: Alain Lescoat

Department of Internal Medicine, CHU Rennes, South Hospital, 16 bd de Bulgarie - 35203 RENNES Cedex 2 - BP 90347

E-Mail : Alain.lescoat@chu-rennes.fr

Phone: +33 2 99 26 71 28 fax : +33 2 99 26 71 98

Financial disclosures:

A Lescoat, C Caimmi and G Coiffier report no conflict of interest.

E Hachulla received speaking fees from Actelion Pharmaceuticals Ltd, Glaxosmithkline and Bayer (<€10,000 each) outside of the current study. P Jégo received speaking fees from Actelion Pharmaceuticals Ltd and Bayer (<€10 000 each) outside of the current study.

Funding:

No specific funding was received to carry out the work described in this manuscript.

Keywords:

Systemic sclerosis, Scleroderma, Digital Ulcers, Power Doppler Ultrasonography, Ultrasound, Skin.

Dear Editor,

In this issue of *Arthritis Care & Research*, T. Santiago and colleagues [1] have conducted an informative systematic literature review on the relevance of ultrasound (US) for the assessment of skin involvement in systemic sclerosis (SSc). In this *Original Article*, the authors highlight the heterogeneous results concerning the reliability of US in studies published to date, but also underscore very promising data from recently published articles [2]. Moreover, this work raises the issue considering US evaluation as an interesting alternative to mRSS for skin assessment in SSc. Therefore, beyond imaging considerations, this article questions the place of US evaluation in the global management of SSc. This topic is still a controversial matter as new assessment technics are emerging for microvascular, macrovascular and musculoskeletal evaluations in SSc [3,4].

Nonetheless, a central question remains concerning these new evaluation tools for this disease: new imaging technologies may come, indeed, but for what purpose in the end?

In addressing this issue from a purely descriptive perspective, as the question of reframing and reshaping phenotypes in SSc is arising [5], US evaluation may offer a more precise description of vascular and musculoskeletal manifestations of the disease. This is all the more relevant as always more performant probes (>18Mhz) lead to always more precise descriptions and high-quality images. US characterization of these manifestations, combined with a precise clinical phenotyping [6] may help to enlarge the current view of SSc bringing the disease into the “technicolor” enriched vision of SSc suggested by Ligon and Wigley [5]. Beyond echocardiography and the detection of Pulmonary Arterial Hypertension, US may also have a part to play in the early detection of visceral manifestations of the disease such as lung fibrosis, as suggested by recent studies evaluating the diagnostic performances of US in this field [7].

In a concrete and practical approach, US may be one step ahead of many assessment tools as it is both widely available and non-invasive/non-irradiating. The issue of reproducibility and observer-dependent reliability may be easily addressed when US evaluation is performed by a trained rheumatologist [8]. As musculoskeletal US assessment is now included in many training programs for young Rheumatologists, switching from joints to other organs might be all the easier. Further studies are nonetheless needed to sketch simple and reproducible consensual evaluation criteria, constituting the first step for a coordinate initiative to homogenize practices concerning vascular, skin, lung and musculoskeletal US evaluation in SSc. This step is essential to foster future multi-centric studies.

From a rheumatologic perspective, US assessment may deserve a specific attention as it could have a part to play in the management of SSc in the future. Considering hand evaluation, one of the main challenges for clinicians dealing with SSc, is the multiplicity of causes that lead to hand functional disability [9,10]: skin roughness limiting mobility; Raynaud phenomenon, digital ulcers, calcinosis or acro-osteolysis hindering precise gripping; synovial or tenosynovial involvement responsible for friction rubs and pain... Deciphering the cause of pain itself can also be challenging in SSc patients. Arthralgia or joint pain can result from numerous and co-existing causes, ranging from specific inflammatory synovitis to sclerotic tenosynovitis, mechanical digital ulcers, osteoarthritis or intra-articular calcinosis. Understanding the precise cause of pain and disability is also a central question as it may lead to a guided and more adapted therapeutic management. US offers the unique opportunity of combined and simultaneous dynamic evaluations of vascular, musculoskeletal and skin involvement: a single tool to assess a multi-causal impairment [11].

In addressing the issue of new imaging technics in SSc, one must not forget that clinical evaluation must remain the keystone for the management of SSc patients. Nonetheless, clinical examination may rise unsolved questions that could be answered through US assessment.

References :

- 1- Santiago T, Santiago M, Ruaro B, Salvador MJ, Cutolo M, da Silva JAP. Ultrasonography for the assessment of skin in systemic sclerosis: a systematic review. *Arthritis Care Res (Hoboken)*. 2018 May 21.
- 2- Sulli A, Ruaro B, Smith V, Paolino S, Pizzorni C, Pesce G, et al. Subclinical dermal involvement is detectable by high frequency ultrasound even in patients with limited cutaneous systemic sclerosis. *Arthritis Res Ther*. 2017;19(1):61.
- 3- Cutolo M, Damjanov N, Ruaro B, et al. Imaging of connective tissue diseases: Beyond visceral organ imaging? *Best Pract Res Clin Rheumatol*. 2016;30(4):670-87.
- 4- Cutolo M, Ruaro B, Smith V. Macrocirculation versus microcirculation and digital ulcers in systemic sclerosis patients: Macro-microcirculation and scleroderma. *Rheumatology (Oxford)*. 2017;56(11):1834-1836.
- 5- Ligon CB, Wigley FM. Editorial: Scleroderma: Bringing a Disease From Black-and-White Into Technicolor. *Arthritis Rheumatol*. 2015;67(12):3101-3.
- 6- Allanore Y, Distler O, Matucci-Cerinic M, Denton CP. Review: Defining a Unified Vascular Phenotype in Systemic Sclerosis. *Arthritis Rheumatol*. 2018;70(2):162-170.
- 7- Tardella M, Di Carlo M, Carotti M, Filippucci E, Grassi W, Salaffi F. Ultrasound B-lines in the evaluation of interstitial lung disease in patients with systemic sclerosis: Cut-off point definition for the presence of significant pulmonary fibrosis. *Medicine (Baltimore)*. 2018;97(18):e0566.
- 8- Ruaro B, Sakellariou G, Cutolo M, Iagnocco A. Well-trained sonographers are worth their weight in gold: ultrasound in systemic sclerosis. *Rheumatology (Oxford)*. 2018 Jan 31. doi:10.1093/rheumatology/kex481. [Epub ahead of print]
- 9- Hachulla E, Hatron PY, Carpentier P, Agard C, Chatelus E, Jegou P et al; (SEDUCE study group). Efficacy of sildenafil on ischaemic digital ulcer healing in systemic sclerosis: the placebo-controlled SEDUCE study. *Ann Rheum Dis*. 2016;75(6):1009-15.
- 10- Mouthon L, Carpentier PH, Lok C, Clerson P, Gressin V, Hachulla E, et al; (ECLIPSE Study Investigators). Ischemic digital ulcers affect hand disability and pain in systemic sclerosis. *J Rheumatol*. 2014;41(7):1317-23.
- 11- Lescoat A, Coiffier G, de Carlan M, Droitcourt C, Ballerie A, Cazalets C, et al. Combination of Capillaroscopic and Ultrasonographic Evaluations in Systemic Sclerosis: Results of a Cross-Sectional Study. *Arthritis Care Res (Hoboken)*. 2018;70(6):938-43.