Positive correlations between free vitamin D and bone variables in a group of young Lebanese men Article type

Abir Alwan, Maroun Rizkallah, Ghassan Maalouf, Joseph Matta, Fabienne Frenn, Abdel Jalil Berro, Angélique Barakat, Falah Bachour, Amer Sebaaly, Mirza Howayek, et al.

► To cite this version:

Abir Alwan, Maroun Rizkallah, Ghassan Maalouf, Joseph Matta, Fabienne Frenn, et al.. Positive correlations between free vitamin D and bone variables in a group of young Lebanese men Article type. *Journal of Clinical Densitometry*, 2018, 21 (3), pp.459-461. 10.1016/j.jocd.2018.02.001 . hal-01877290

HAL Id: hal-01877290

<https://hal-univ-rennes1.archives-ouvertes.fr/hal-01877290>

Submitted on 19 Sep 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Positive correlations between free vitamin D and bone variables in a group of young Lebanese men

Article type: Letter to the Editor

Abir Alwan^{1,2}, Maroun Rizkallah³, Ghassan Maalouf³, Joseph Matta^{4,5}, Fabienne Frenn⁵, Abdel-Jalil Berro^{1,6}, Angélique Barakat³, Falah Bachour³, Amer Sebaaly³, Mirza Howayek³, Hassane Zouhal², **Rawad El Hage**^{1,*}

1. Department of Physical Education, Faculty of Arts and Social Sciences, University of Balamand, El-Koura, Lebanon.

2. Movement Sport and Health Sciences Laboratory, University of Rennes 2-ENS Cachan, Rennes, France.

3. Bellevue University Medical Center, Faculty of Medicine, Saint Joseph University, Mansourieh, Lebanon.

4. Industrial Research Institute, Baabda, Lebanon.

5. Department of Nutrition, Faculty of Pharmacy, Saint Joseph University, Lebanon.

6. EA-3300: APERE, Exercise Physiology and Rehabilitation Laboratory, Picardie Jules Verne University, Sport Sciences Department, Avenue Paul Claudel, F-80025, Amiens Cedex, France.

* Corresponding author: Rawad El Hage

E-mail address: rawadelhage21@hotmail.com

Tel.: 00961/3/713605

Fax: 00961/6/930278

Department of Physical Education

Faculty of Arts and Social Sciences

University of Balamand, P.O. Box 100 Tripoli, Lebanon.

Dear Editor,

Vitamin D deficiency is a major public health problem in many countries (1-7). The prevalence of vitamin D deficiency is also very common in Lebanese adults (1-7). Although vitamin D is very important for bone health, several studies have failed to find a positive relationship between 25-hydroxyvitamin D [25(OH)D] and bone mineral density (8-10). However, we have previously shown that serum vitamin D is a positive determinant of composite indices of femoral neck strength in young adults (11, 12). Bone strength is not only influenced by bone mineral density (BMD) but also by bone geometry (13). Several DXA-based variables other than BMD can predict fracture risk in elderly subjects such as bone mineral apparent density (BMAD), cross-sectional area (CSA) of the femoral neck, compressive strength index (CSI), bending strength index (BSI), impact strength index (ISI) and trabecular bone score (TBS) (14, 15). In order to maintain healthy bones, adequate serum vitamin D levels should be maintained (16). Approximately 90% of 25(OH)D is bound to binding protein (16, 17). In general, the free fraction of a hormone is the biological active component (16, 17). Accordingly, one would expect the bone strength variables to correlate more strongly to free fraction of vitamin D than to total vitamin D (18, 19). In line with this hypothesis, Powe et al. (20) have shown that free vitamin D is positively correlated to BMD in a group of young adults. Johnsen et al. (21) have obtained similar results in a population of postmenopausal women. These previous studies did not focus on evaluating other bone health parameters such as bone geometry and architecture. In a recent study, we have demonstrated free vitamin D serum level is a stronger positive determinant of bone parameters and hip bone strength indices in young female adults than total serum Vitamin D (22). The aim of the current study was to explore the relationships between serum free vitamin D and several bone parameters (bone mass, BMD, hip bone geometry

indices, composite indices of femoral neck strength and TBS) in a group of young Lebanese men.

Forty young men whose ages ranged from 18 to 35 years participated in the present study. The 40 participants were recruited from 2 private universities located in North Lebanon. All participants were nonsmokers and had no history of major orthopedic problems or other disorders known to affect bone metabolism or physical tests of the study. Other inclusion criteria included no diagnosis of comorbidities and no history of fracture. An informed written consent was obtained from the participants. The current study was approved by the University of Balamand Ethics Committee. Weight and height were measured, and body mass index (BMI) was calculated. BMC (in gram) and BMD (in gram per square centimeter) were determined for each individual by DXA (GE Healthcare, Madison, WI, USA) at whole body (WB), lumbar spine (L1–L4), total hip (TH), and femoral neck (FN). Femoral neck (FN) cross-sectional area (CSA), FN cross-sectional moment of inertia (CSMI), and L1–L4 TBS were also evaluated by DXA (22). In our laboratory, the coefficients of variation were <1% for BMC and BMD (22). Composite indices of FN strength were calculated as previously described (14, 15). FN compressive strength index (CSI), FN bending strength index (BSI), and FN impact strength index (ISI) were calculated (14, 15). Compressive strength $((FN\ BMD * FN\ width) / weight)$ and bending strength $((FN\ BMD * [FN\ width]^2) / (hip\ axis\ length * weight))$ express the forces that the FN has to withstand in weight-bearing, whereas impact strength $((FN\ BMD * FN\ width * hip\ axis\ length) / (height * weight))$ expresses the energy that the FN has to absorb in an impact from standing height (14, 15). Free vitamin D, vitamin D binding protein and total vitamin D were measured as previously described (22).

Free vitamin D was positively correlated to WB BMC ($r = 0.39$; $p = 0.02$), WB BMD ($r = 0.35$; $p = 0.04$), TBS ($r = 0.43$; $p = 0.01$), TH BMD ($r = 0.47$; $p = 0.003$), FN BMD ($r = 0.45$; $p = 0.006$), ISI ($r = 0.41$; $p = 0.01$), CSA ($r = 0.45$; $p = 0.006$) and CSMI ($r = 0.41$; $p = 0.01$). Vitamin D binding protein was positively correlated to TBS ($r = 0.44$; $p = 0.008$) and CSI ($r = 0.51$; $p = 0.001$). Total vitamin D was not correlated to bone variables. The present study conducted on a group of young Lebanese men mainly shows that serum free vitamin D is a positive determinant of several bone variables (BMC, BMD, TBS and bone geometry indices). Although the strengths of the associations between free vitamin D and bone variables were moderate, this is the first study to show positive associations between serum free vitamin D and several DXA variables in young men. This correlation is of great importance making free vitamin D a variable to test routinely in youth. Accordingly, our study suggests that treatment for free vitamin D deficiency may improve bone health in youth, and therefore bone fragility fractures at a later age would be reduced. Enough evidence is now present to support the realization of a larger-scale powered study evaluating the association between free vitamin D and bone parameters in young and elderly populations. Such correlations, if present, should make free vitamin D dosing a routine in young and elderly subjects, especially if diagnosing a deficiency and treating it could improve bone quality and therefore reduce fragility fracture incidence at a later age.

References

1. Lips P. 2007 Vitamin D status and nutrition in Europe and Asia. *J Steroid Biochem Mol Biol* 103(3-5):620–625.

2. Medlej-Hashim M, Jounblat R, Hamade A, et al. 2015 Hypovitaminosis D in a Young Lebanese Population: Effect of GC Gene Polymorphisms on Vitamin D and Vitamin D Binding Protein Levels. *Ann Hum Genet* 79(6):394–401.
3. Arabi A, El Rassi R, El-Hajj Fuleihan G. 2010 Hypovitaminosis D in developing countries-prevalence, risk factors and outcomes. *Nat Rev Endocrinol* 6(10):550–561.
4. Gannagé-Yared MH, Chemali R, et al. 2000 Hypovitaminosis D in a sunny country: relation to lifestyle and bone markers. *J Bone Miner Res* 15(9):1856–1862.
5. Fuleihan GE, Deeb M. 1999 Hypovitaminosis D in a sunny country. *N Engl J Med* 340(23):1840–1841.
6. Arabi A, Baddoura R, El-Rassi R, et al. 2012 PTH level but not 25 (OH) vitamin D level predicts bone loss rates in the elderly. *Osteoporos Int* 23(3):971–980.
7. El-Hajj Fuleihan G, Nabulsi M, Choucair M, et al. 2001 Hypovitaminosis D in healthy schoolchildren. *Pediatrics* 107(4):E53.
8. Jemielita TO, Leonard MB, Baker J, et al. 2016 Association of 25-hydroxyvitamin D with areal and volumetric measures of bone mineral density and parathyroid hormone: impact of vitamin D-binding protein and its assays. *Osteoporos Int* 27(2):617–626.
9. Kremer R, Campbell PP, Reinhardt T, et al. 2009 Vitamin D status and its relationship to body fat, final height, and peak bone mass in young women. *J Clin Endocrinol Metab* 94(1):67–73.
10. Sherman SS, Tobin JD, Hollis BW, et al. 1992 Biochemical parameters associated with low bone density in healthy men and women. *J Bone Miner Res* 7(10):1123–1130.
11. Alwan A, El Khoury C, Zouhal H, et al. 2016 Vitamin D Level and Composite Indices of Femoral Neck Strength in a Group of Young Lebanese Men. *J Clin Densitom* 19(4):492–493.

12. Berro AJ, Alwan A, Zouhal H, et al. 2016 Vitamin D Level and Composite Indices of Femoral Neck Strength in a Group of Young Lebanese Women. *J Clin Densitom.* In press.
13. Beck TJ. 2003 Measuring the structural strength of bones with dual-energy X-ray absorptiometry: principles, technical limitations, and future possibilities. *Osteoporos Int* 14:S81–S88.
14. Ayoub ML, Maalouf G, Bachour F, et al. 2014 DXA-based variables and osteoporotic fractures in Lebanese postmenopausal women. *Orthop Traumatol Surg Res* 100:855–858.
15. Karlamangla AS, Barrett-Connor E, Young J, et al. 2004 Hip fracture risk assessment using composite indices of femoral neck strength: the Rancho Bernardo study. *Osteoporos Int* 15:62–70.
16. Fuleihan Gel-H, Bouillon R, Clarke B, et al. 2015 Serum 25-Hydroxyvitamin D Levels: Variability, Knowledge Gaps, and the Concept of a Desirable Range. *J Bone Miner Res* 30(7):1119–1133.
17. Chun RF, Peercy BE, Orwoll ES, et al. 2014 Vitamin D and DBP: the free hormone hypothesis revisited. *J Steroid Biochem Mol Biol* 144 Pt A:132–137.
18. Bouillon R. 2016 Free or Total 25OHD as Marker for Vitamin D Status? *J Bone Miner Res* 31(6):1124–1127.
19. Bikle D, Bouillon R, Thadhani R, et al. 2017 Vitamin D metabolites in captivity? Should we measure free or total 25(OH)D to assess vitamin D status? *J Steroid Biochem Mol Biol.* In press.
20. Powe CE, Ricciardi C, Berg AH, et al. 2011 Vitamin D-binding protein modifies the vitamin D-bone mineral density relationship. *J Bone Miner Res* 26(7):1609–1616.

21. Johnsen MS, Grimnes G, Figenschau Y, et al. 2014 Serum free and bio-available 25-hydroxyvitamin D correlate better with bone density than serum total 25-hydroxyvitamin D. *Scand J Clin Lab Invest* 74(3):177–183.
22. Alwan A, Rizkallah M, Maalouf G, et al. 2018. Positive correlations between free vitamin D and bone variables in a group of young Lebanese women. *J Clin Densitom.* In press.

Accepted Manuscript