Urbanization-driven changes in web-building and body size in an orb-web spider

Maxime Dahirel, Maarten de Cock, Pieter Vantieghem, Dries Bonte

► To cite this version:

Maxime Dahirel, Maarten de Cock, Pieter Vantieghem, Dries Bonte. Urbanization-driven changes in web-building and body size in an orb-web spider. *Journal of Animal Ecology*, 2019, 88 (1), pp.79-91. 10.1111/1365-2656.12909 . hal-01903122

HAL Id: hal-01903122

<https://hal-univ-rennes1.archives-ouvertes.fr/hal-01903122>

Submitted on 9 Nov 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Journal of Animal Ecology

MR MAXIME DAHIREL (Orcid ID : 0000-0001-8077-7765)

DR DRIES BONTE (Orcid ID : 0000-0002-3320-7505)

Article type : Research Article

Editor : Niels Dingemanse

Urbanization-driven changes in web-building and body size in an orb-web spider

Maxime Dahirel^{1,2*}, Maarten De Cock¹, Pieter Vantieghem¹ and Dries Bonte¹

¹ Ghent University, Department of Biology, Terrestrial Ecology unit, K. L. Ledeganckstraat
35, B-9000 Gent, Belgium

² Univ Rennes, CNRS, ECOBIO (Ecosystèmes, biodiversité, évolution) - UMR 6553, F-
35000 Rennes, France

* corresponding author: maxime.dahirel@yahoo.fr

Abstract

1. In animals, behavioural responses may play an important role in determining population persistence in the face of environmental changes. Body size is a key trait central to many life history traits and behaviours. Correlations with body size may constrain behavioural variation in response to environmental changes, especially when size itself is influenced by environmental conditions.

- Accepted Article
2. Urbanization is an important human-induced rapid environmental change that imposes multiple selection pressures on both body size and (size-constrained) behaviour. How these combine to shape behavioural responses of urban-dwelling species is unclear.
 3. Using web-building, an easily quantifiable behaviour linked to body size, and the garden spider *Araneus diadematus* as a model, we evaluated direct behavioural responses to urbanization and body size constraints across a network of 63 selected populations differing in urbanization intensity. We additionally studied urbanization at two spatial scales to account for some environmental pressures varying across scales and to obtain first qualitative insights about the role of plasticity and genetic selection.
 4. Spiders were smaller in highly urbanized sites (local scale only), in line with expectations based on reduced prey biomass availability and the Urban Heat Island effect. Web surface and mesh width decreased with urbanization at the local scale, while web surface also increased with urbanization at the landscape scale. The latter two responses are expected to compensate, at least in part, for reduced prey biomass availability in cities. The use of multivariate mixed modelling reveals that although web traits and body size are correlated within populations, behavioural responses to urbanization do not appear to be constrained by size: there is no evidence of size-web correlations among populations or among landscapes, and web traits appear independent from each other.
 5. Our results demonstrate that responses in size-dependent behaviours may be decoupled from size changes, thereby allowing fitness maximisation in novel environments. The spatial scale at which traits respond suggests contributions of both genetic adaptation (for web investment) and plasticity (for mesh width). Although fecundity decreased with local-scale urbanization, *Araneus diadematus* abundances

were similar across urbanization gradients; behavioural responses thus appear overall successful at the population level.

Key words: adaptation, fecundity, foraging, multivariate mixed model, spider web, temperature-size rule

Introduction

In animals, behaviour is often considered as the first route of adaptation to rapid environmental changes (Wong & Candolin, 2015). Numerous examples of both adaptive and maladaptive behavioural changes in response to human-induced environmental changes have now been recorded (Lowry, Lill, & Wong, 2013); the potential costs and constraints associated with these behavioural changes are, however, poorly understood. Behaviours may be linked to metabolic and physiological processes that directly impact fitness (Bonte et al., 2012; Debecker, Sanmartín-Villar, de Guinea-Luengo, Cordero-Rivera, & Stoks, 2016). When behaviour is correlated to other traits, conflicting selection pressures may hinder adaptation, leading to mismatches between the expressed and optimal behaviours in the new environment (Sih, 2013; Wong & Candolin, 2015). In particular, many behaviours are correlated to body size (e.g. Gregorič, Kuntner, & Blackledge, 2015; Stevens et al., 2014), a key trait that can itself be directly impacted by environmental changes (Oliveira, Freitas, Scheper, & Kleijn, 2016; Renault, Hutchinson, Loeb, Poveda, & Connelly, 2016).

Urbanization is one of the most prominent human-induced environmental changes, with cities now harbouring more than half of the global human population (Seto, Güneralp, & Hutyrá, 2012; United Nations Population Division, 2015). Direct and indirect impacts of the urbanization process include habitat fragmentation, increased temperatures (the “Urban Heat

Island” effect), elevated levels of pollution and changes in resource availability due to the decline of key species and/or the increased availability of anthropogenic food sources (Alberti, 2015; Parris, 2016). This cocktail of environmental changes means cities present novel ecological conditions never encountered before in organisms’ evolutionary histories (Alberti, 2015; Hendry, Gotanda, & Svensson, 2017; Johnson & Munshi-South, 2017). This drives changes in community taxonomic and functional composition (Dahirel, Dierick, De Cock, & Bonte, 2017; Piano et al., 2017) as well as intraspecific phenotypic changes (plastic and/or genetic: Alberti et al., 2017; Brans, Jansen, et al., 2017; Johnson & Munshi-South, 2017; Lowry et al., 2013) at an unprecedented rate (Alberti et al., 2017). Cities can therefore be seen as key “natural experiments” to understand eco-evolutionary consequences of global change (Johnson & Munshi-South, 2017).

Orb web spiders (Araneae, Araneidae) are a unique model for the study of behaviour, as their foraging behaviour is archived in their web (Foelix, 2010; Fig. 1). Detailed quantification of the orb web structure therefore allows inferences on both the implemented foraging strategies (from web architecture) and the energetic investments in web production (Sherman, 1994). Keeping total investment equal, larger spacing of the capture threads allows the construction of larger webs and an increase in prey interception; conversely, pressures to build small meshes to e.g. increase prey retention (Blackledge & Zevenbergen, 2006) come at the cost of reduced capture areas (Sandoval, 1994). Spiders can maximise both retention and interception simultaneously only by increasing the quantity of capture silk per web (Eberhard, 2013; Sherman, 1994). Variation in web building behaviour has been recorded in relation to aging and growth (e.g. Witt, Rawlings, & Reed, 1972), and in response to a wide variety of environmental changes. Spiders experiencing lower prey biomass have for instance been shown to increase the capture area of their web to increase its efficiency (Mayntz, Toft, & Vollrath, 2009). Spiders may also use information on the size of recently caught prey to

optimize the web mesh width and capture area (Schneider & Vollrath, 1998). The architecture and size of webs are influenced by weather, both through changes in the physical pressures the web has to withstand and through changes in the internal physiological state of the building spider (Barghusen, Claussen, Anderson, & Bailer, 1997; Vollrath, Downes, & Krackow, 1997). Web building can also be disturbed by exposure to various chemical pollutants (Benamú, Schneider, González, & Sánchez, 2013), and constrained by space availability in the current habitat (Vollrath et al., 1997). Most of these environmental drivers are highly altered by urbanization (Dahirel et al., 2017; Parris, 2016) and may either impose constraining or adaptive changes in web building.

Besides these direct responses, environmental changes may also influence spider foraging behaviour indirectly, as web-building is coupled to body size both within and among species.

Larger spiders produce on average larger, and often less densely structured webs (Gregorič et al., 2015; Heiling & Herberstein, 1998). This may stem from physiological constraints such as the quantity of silk available or the need for larger spiders to capture larger prey to fulfil energy requirements), as well as physical constraints, as spiders seem to use their own body as a “measurement stick” during web building (Gregorič, Kuntner, & Blackledge, 2015; Venner, Bel-Venner, Pasquet, & Leborgne, 2003, Vollrath, 1987). Increased temperatures in urban areas are expected to lead to reduced body size in ectotherms through increases in metabolic rates (Brans, Jansen, et al., 2017; Horne, Hirst, & Atkinson, 2015; Sheridan & Bickford, 2011). We may expect urbanization-induced body size reduction to be even stronger when urbanization decreases resource availability during development (Dahirel et al., 2017), as spider adult size is also highly resource-dependent (Mayntz, Toft, & Vollrath, 2003). This expected size reduction may thus constrain optimal web-building behaviour, for instance limiting the ability of spiders to increase web size to catch more prey when needed.

Behavioural responses to urbanization and other human-induced rapid environmental changes, whether adaptive or not, can result from plasticity and/or genetic changes (Sih, Stamps, Yang, McElreath, & Ramenofsky, 2010; Tuomainen & Candolin, 2011). A formal understanding of the underlying sources of phenotypic (behavioural) variation in response to urbanization can only be achieved by experimental work (Johnson & Munshi-South, 2017; Merilä & Hendry, 2014). However, the spatial scale of phenotypic variation, relative to the scale of dispersal, may provide first insights. Spiders disperse passively by “ballooning” (Bonte, 2012); although very long-distance dispersal events occur, most dispersers are thought to land within a few hundred metres from their departure point (Bell, Bohan, Shaw, & Weyman, 2005; Croucher, Oxford, & Gillespie, 2011; Reed, Teoh, Stratton, & Hataway, 2011; Reynolds, Bohan, & Bell, 2007). Once airborne, individuals have no control over their trajectory, meaning adaptive habitat choice (Edelaar & Bolnick, 2012) is unlikely. Gene flow is thus expected to swamp genetic differentiation at small spatial scales; any response to environmental variation with a spatial grain equal or smaller than the typical dispersal event is more likely to result from plasticity. By contrast, evidence of phenotype-environment matching at larger spatial scales (several kilometres) suggests putative genetic adaptation (Macdonald, Llewelyn, & Phillips, 2018; Richardson, Urban, Bolnick, & Skelly, 2014).

Here, we study and quantify shifts in body size and web building in response to urbanization, then use a multivariate mixed modelling approach in order to identify the contribution of size constraints to (adaptive) behavioural variation. We used the garden spider *Araneus diadematus* as a model; this species is one of the most common species in both urban and non-urban orb web spider communities in western Europe (Dahirel et al., 2017). *A. diadematus* adjusts its web-building behaviour depending on abiotic conditions and the availability/ characteristics of potential prey (Bonte, Lanckacker, Wiersma, & Lens, 2008; Schneider & Vollrath, 1998; Vollrath et al., 1997); webs are recycled and rebuilt daily,

allowing spiders to match currently/recently experienced environmental conditions (Breed, Levine, Peakall, & Witt, 1964).

We studied responses to urbanization at two independent spatial scales to obtain indications on the roles of genetic adaptation vs. plasticity, and to account for the fact that environmental correlates of urbanization may be scale-dependent (Kaiser et al., 2016; McDonnell & Hahs, 2015; Merckx et al., 2018). We used a well-studied network of urban and non-urban sites in which differences in resource availability and temperature have been previously recorded (Dahirel et al., 2017; Kaiser, Merckx, & Van Dyck, 2016; Merckx et al., 2018). Although prey numbers are roughly constant across urbanization levels, prey biomass is expected to be lower in highly urbanized sites as the size of the average prey declines by about 18% (Dahirel et al., 2017). Average temperatures are higher in highly urbanized sites, but this effect is mainly detected when considering urbanization at the most local scale (about +1-2°C) and weaker to non-significant at larger spatial scales (Merckx et al., 2018).

In this context, we expected *Araneus diadematus* individuals to become smaller with urbanization, and the decrease in size to be stronger with local-scale urbanization as the effects of resource reduction and increased temperature cumulate. We expected *Araneus diadematus* to present adaptive shifts in web-building behaviour in response to urbanization-related resource reduction, namely an increase in web investment/ web area to increase interception and a narrowing of mesh width to increase retention. If spiders were unable to display such adaptive behavioural responses, we expected to see negative fitness consequences through lower individual fecundity and/or population success, measured by spider density. Assuming correlations between web traits and body size were important and persisted across urbanization contexts (which may not be the case; Fischer, Ghalambor, & Hoke, 2016; Peiman & Robinson, 2017), we expected the mesh reduction response to be facilitated by reduced body size, while a smaller body size should have hindered spiders'

ability to build larger webs. Mesh width and web area responses should thus be stronger at the scale at which web trait and body size are the most/ the least strongly correlated, respectively.

Material and Methods

Study species

Araneus diadematus Clerck 1757 is a common orb-weaving spider present across the Holarctic in a wide range of natural and human-altered environments, building its web in shrubs and tall herbaceous vegetation (Lee & Thomas, 2002; Nentwig, Blick, Gloor, Hänggi, & Kropf, 2016). Its distinctive dorsal cross pattern makes field identification easy (Roberts, 1993; Fig. 1). Females usually become mature in late summer, and can survive through to late autumn (Lee & Thomas, 2002). In cities, *Araneus diadematus* mostly settles in gardens and green spaces as opposed to roadsides or close to buildings (Van Keer, Vanuytven, De Koninck, & Van Keer, 2010).

Study sites

We sampled *A. diadematus* females in 63 sites (hereafter “populations”) across a well-studied network of urban, rural and natural landscapes in northern Belgium (Brans, Govaert, et al., 2017; Dahirel et al., 2017; Kaiser et al., 2016; Merckx et al., 2018; Piano et al., 2017; Fig. 2), one of the most urbanized and densely populated regions in Europe (United Nations Population Division, 2015). Urbanization was studied at two different spatial scales thanks to a 2-step stratified selection design, with 3×3 km landscapes (“landscape scale” urbanization) being selected first, then three 200×200 m sites (“local scale” urbanization) chosen within each landscape. We used the percentage of surfaces occupied by buildings as our proxy for

urbanization (extracted from the Large-scale Reference Database, a reference map of Flanders; <https://www.agiv.be/international/en/products/grb-en>). This metric is precise down to the individual building, and thus usable at both spatial scales, but excludes other artificialized surfaces such as roads, railways, parking spaces or pavements; hence percentages higher than 10% already correspond to highly urbanized contexts. We cross-checked this metric with the percentage of artificial surfaces based on the CORINE land cover database (European Environmental Agency, 2016)(Level 1: “Artificial surfaces”, excluding classes 1.41 and 1.42 corresponding to human-created urban green spaces). This second metric includes all artificialized surfaces but, due to its coarser resolution (minimum mapping unit 25 ha) cannot be used at smaller spatial scales. Both metrics were highly correlated ($N = 21$ landscapes, Spearman correlation coefficient = 0.95, $p = 5.17 \times 10^{-6}$, Supplementary Figure S1). Twenty-one 3×3 km landscapes were selected and sorted into three urbanization levels (7 landscapes by level). High-urbanization landscapes had more than 10% of their area covered by buildings (equivalent to more than 70% of artificial surfaces based on CORINE). Low-urbanization landscapes had less than 3% of their surfaces occupied by buildings (less than 20% of artificial surfaces based on CORINE); additionally, they were selected so that more than 20% of their surface was covered by so-called “ecologically valuable areas” (areas with rare, vulnerable or highly diverse vegetation based on the Flanders Biological Valuation Map; Vriens et al., 2011). A third class of “intermediate” landscapes had between 5 and 10% of their surface covered by buildings (between 20 and 70% of artificial surfaces based on CORINE). Within each landscape, three 200×200 m sites (one per urbanization level) were selected this time based on the percentage covered by buildings only. Vegetated areas in chosen sites were unforested and dominated by low vegetation (grasslands/lawns with low shrubs and occasional trees in e.g. gardens, parks or hedgerows). Sites were separated on average by 39.4 km (range: 0.2 to 91.3 km, SD: 20.7 km). Sites within the same landscape

were separated on average by 1.4 km (range 0.2 to 3.3 km, SD: 0.8 km), while sites belonging to different landscapes were separated on average by 40.33 km (range: 5.6 to 91.3 km, SD: 19.75 km, Fig. 2).

Spider collection and phenotypic measurements

Populations were sampled from 25 August to 5 October 2014, i.e. during the first part of the reproductive period (Lee & Thomas, 2002). One landscape (3 sites) was visited per day; there was no significant link between landscape-level urbanization and sampling date (ANOVA; $N = 21$ landscapes, $F_{2,18} = 0.009$, $p = 0.991$). In each visited site, between 7 and 11 adult females per population ($N_{total} = 621$, average $\pm$ SD : 9.86 ± 0.74) were sampled on their webs and stored in 70% ethanol; spiders' cephalothorax width was measured under binocular microscope and used as a proxy for body size (Bonte et al., 2008; Fig. 1). Out of these 621 spiders, 193 individuals caught in the 9 landscapes of the Ghent region (Fig. 2) were also dissected and the number of mature eggs recorded. As *Araneus diadematus* webs are conspicuous and located in similar habitats (grassland-shrub mosaics taken in the broad sense) independently of urbanization levels (Lee & Thomas, 2002; Van Keer et al., 2010), we were also able to collect reliable population density data (number of adult female spiders observed per 200×200 m site in 4.5 person-hours) in the 62 sites (out of 63) sampled in the community-level study by Dahirel *et al.* (2017).

Based on measurements taken in the field (vertical and horizontal diameters of whole web and free central zone, number of sticky silk spirals in each web quadrant, Fig. 1), we estimated two design parameters for the webs belonging to sampled spiders: the web capture area surface (considering orb webs as ellipses, following Herberstein & Tso, 2000), and the mesh width (interval between sticky spirals, averaged over the horizontal and vertical axes).

Statistical analysis

All analyses were carried out using R, version 3.5 (R Core Team, 2018). Body size, web traits and sampling date values were scaled to unit standard deviation prior to inclusion in models (see Table 1 for descriptive statistics), in order to improve model convergence and make results comparable across parameters and variables.

The number of adult spiders per site was analysed using a Poisson generalized linear mixed model. Spider numbers were modelled as a function of local and landscape level of urbanization, as well as sampling date, with a random effect of landscape identity added to account for spatial clustering of populations in landscapes.

We followed broadly Araya-Ajoy and Dingemanse (2014)'s approach and analysed spider traits and their correlations in two steps.

First, each trait (body size, web surface, mesh width, egg number) was analysed separately using univariate mixed models. These models included fixed effects of local and landscape level of urbanization as well as sampling date, with random effects of landscape and site of origin to account for non-independence of observations. Body size, web surface and mesh width were analysed using linear mixed models, egg number was analysed using a Poisson generalized linear mixed model.

Second, we used a multivariate mixed model (Araya-Ajoy & Dingemanse, 2014; Dingemanse & Dochtermann, 2013) to estimate trait covariances/correlations and partition them across hierarchical levels (correlations among landscapes, among populations, and (residual) within populations). Fixed effects, random intercepts and error families for each trait were included as in univariate models.

All models were fitted using the MCMCglmm R package (Hadfield, 2010). We used uninformative default priors for fixed effects, inverse-Wishart priors for the residual (co)variances and parameter expanded priors for among-landscapes and among-population (co)variance matrices. Degrees of belief ν for the (co)variance matrices were set to weakly informative values, i.e. the dimension of the corresponding matrix (1 for univariate models, 4 for the multivariate model). Chains ran for (univariate/ multivariate model) 1000000/ 3300000 iterations, with the first 250000/ 300000 iterations discarded as burn-in and sampling every 750/ 3000 iteration post burn-in, resulting in 1000 samples per chain with serial autocorrelation < 0.1 for all model parameters. We ran three chains per model, checked mixing graphically and confirmed chain convergence using the Gelman-Rubin statistic (using the coda R package; Plummer, Best, Cowles, & Vines, 2006; all $\hat{R}$ were ≤ 1.03). There was no evidence of residual spatial autocorrelation in any of the models; 95% confidence intervals of spline correlograms (fitted using the ncf package; Bjornstad, 2016) overlapped with 0 at all distances and for all response variables. Model diagnostic plots are presented in Supplementary Figures S2 to S30c.

Results

Spider population density

The number of spiders found per site was not influenced by urbanization at either spatial scale (95% Credible Intervals for all model coefficients β overlapped with 0, Table 2). There was also no temporal trend in spider abundance (Table 2).

Sources of variation in spider traits: univariate models (Table 2, Fig 3)

Body size, as measured by cephalothorax width, decreased with local-scale urbanization, with spiders being 8.30% smaller [95% Credible Interval: -14.03%, -2.34%] in highly urbanized sites compared to populations from low-urbanization sites. There was no effect of landscape-scale urbanization or sampling date.

Spiders built webs with narrower mesh when local scale urbanization increased (low to high urbanization: -9.67% [-14.96%, -4.07%]). Mesh width was not influenced by landscape-scale urbanization, but increased with sampling date. Web surface decreased with increasing urbanization at the local scale (low to intermediate urbanization: -12.93% [-23.83%, -1.62%]; low to high urbanization: -20.28% [-31.12%, -9.37%]). It increased with landscape-scale urbanization (low to intermediate urbanization: +26.19% [+2.25%, +54.63%]; low to high urbanization: +25.43% [+0.5%, +54.54%]) and with time.

Local urbanization (both intermediate and high levels) decreased fecundity (number of eggs per spider) at the local scale (low to intermediate urbanization: -9.64% [-14.51%, -4.77%]; low to high urbanization: -16.62% [-20.81%, -12.12%]). There was no supported effect of landscape-scale urbanization or sampling date on fecundity.

Correlations among traits: multivariate models

Inferences based on fixed effects from this multivariate model were virtually the same as those obtained from the univariate models (see Table 2 vs. Supplementary Table S1 for a comparison of parameters). At the landscape and population levels, none of the estimated trait covariances were different from zero, based on credible intervals (Table 3). At the within-population level, web traits were positively correlated with each other and with body size (Table 3).

Discussion

Using a standardized hierarchical sampling design and multivariate modelling, we found *Araneus diadematus* to decrease in size with increasing levels of local urbanization, and independently, to adjust different components of its web-building behaviour. Web-building changes, in particular the reduction of mesh width at the local scale, and the increase in web investment at the landscape scale, appear to be adaptive (i.e. beneficial) and follow predictions based on lower prey biomass in cities. The putative adaptive value of these behavioural changes is supported by the fact that *A. diadematus* is able to maintain consistent population densities across broad gradients of urbanization, when other orb web spider species in the same communities cannot (Dahirel et al., 2017). An alternative explanation is that urban habitats are population sinks that are kept afloat by a supply of immigrant ballooning spiders from natural habitats (Parris, 2016). While we have no reason to anticipate such an asymmetry in dispersal, we should then expect landscape-scale urbanization to decrease local population density, as it would limit this supply. This is contrary to our observations. Interestingly, while web building behaviours are intrinsically connected to body size, as shown by within-population correlations, their responses to urbanization are decoupled (no among-populations or among-landscapes correlations).

Araneus diadematus experienced size reduction (by about 8%) in response to local urbanization only. The temperature-size rule, found in many arthropods, predicts that individuals should be smaller at higher temperatures (Atkinson, 1994; Horne et al., 2015). As the spatial scale of urban heating matches the spatial scale at which size reduction is observed (Kaiser et al., 2016; Merckx et al., 2018), the Urban Heat Island Effect provides a valid explanation for the observed patterns (Merckx 2018, but see Entling, Schmidt-Entling, Bacher, Brandl, & Nentwig, 2010; Puzin, Leroy, & Pétillon, 2014). However, the impact of small temperature changes on spider body size remains to be studied, and resource availability

Accepted Article

during development is a well-documented influence and a potentially stronger driver of spider adult size (DiRienzo & Montiglio, 2016; Kralj-Fišer et al., 2014; Mayntz et al., 2003). Higher prey abundance in urbanized spots of the Sydney area was for instance associated with a larger body size in the spider *Nephila plumipes* (Lowe et al., 2014; Lowe, Wilder, & Hochuli, 2016). In our study region, prey size and prey biomass availability decreased with urbanization (by about 18%; Dahirel et al., 2017). The among-studies discrepancy in spider body size shifts cannot be easily reconciled with a major role of the Urban Heat Island effect, but is fully consistent with a prominent role of resource availability. Interestingly, we did not detect a spider size reduction in response to landscape-level urbanization (Table 2, Fig. 3), even though prey body size declines equally in response to urbanization at both scales (Dahirel et al., 2017). This result is in accordance with our prediction that size reduction should be stronger at the spatial scale at which more urbanization-induced environmental changes cumulate, i.e. the local scale. Pollutants such as heavy metals also have a negative impact on body size (Ramirez et al., 2011), but we expect their influence here to be limited, as pesticide and heavy metal levels appear unrelated to urbanization in our study area (Merckx et al., 2018; T. Merckx, pers. comm.). Alternatively, resource loss may be the main mechanism driving size-reduction with urbanization, which is only observed at the local scale because spiders are able to increase their investment to compensate for prey biomass loss at the landscape, but not at the local scale.

Changes in foraging behaviour in response to urbanization were scale-dependent: spiders built smaller webs with smaller mesh width in response to local urbanization, but increased their web surface with urbanization at the landscape scale (Fig. 3). Additionally, web surface responded to urbanization at both scales independently of mesh width variation (surface – mesh correlation ≈ 0 ; Table 3), meaning that web surface changes can tentatively be interpreted in terms of corresponding silk investment changes. Webs with smaller mesh are

considered better at retaining prey (Blackledge & Zevenbergen, 2006), at the cost of a smaller web surface and therefore fewer prey intercepted for a given investment in silk production. When mesh width is held constant, as it is in response to landscape-level urbanization, increased silk investment leads to an increase in web surface (Figs 1, 3) and therefore an increase in the number of prey caught (Prokop & Grygláková, 2005; Venner & Casas, 2005). Because of their larger diameter, larger webs are also built with longer radial silk threads, meaning they can stop prey with higher kinetic energy (i.e. bigger prey) without breaking down (Harmer, Clausen, Wroe, & Madin, 2015). Although the importance of these large prey in particular for orb web evolution (Blackledge, 2011; Eberhard, 2013; Harmer et al., 2015) and individual performance (Harmer et al., 2015; Venner & Casas, 2005) remains controversial, selective advantages to having webs that maximize prey biomass in general are clear (Harmer et al., 2015). Temperature can also have a positive effect on silk production and hence web surface (Barghusen et al., 1997; Vollrath et al., 1997). However, the Urban Heat Island effect is stronger at the local, rather than landscape scale in our study region (Kaiser et al., 2016; Merckx et al., 2018), and is too weak (about 1-2°C) to explain the observed increase in web surface (in *Araneus diadematus*, a 12°C increase was needed to obtain surface changes similar to those we observed; Vollrath et al., 1997). Prey size (and thus biomass) decreased with urbanization at both scales (Dahirel et al., 2017). In this context, both the reduction of mesh size at local scales and the increase in capture area at larger spatial scales can be seen, in accordance with our predictions, as adaptive responses to urbanization, respectively increasing prey retention and prey capture efficiency. These adaptations potentially contribute to the persistence of *Araneus diadematus* across urbanization gradients, and confirm that biotic interactions can be important drivers of phenotypic changes in urban environments (Alberti et al., 2017).

Web-building in orb-weaving spiders has previously been shown to be both size-constrained (Bonte et al., 2008; Gregorič et al., 2015), but also highly variable depending on the environment (Herberstein & Tso, 2011); the net effect in the case of environmental changes influencing body size was so far unknown. We show that correlations among our studied traits are only present within populations (residual covariance matrix) and not at the among-population and among-landscape levels (Table 3), implying that traits respond independently to urbanization. This must be interpreted with some caution however, as our ability to accurately detect weak among-population correlations ($|r| < 0.3$) may be limited (Dingemanse & Dochtermann, 2013). Within-population (phenotypic) correlations likely result from physiological and possibly physical constraints on web-building, with larger spiders having more reserves to build large webs and spiders using their own body as measurement during web building (Gregorič, Kuntner, & Blackledge, 2015; Venner, Bel-Venner, Pasquet, & Leborgne, 2003; Vollrath, 1987). They may also result from small scale within-site environmental variation influencing both webs and body size independently, for instance if web-building micro-habitats leading to smaller webs for structural reasons (Blamires, Thompson, & Hochuli, 2007; Vollrath, Downes, & Krackow, 1997) are also sites poorer in prey. In any case, our results suggest that while body size is correlated with web-building strategy within populations, it does not substantially limit the possibilities of spiders to adapt their web-building behaviour in response to urbanization. The observed changes in web building are thus a direct response to urban selection pressures, and not an indirect consequence of body size shifts. Additionally, these results confirm that the existence of trait correlations/syndromes in one environmental context or at one biological organisation level may not be informative with respect to their maintenance across contexts/ levels (e.g. Fischer, Ghalambor, & Hoke, 2016; Peiman & Robinson, 2017).

Changes in web-building behaviour in relation to urbanization may both originate from plasticity and/or evolutionary changes (Herberstein & Tso, 2011). Disentangling the relative contribution of both is however difficult in observational studies (Merilä & Hendry, 2014). We take advantage of our two-spatial scales design, and the fact that plasticity and genetic adaptation are generally expected to appear in response to finer-grained and coarser-grained environmental variation, respectively (Richardson et al., 2014), to emit hypotheses on the relative contributions of these two mechanisms. Mesh width varies with urbanization at the local but not at the landscape scale. This would indicate a more important role of plastic responses, in accordance with experimental work demonstrating high within-individual behavioural flexibility in response to food availability and prey spectrum (reviewed by Herberstein & Tso, 2011). By contrast, silk investment, reflected in web surface, adaptively increased in response to landscape-scale urbanization. With the caveat that some drivers of web building variation may have been overlooked, this suggests here a putative role of genetic adaptation in urbanization-driven silk production changes. We acknowledge our results only provide first hints in this direction, and that further research will be needed to quantify the contribution of genetic selection relative to developmental and behavioural plasticity.

Costs associated with urban life were especially prominent at the local scale. Indeed, at this scale, body sizes and web capture surfaces were smaller, and lower fecundity was observed despite stable population densities (Fig. 3, Table 2). Egg load did not, however, covary with body size or web traits at either the inter- or intra-population levels. Reduced fecundity is therefore unlikely to be a direct consequence of increased energetic investment in webs or a reduced body size, but rather an additive response to environmental changes associated with urbanization. Food limitation has strong negative impacts on spider lifetime fecundity even in the absence of survival costs (Kleinteich, Wilder, & Schneider, 2015) or body size changes (Miyashita, 1990). The lower fecundities observed in locally urbanized sites may therefore

indicate that spiders were unable to fully compensate for reduced prey biomass at this scale despite shifts in web-building, and despite smaller size leading to reduced requirements. Other –here unmeasured– stressors related to urbanization may also lead to reduced fecundity and capture area in locally urbanized sites: heat waves (Kingsolver, Diamond, & Buckley, 2013), pollution (fecundity costs: Hendrickx, Maelfait, Speelmans, & Straalen, 2003; web surface costs: Benamú et al., 2013; but see Ramirez et al., 2011), or repeated web destructions and rebuilding following human disturbance (Tew, Adamson, & Hesselberg, 2015). Overall, *Araneus diadematus* was comparatively much better at dealing with landscape-scale urbanization; indeed, both fecundity and population densities remained unaffected by urbanization at this scale. This suggests that traits that successfully responded at this scale (namely web surface) are more important to prey capture success and biomass gain than the others (mesh width) (Blackledge & Eliason, 2007). Conversely, benefits, costs and trade-offs associated with increased silk production may be detectable in other life-history dimensions than the ones we explored in the present study, such as development time and adult longevity (both expected to increase with resource restriction; Kleinteich et al., 2015; Kralj-Fišer et al., 2014), or dispersal, which is strongly counter-selected in urban fragmented environments for passively dispersing organisms (Cheptou, Carrue, Rouifed, & Cantarel, 2008).

From an applied perspective, our results confirm the oft-stated importance of maintaining local green spaces in highly urbanized landscapes (e.g. Philpott et al., 2013), as even generalist “winning” species such as *A. diadematus* may benefit from them. Additionally, even though spider abundance stayed constant, webs were up to 20% smaller in locally urbanized sites. Spiders are important predators in natural and anthropogenic landscapes (Birkhofer, Entling, & Lubin, 2013; Foelix, 2010): our results highlight the necessity of accounting for changes in functional, and not only numerical predator responses to accurately

quantify ecosystem services provided by green infrastructures in urban environments. Further comparisons among successful species that differ in their silk production strategy (e.g. *A. diadematus*, which destroys and recreates webs regularly, versus *Nephila plumipes*, which build semi-permanent webs; Lowe et al., 2014) may highlight commonalities and help shed further light on the costs/ benefits balance of adaptation to urban life.

Authors' contributions

DB conceived the study and designed methodology; MDC and PV collected the data; MD and MDC analysed the data; MD led the writing of the manuscript. All authors contributed critically to the draft and gave final approval for publication.

Acknowledgments

We are grateful to Jasper Dierick for his assistance during field sampling, and to Hans Matheve for GIS calculations and site selection. We thank Niels Dingemanse and four anonymous reviewers for their comments on previous versions of this paper. This article is part of the Belspo-funded IAP project P7/04 SPEEDY (SPatial and environmental determinants of Eco-Evolutionary DYnamics: anthropogenic environments as a model). MD is funded by a postdoctoral grant from the Fyssen Foundation.

Data accessibility

Data and R code are available from the Dryad Digital Repository: <https://doi.org/10.5061/dryad.j886dg0> (Dahirel et al., 2018).

References

- Alberti, M. (2015). Eco-evolutionary dynamics in an urbanizing planet. *Trends in Ecology & Evolution*, 30(2), 114–126. doi:10.1016/j.tree.2014.11.007
- Alberti, M., Correa, C., Marzluff, J. M., Hendry, A. P., Palkovacs, E. P., Gotanda, K. M., ... Zhou, Y. (2017). Global urban signatures of phenotypic change in animal and plant populations. *Proceedings of the National Academy of Sciences*, 201606034. doi:10.1073/pnas.1606034114
- Araya-Ajoy, Y. G., & Dingemanse, N. J. (2014). Characterizing behavioural ‘characters’: an evolutionary framework. *Proceedings of the Royal Society of London B: Biological Sciences*, 281(1776), 20132645. doi:10.1098/rspb.2013.2645
- Atkinson, D. (1994). Temperature and organism size—a biological law for ectotherms? *Advances in Ecological Research*, 25, 1–58. doi:10.1016/S0065-2504(08)60212-3
- Barghusen, L. E., Claussen, D. L., Anderson, M. S., & Bailer, A. J. (1997). The effects of temperature on the web-building behaviour of the common house spider, *Achaearanea tepidariorum*. *Functional Ecology*, 11(1), 4–10. doi:10.1046/j.1365-2435.1997.00040.x
- Bell, J. R., Bohan, D. A., Shaw, E. M., & Weyman, G. S. (2005). Ballooning dispersal using silk: world fauna, phylogenies, genetics and models. *Bulletin of Entomological Research*, 95(02). doi:10.1079/BER2004350
- Benamú, M. A., Schneider, M. I., González, A., & Sánchez, N. E. (2013). Short and long-term effects of three neurotoxic insecticides on biological and behavioural attributes of the orb-web spider *Alpaida veniliae* (Araneae, Araneidae): implications for IPM programs. *Ecotoxicology*, 22(7), 1155–1164. doi:10.1007/s10646-013-1102-9
- Birkhofer, K., Entling, M. H., & Lubin, Y. (2013). Agroecology: Trait composition, spatial relationships, trophic interactions. In D. Penney (Ed.), *Spider Research in the 21st Century: Trends and Perspectives* (pp. 200–229). Manchester: Siri Scientific Press.
- Bjornstad, O. N. (2016). *R package ncf: Spatial Nonparametric Covariance Functions*. R. Retrieved from <https://CRAN.R-project.org/package=ncf>

- Blackledge, T. A. (2011). Prey capture in orb weaving spiders: are we using the best metric? *Journal of Arachnology*, 39(2), 205–210. doi:10.1636/Chi10-52.1
- Blackledge, T. A., & Eliason, C. M. (2007). Functionally independent components of prey capture are architecturally constrained in spider orb webs. *Biology Letters*, 3(5), 456–458. doi:10.1098/rsbl.2007.0218
- Blackledge, T. A., Kuntner, M., & Agnarsson, I. (2011). The form and function of spider orb webs: evolution from silk to ecosystems. In J. Casas (Ed.), *Advances in Insect Physiology* (Vol. 41, pp. 175–262). Academic Press. Retrieved from <http://www.sciencedirect.com/science/article/pii/B9780124159198000045>
- Blackledge, T. A., & Zevenbergen, J. M. (2006). Mesh width influences prey retention in spider orb webs. *Ethology*, 112(12), 1194–1201. doi:10.1111/j.1439-0310.2006.01277.x
- Blamires, S. J., Thompson, M. B., & Hochuli, D. F. (2007). Habitat selection and web plasticity by the orb spider *Argiope keyserlingi* (Argiopidae): Do they compromise foraging success for predator avoidance? *Austral Ecology*, 32(5), 551–563. doi:10.1111/j.1442-9993.2007.01727.x
- Bonte, D. (2012). Case study II: Spiders as a model in dispersal ecology and evolution. In J. Clobert, M. Baguette, T. G. Benton, & J. M. Bullock (Eds.), *Dispersal ecology and evolution* (pp. xxvi–xxviii). Oxford: Oxford University Press.
- Bonte, D., Lanckacker, K., Wiersma, E., & Lens, L. (2008). Web building flexibility of an orb-web spider in a heterogeneous agricultural landscape. *Ecography*, 31(5), 646–653. doi:10.1111/j.0906-7590.2008.05508.x
- Bonte, D., Van Dyck, H., Bullock, J. M., Coulon, A., Delgado, M., Gibbs, M., ... Travis, J. M. J. (2012). Costs of dispersal. *Biological Reviews*, 87(2), 290–312. doi:10.1111/j.1469-185X.2011.00201.x
- Brans, K. I., Govaert, L., Engelen, J. M. T., Gianuca, A. T., Souffreau, C., & Meester, L. D. (2017). Eco-evolutionary dynamics in urbanized landscapes: evolution, species sorting and the change in zooplankton body size along urbanization gradients. *Philosophical Transactions of the Royal Society of London B*, 372(1712), 20160030. doi:10.1098/rstb.2016.0030

- Brans, K. I., Jansen, M., Vanoverbeke, J., Tüzün, N., Stoks, R., & De Meester, L. (2017). The heat is on: Genetic adaptation to urbanization mediated by thermal tolerance and body size. *Global Change Biology*, 23(12), 5218–5227. doi:10.1111/gcb.13784
- Breed, A. L., Levine, V. D., Peakall, D. B., & Witt, P. N. (1964). The fate of the intact orb web of the spider *Araneus diadematus*. *Behaviour*, 232(1), 43–60.
- Cheptou, P.-O., Carrue, O., Rouifed, S., & Cantarel, A. (2008). Rapid evolution of seed dispersal in an urban environment in the weed *Crepis sancta*. *Proceedings of the National Academy of Sciences*, 105(10), 3796–3799. doi:10.1073/pnas.0708446105
- Croucher, P. J. P., Oxford, G. S., & Gillespie, R. G. (2011). Population structure and dispersal in a patchy landscape: nuclear and mitochondrial markers reveal area effects in the spider *Theridion californicum* (Araneae: Theridiidae). *Biological Journal of the Linnean Society*, 104(3), 600–620. doi:10.1111/j.1095-8312.2011.01734.x
- Dahirel, M., Dierick, J., De Cock, M., & Bonte, D. (2017). Intraspecific variation shapes community-level behavioral responses to urbanization in spiders. *Ecology*, 98(9), 2379–2390. doi:10.1002/ecy.1915
- Dahirel, M., De Cock, M., Vantieghem P. & Bonte, D. (2018). Data from: Urbanization-driven changes in web-building and body size in an orb-web spider. Dryad Digital Repository. doi:10.5061/dryad.j886dg0
- Debecker, S., Sanmartín-Villar, I., de Guinea-Luengo, M., Cordero-Rivera, A., & Stoks, R. (2016). Integrating the pace-of-life syndrome across species, sexes and individuals: covariation of life history and personality under pesticide exposure. *Journal of Animal Ecology*, 85(3), 726–738. doi:10.1111/1365-2656.12499
- Dingemanse, N. J., & Dochtermann, N. A. (2013). Quantifying individual variation in behaviour: mixed-effect modelling approaches. *Journal of Animal Ecology*, 82(1), 39–54. doi:10.1111/1365-2656.12013

- DiRienzo, N., & Montiglio, P.-O. (2016). The contribution of developmental experience vs. condition to life history, trait variation and individual differences. *Journal of Animal Ecology*, 85(4), 915–926. doi:10.1111/1365-2656.12512
- Eberhard, W. G. (2013). The rare large prey hypothesis for orb web evolution: a critique. *Journal of Arachnology*, 41(1), 76–80. doi:10.1636/B12-34.1
- Edelaar, P., & Bolnick, D. I. (2012). Non-random gene flow: an underappreciated force in evolution and ecology. *Trends in Ecology & Evolution*, 27(12), 659–665. doi:10.1016/j.tree.2012.07.009
- Entling, W., Schmidt-Entling, M. H., Bacher, S., Brandl, R., & Nentwig, W. (2010). Body size–climate relationships of European spiders. *Journal of Biogeography*, 37(3), 477–485. doi:10.1111/j.1365-2699.2009.02216.x
- European Environmental Agency. (2016). *Corine Land Cover, 2012, Version 18.5.1*. European Union, Copernicus Land Monitoring Service. Retrieved from <https://land.copernicus.eu/pan-european/corine-land-cover/clc-2012>
- Fischer, E. K., Ghalambor, C. K., & Hoke, K. L. (2016). Plasticity and evolution in correlated suites of traits. *Journal of Evolutionary Biology*, 29(5), 991–1002. doi:10.1111/jeb.12839
- Foelix, R. (2010). *Biology of Spiders* (3rd ed.). Oxford, UK: Oxford University Press.
- Gregorič, M., Kuntner, M., & Blackledge, T. A. (2015). Does body size predict foraging effort? Patterns of material investment in spider orb webs. *Journal of Zoology*, 296(1), 67–78. doi:10.1111/jzo.12219
- Hadfield, J. D. (2010). MCMC methods for multi-response generalized linear mixed models: the MCMCglmm R package. *Journal of Statistical Software*, 33(2). doi:10.18637/jss.v033.i02
- Harmer, A. M. T., Clausen, P. D., Wroe, S., & Madin, J. S. (2015). Large orb-webs adapted to maximise total biomass not rare, large prey. *Scientific Reports*, 5. doi:10.1038/srep14121
- Heiling, A. M., & Herberstein, M. E. (1998). The web of *Nuctenea sclopetaria* (Araneae, Araneidae): relationship between body size and web design. *The Journal of Arachnology*, 26(1), 91–96.
- Hendrickx, F., Maelfait, J.-P., Speelmans, M., & Straalen, N. M. V. (2003). Adaptive reproductive variation along a pollution gradient in a wolf spider. *Oecologia*, 134(2), 189–194. doi:10.1007/s00442-002-1031-4

- Hendry, A. P., Gotanda, K. M., & Svensson, E. I. (2017). Human influences on evolution, and the ecological and societal consequences. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 372(1712), 20160028. doi:10.1098/rstb.2016.0028
- Herberstein, Marie E., & Tso, I.-M. (2000). Evaluation of formulae to estimate the capture area and mesh height of orb webs (Araneoidea, Araneae). *Journal of Arachnology*, 28(2), 180–184. doi:10.1636/0161-8202(2000)028[0180:EOFTET]2.0.CO;2
- Herberstein, Marie E., & Tso, I.-M. (2011). Spider webs: evolution, diversity and plasticity. In Marie Elisabeth Herberstein (Ed.), *Spider Behaviour: Flexibility and Versatility* (pp. 57–98). Cambridge, UK: Cambridge University Press. doi:10.1017/CBO9780511974496.004
- Horne, C. R., Hirst, A. G., & Atkinson, D. (2015). Temperature- size responses match latitudinal- size clines in arthropods, revealing critical differences between aquatic and terrestrial species. *Ecology Letters*, 18(4), 327–335. doi:10.1111/ele.12413
- Johnson, M. T. J., & Munshi-South, J. (2017). Evolution of life in urban environments. *Science*, 358(6363), eaam8327. doi:10.1126/science.aam8327
- Kaiser, A., Merckx, T., & Van Dyck, H. (2016). The Urban Heat Island and its spatial scale dependent impact on survival and development in butterflies of different thermal sensitivity. *Ecology and Evolution*, 6(12), 4129–4140. doi:10.1002/ece3.2166
- Kingsolver, J. G., Diamond, S. E., & Buckley, L. B. (2013). Heat stress and the fitness consequences of climate change for terrestrial ectotherms. *Functional Ecology*, 27(6), 1415–1423. doi:10.1111/1365-2435.12145
- Kleinteich, A., Wilder, S. M., & Schneider, J. M. (2015). Contributions of juvenile and adult diet to the lifetime reproductive success and lifespan of a spider. *Oikos*, 124(2), 130–138. doi:10.1111/oik.01421
- Kralj-Fišer, S., Čelik, T., Lokovšek, T., Šuen, K., Šiling, R., & Kuntner, M. (2014). Development and growth in synanthropic species: plasticity and constraints. *Naturwissenschaften*, 101(7), 565–575. doi:10.1007/s00114-014-1194-y

- Lee, P., & Thomas, T. (2002). *Araneus diadematus*. In P. R. Harvey, D. R. Nellist, & M. G. Telfer (Eds.), *Provisional atlas of British spiders (Arachnida, Araneae), volumes 1 and 2* (p. 225). Huntingdon, UK: Biological Records Centre.
- Lowe, E. C., Wilder, S. M., & Hochuli, D. F. (2016). Persistence and survival of the spider *Nephila plumipes* in cities: do increased prey resources drive the success of an urban exploiter? *Urban Ecosystems*, 19(2), 705–720. doi:10.1007/s11252-015-0518-9
- Lowe, Elizabeth C., Wilder, S. M., & Hochuli, D. F. (2014). Urbanisation at multiple scales is associated with larger size and higher fecundity of an orb-weaving spider. *PLoS ONE*, 9(8), e105480. doi:10.1371/journal.pone.0105480
- Lowry, H., Lill, A., & Wong, B. B. M. (2013). Behavioural responses of wildlife to urban environments. *Biological Reviews*, 88(3), 537–549. doi:10.1111/brv.12012
- Macdonald, S. L., Llewelyn, J., & Phillips, B. L. (2018). Using connectivity to identify climatic drivers of local adaptation. *Ecology Letters*, 21(2), 207–216. doi:10.1111/ele.12883
- Mayntz, D., Toft, S., & Vollrath, F. (2003). Effects of prey quality and availability on the life history of a trap-building predator. *Oikos*, 101(3), 631–638. doi:10.1034/j.1600-0706.2003.12408.x
- Mayntz, D., Toft, S., & Vollrath, F. (2009). Nutrient balance affects foraging behaviour of a trap-building predator. *Biology Letters*, 5(6), 735–738. doi:10.1098/rsbl.2009.0431
- McDonnell, M. J., & Hahs, A. K. (2015). Adaptation and adaptedness of organisms to urban environments. *Annual Review of Ecology, Evolution, and Systematics*, 46(1), 261–280. doi:10.1146/annurev-ecolsys-112414-054258
- Merckx, T., Souffreau, C., Kaiser, A., Baardsen, L., Backeljau, T., Bonte, D., ... Van Dyck, H. (2018). Body size shifts in aquatic and terrestrial urban communities. *Nature*, provisionally accepted.
- Merilä, J., & Hendry, A. P. (2014). Climate change, adaptation, and phenotypic plasticity: the problem and the evidence. *Evolutionary Applications*, 7(1), 1–14. doi:10.1111/eva.12137
- Miyashita, T. (1990). Decreased reproductive rate of the spider, *Nephila clavata*, inhabiting small woodlands in urban areas. *Ecological Research*, 5(3), 341–351. doi:10.1007/BF02347009

- Nakagawa, S., & Schielzeth, H. (2013). A general and simple method for obtaining R^2 from generalized linear mixed-effects models. *Methods in Ecology and Evolution*, 4(2), 133–142. doi:10.1111/j.2041-210x.2012.00261.x
- Nentwig, W., Blick, T., Gloor, D., Hänggi, A., & Kropf, C. (2016). Spiders of Europe. Retrieved November 17, 2016, from <http://www.araneae.unibe.ch>
- Oliveira, M. O., Freitas, B. M., Scheper, J., & Kleijn, D. (2016). Size and sex-dependent shrinkage of Dutch bees during one-and-a-half centuries of land-use change. *PLOS ONE*, 11(2), e0148983. doi:10.1371/journal.pone.0148983
- Parris, K. M. (2016). *Ecology of Urban Environments*. Chichester, UK: Wiley-Blackwell.
- Peiman, K. S., & Robinson, B. W. (2017). Comparative analyses of phenotypic trait covariation within and among populations. *The American Naturalist*, 190(4), 451–458. doi:10.1086/693482
- Philpott, S. M., Cotton, J., Bichier, P., Friedrich, R. L., Moorhead, L. C., Uno, S., & Valdez, M. (2013). Local and landscape drivers of arthropod abundance, richness, and trophic composition in urban habitats. *Urban Ecosystems*, 17(2), 513–532. doi:10.1007/s11252-013-0333-0
- Piano, E., De Wolf, K., Bona, F., Bonte, D., Bowler, D. E., Isaia, M., ... Hendrickx, F. (2017). Urbanization drives community shifts towards thermophilic and dispersive species at local and landscape scales. *Global Change Biology*, 23(7), 2554–2564. doi:10.1111/gcb.13606
- Plummer, M., Best, N., Cowles, K., & Vines, K. (2006). CODA: Convergence Diagnosis and Output Analysis for MCMC. *R News*, 6(1), 7–11.
- Prokop, P., & Grygláková, D. (2005). Factors affecting the foraging success of the wasp-like spider *Argiope bruennichi* (Araneae): role of web design. *Biologia*, 60(2), 165–169.
- Puzin, C., Leroy, B., & Pétillon, J. (2014). Intra- and inter-specific variation in size and habitus of two sibling spider species (Araneae: Lycosidae): taxonomic and biogeographic insights from sampling across Europe. *Biological Journal of the Linnean Society*, 113(1), 85–96. doi:10.1111/bij.12303

- R Core Team. (2018). R: a language and environment for statistical computing (Version 3.5.1). Vienna, Austria: R Foundation for Statistical Computing. Retrieved from <https://www.R-project.org/>
- Ramirez, M. G., McCallum, J. E. B., Landry, J. M., Vallin, V. A., Fukui, S. A., Gergus, H. E., ... Sy, C. L. (2011). Relationships between physiological characteristics and trace metal body burdens of banded garden spiders *Argiope trifasciata* (Araneae, Araneidae). *Ecotoxicology and Environmental Safety*, 74(4), 1081–1088. doi:10.1016/j.ecoenv.2011.02.003
- Reed, D. H., Teoh, V.-H., Stratton, G. E., & Hataway, R. A. (2011). Levels of gene flow among populations of a wolf spider in a recently fragmented habitat: current versus historical rates. *Conservation Genetics*, 12(1), 331–335. doi:10.1007/s10592-009-9995-9
- Renauld, M., Hutchinson, A., Loeb, G., Poveda, K., & Connelly, H. (2016). Landscape simplification constrains adult size in a native ground-nesting bee. *PLOS ONE*, 11(3), e0150946. doi:10.1371/journal.pone.0150946
- Reynolds, A. M., Bohan, D. A., & Bell, J. R. (2007). Ballooning dispersal in arthropod taxa: conditions at take-off. *Biology Letters*, 3(3), 237–240. doi:10.1098/rsbl.2007.0109
- Richardson, J. L., Urban, M. C., Bolnick, D. I., & Skelly, D. K. (2014). Microgeographic adaptation and the spatial scale of evolution. *Trends in Ecology & Evolution*, 29(3), 165–176. doi:10.1016/j.tree.2014.01.002
- Roberts, M. (1993). *The Spiders of Great Britain and Ireland* (Vols. 1–2). Stenstrup, Denmark: Apollo Books (original edition by Harley Books).
- Sandoval, C. P. (1994). Plasticity in web design in the spider *Parawixia bistriata*: a response to variable prey type. *Functional Ecology*, 8(6), 701–707. doi:10.2307/2390229
- Schneider, J. M., & Vollrath, F. (1998). The effect of prey type on the geometry of the capture web of *Araneus diadematus*. *Naturwissenschaften*, 85(8), 391–394. doi:10.1007/s001140050521
- Seto, K. C., Güneralp, B., & Hutyra, L. R. (2012). Global forecasts of urban expansion to 2030 and direct impacts on biodiversity and carbon pools. *Proceedings of the National Academy of Sciences*, 109(40), 16083–16088. doi:10.1073/pnas.1211658109

- Sheridan, J. A., & Bickford, D. (2011). Shrinking body size as an ecological response to climate change. *Nature Climate Change*, 1(8), 401–406. doi:10.1038/nclimate1259
- Sherman, P. M. (1994). The orb-web: an energetic and behavioural estimator of a spider's dynamic foraging and reproductive strategies. *Animal Behaviour*, 48(1), 19–34. doi:10.1006/anbe.1994.1208
- Sih, A. (2013). Understanding variation in behavioural responses to human-induced rapid environmental change: a conceptual overview. *Animal Behaviour*, 85(5), 1077–1088.
- Sih, A., Stamps, J., Yang, L. H., McElreath, R., & Ramenofsky, M. (2010). Behavior as a key component of integrative biology in a human-altered world. *Integrative and Comparative Biology*, 50(6), 934–944. doi:10.1093/icb/icq148
- Stevens, V. M., Whitmee, S., Le Galliard, J.-F., Clobert, J., Böhning-Gaese, K., Bonte, D., ... Baguette, M. (2014). A comparative analysis of dispersal syndromes in terrestrial and semi-terrestrial animals. *Ecology Letters*, 17(8), 1039–1052. doi:10.1111/ele.12303
- Tew, E. R., Adamson, A., & Hesselberg, T. (2015). The web repair behaviour of an orb spider. *Animal Behaviour*, 103, 137–146. doi:10.1016/j.anbehav.2015.02.016
- Tuomainen, U., & Candolin, U. (2011). Behavioural responses to human-induced environmental change. *Biological Reviews*, 86(3), 640–657. doi:10.1111/j.1469-185X.2010.00164.x
- United Nations Population Division. (2015). *World urbanization prospects: the 2014 revision*. New York, USA: United Nations, Department of Economic and Social Affairs.
- Van Keer, K., Vanuytven, H., De Koninck, H., & Van Keer, J. (2010). More than one third of the Belgian spider fauna (Araneae) found within the city of Antwerp: faunistics and some reflections on urban ecology. *Nieuwsbr. Belg. Arachnol. Ver.*, 25(2), 160.
- Venner, S., Bel-Venner, M.-C., Pasquet, A., & Leborgne, R. (2003). Body-mass-dependent cost of web-building behavior in an orb weaving spider, *Zygiella x-notata*. *Naturwissenschaften*, 90(6), 269–272. doi:10.1007/s00114-003-0420-9
- Venner, S., & Casas, J. (2005). Spider webs designed for rare but life-saving catches. *Proceedings of the Royal Society of London B: Biological Sciences*, 272(1572), 1587–1592. doi:10.1098/rspb.2005.3114

Vollrath, F. (1987). Altered geometry of webs in spiders with regenerated legs. *Nature*, 328(6127), 247–248. doi:10.1038/328247a0

Vollrath, F., Downes, M., & Krackow, S. (1997). Design variability in web geometry of an orb-weaving spider. *Physiology & Behavior*, 62(4), 735–743. doi:10.1016/S0031-9384(97)00186-8

Vriens, L., Bosch, H., De Knijf, S., De Saeger, S., Guelinckx, R., Oosterlynck, P., ... Paelinckx, D. (2011). *De Biologische Waarderingskaart - Biotopen en hun verspreiding in Vlaanderen en het Brussels Hoofdstedelijk Gewest [en: The Biological Valuation Map - Habitats and their distribution in Flanders and the Brussels Capital Region]*. Brussels, Belgium: Mededelingen van het Instituut voor Natuur- en Bosonderzoek.

Witt, P. N., Rawlings, J. O., & Reed, C. F. (1972). Ontogeny of web-building behavior in two orb-weaving spiders. *American Zoologist*, 12(3), 445–454. doi:10.2307/3881779

Wong, B. B. M., & Candolin, U. (2015). Behavioral responses to changing environments. *Behavioral Ecology*, 26(3), 665–673. doi:10.1093/beheco/aru183

TABLES

Table 1. Descriptive statistics for the quantitative traits

	Mean	SD
Sampling date (days since January 1 st)	255.00 (September 13)	11.07
Population density (adult female spiders per 4 ha)	16.85	3.16
Body size (cephalothorax width, cm)	0.40	0.07
Mesh width (cm)	0.31	0.08
Web surface (cm ²)	623.02	303.45
Egg load	314.77	42.71

Table 2. Sources of variation in population density and spider traits. Parameters from univariate mixed models are presented with 95% Credible Intervals in parentheses. The proportions of total and among-population variation explained by models are calculated using Nakagawa and Schielzeth (2013)'s formulas for conditional and marginal R^2 .

	Population density (per 4ha)	Spider traits			
		Body size	Mesh width	Web surface	Egg load
Fixed effects	β (95% CI)	β (95% CI)	β (95% CI)	β (95% CI)	β (95% CI)
Intercept	2.70 (2.50; 2.90)	0.00 (-0.36; 0.37)	0.24 (-0.12; 0.61)	-0.10 (-0.45; 0.23)	5.83 (5.76; 5.90)
Local urbanization, medium	0.07 (-0.14; 0.27)	0.04 (-0.29; 0.36)	-0.15 (-0.39; 0.09)	-0.25 (-0.47; -0.03)	-0.10 (-0.16; -0.05)
Local urbanization, high	0.04 (-0.17; 0.24)	-0.46 (-0.80; -0.14)	-0.42 (-0.66; -0.18)	-0.39 (-0.62; -0.17)	-0.18 (-0.24; -0.13)
Landscape urbanization, medium	0.11 (-0.11; 0.33)	0.10 (-0.35; 0.52)	-0.13 (-0.61; 0.36)	0.49 (0.05; 0.94)	0.01 (-0.06; 0.08)
Landscape urbanization, high	0.10 (-0.13; 0.32)	0.32 (-0.12; 0.73)	0.00 (-0.49; 0.49)	0.48 (0.03; 0.93)	-0.03 (-0.10; 0.05)
Sampling date	-0.01 (-0.09; 0.08)	0.14 (-0.03; 0.32)	0.26 (0.07; 0.46)	0.32 (0.14; 0.50)	0.07 (-0.01; 0.15)
Random effects	σ^2 (95% CI)	σ^2 (95% CI)	σ^2 (95% CI)	σ^2 (95% CI)	σ^2 (95% CI)
Landscape-level	0.00 (0.00; 0.02)	0.06 (0.00; 0.21)	0.16 (0.04; 0.39)	0.13 (0.04; 0.32)	0.00 (0.00; 0.01)
Population-level	--	0.23 (0.13; 0.38)	0.08 (0.02; 0.18)	0.06 (0.01; 0.14)	0.00 (0.00; 0.00)
Residual variation	0.06 (0.03; 0.09)	0.70 (0.62; 0.79)	0.74 (0.66; 0.83)	0.70 (0.62; 0.78)	0.02 (0.01; 0.02)
Proportion of variation explained :					
- of total variation, by fixed and random effects	--	0.36 (0.26, 0.49)	0.33 (0.22, 0.48)	0.36 (0.25, 0.49)	0.41 (0.23, 0.64)
- of total variation, by fixed effects only	--	0.11 (0.04, 0.19)	0.12 (0.04, 0.23)	0.19 (0.08, 0.30)	0.37 (0.19, 0.58)
- of population-level variation, by fixed and random effects	0.13 (0.03, 0.29)	0.43 (0.18, 0.69)	0.77 (0.52, 0.94)	0.84 (0.64, 0.98)	0.77 (0.59, 0.90)
- of population-level variation, by fixed effects only	0.10 (0.02, 0.22)	0.30 (0.12, 0.48)	0.37 (0.13, 0.59)	0.52 (0.25, 0.72)	0.73 (0.49, 0.87)

Body size, Mesh width, Web surface and sampling date are centered and scaled to unit standard deviation

Reference category for urbanization levels is "low urbanization"

Family is Gaussian for Body size, Mesh width, Web surface and Poisson for Population density and Egg load

$N_{landscapes} = 21$; $N_{sites} = 63$; $N_{observations} = 621$ except for Population density ($N_{sites} = N_{observations} = 62$) and Egg load ($N_{landscapes} = 9$; $N_{sites} = 27$; $N_{observations} = 193$)

Table 3. Trait variance-covariance matrices for each hierarchical level under study, based on the multivariate mixed model (Fixed effects for this model are virtually identical to those from the univariate models presented Table 2, and are presented Supplementary Table S1). In each case, among landscapes/populations/individual variances are given on diagonals, with between-trait covariances below and corresponding between-trait correlations above the diagonal. Covariances/correlations involving Egg load are on the latent (log) scale for this trait. 95% credible intervals are given in parentheses

	Body size	Mesh width	Web surface	Egg load
(a) Among landscapes				
Body size	0.07 (0.00; 0.23)	-0.05 (-0.75; 0.64)	0.49 (-0.35; 0.92)	-0.05 (-0.90; 0.83)
Mesh width	-0.00 (-0.08; 0.08)	0.17 (0.05; 0.42)	0.04 (-0.56; 0.58)	-0.02 (-0.78; 0.78)
Web surface	0.05 (-0.01; 0.16)	0.01 (-0.08; 0.11)	0.13 (0.03; 0.30)	-0.09 (-0.87; 0.81)
Egg load	-0.00 (-0.01; 0.01)	-0.00 (-0.02; 0.02)	-0.00 (-0.02; 0.01)	0.00 (0.00; 0.01)
(b) Among populations				
Body size	0.22 (0.11; 0.37)	0.36 (-0.12; 0.74)	0.18 (-0.42; 0.63)	0.13 (-0.73; 0.84)
Mesh width	0.05 (-0.01; 0.12)	0.08 (0.02; 0.16)	0.23 (-0.49; 0.72)	0.07 (-0.75; 0.81)
Web surface	0.02 (-0.03; 0.09)	0.02 (-0.02; 0.07)	0.06 (0.00; 0.14)	0.04 (-0.73; 0.76)
Egg load	0.00 (-0.01; 0.01)	0.00 (-0.00; 0.01)	0.00 (-0.00; 0.01)	0.00 (0.00; 0.00)
(c) Within populations (residual)				
Body size	0.71 (0.63; 0.79)	0.33 (0.26; 0.41)	0.33 (0.26; 0.41)	0.14 (-0.03; 0.30)
Mesh width	0.24 (0.18; 0.31)	0.75 (0.66; 0.84)	0.56 (0.50; 0.61)	0.09 (-0.08; 0.24)
Web surface	0.24 (0.18; 0.30)	0.40 (0.34; 0.47)	0.71 (0.63; 0.80)	0.11 (-0.02; 0.24)
Egg load	0.02 (-0.01; 0.05)	0.01 (-0.01; 0.04)	0.02 (-0.00; 0.04)	0.03 (0.03; 0.04)

FIGURE LEGENDS

Figure 1. Left: photograph of a female *Araneus diadematus* showing the distinctive cross pattern on the abdomen (credit: Maxime Dahirel); CT_w : cephalothorax width, used as a proxy of body size. Right: diagram of an *Araneus diadematus* web, with radii spanning from the central hub supporting a sticky silk thread spiral (bold). D_v and D_h refer to the vertical and horizontal diameters of the full web (D_{web}) and the free zone (D_{free}), used to estimate the capture area (in grey); the number of threads of the capture spiral on the horizontal and vertical axes of the capture area is used to calculate average mesh width.

Figure 2. Location of the 21 studied landscapes within northern Belgium. Colour denotes landscape-scale urbanization level. Three 200×200 m sites were sampled per landscape. Greyed areas in background correspond to artificialized areas after CORINE Land Cover (European Environmental Agency, 2016). The dotted line rectangle encompasses the nine landscapes in which information on prey availability was obtained by Dahirel et al. (2017), and in which information about spider fecundity was obtained in the present study. Inset: Location of the study area within the broader western Europe.

Figure 3. Predicted mean *Araneus diadematus* trait values $\pm 50\%$ (thick bars) and 95% (thin lines) credible intervals as a function of urbanization intensity at two spatial scales. Predictions are based on fixed effects from univariate models (Table 2), fixed effects from the multivariate model being virtually identical (Supplementary Table S1). Predictions at one spatial scale are made by averaging over the effects of sampling date and urbanization at the other spatial scale. Grey points represent observed individual site (local scale) and landscape average values.