

Health differentials between citizens and immigrants in Europe: A heterogeneous convergence

Marwân-Al-Qays Bousmah, Jean-Baptiste, Simon Combes, Mohammad Abu-Zaineh

► **To cite this version:**

Marwân-Al-Qays Bousmah, Jean-Baptiste, Simon Combes, Mohammad Abu-Zaineh. Health differentials between citizens and immigrants in Europe: A heterogeneous convergence. Health Policy, Elsevier, 2019, 123 (2), pp.235-243. 10.1016/j.healthpol.2018.12.005 . hal-01977547v2

HAL Id: hal-01977547

<https://hal-univ-rennes1.archives-ouvertes.fr/hal-01977547v2>

Submitted on 20 Mar 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Health differentials between citizens and immigrants in Europe: A heterogeneous convergence

Marwân-al-Qays Bousmah^{*, a, b, c}, Jean-Baptiste Combes^{†, d} and
Mohammad Abu-Zaineh^{‡, c}

^aAix Marseille Univ, INSERM, IRD, SESSTIM, Marseille, France

^bORS PACA, Observatoire régional de la santé Provence-Alpes-Côte d'Azur, Marseille,
France

^cAix-Marseille Univ, CNRS, EHESS, Centrale Marseille, AMSE, and IDEP, Marseille,
France

^dUniv Rennes, EHESP, CNRS, ARENES – UMR 6051, F-35000 Rennes, France

***Corresponding author.** SESSTIM, 19-21 boulevard Jean Moulin, 13305 Marseille, France.
Tel.: +33491324600. E-mail address: marwan-al-qays.bousmah@univ-amu.fr

†EHESP, 2 avenue Gaston Berger, 35000 Rennes, France. Tel.: +33299022718. E-mail address:
jean-baptiste.combes@ehesp.fr

‡AMSE, 5-9 Boulevard Bourdet, CS 50498, 13205 Marseille, France. Tel.: +33413552520. E-mail
address: mohammad.abu-zaineh@inserm.fr

Acknowledgments

This paper uses data from SHARE Waves 1, 2, 3 (SHARELIFE), 4 and 5 (DOIs: 10.6103/SHARE.w1.500, 10.6103/SHARE.w2.500, 10.6103/SHARE.w3.500, 10.6103/SHARE.w4.500, 10.6103/SHARE.w5.500), see Börsch-Supan et al. (2013) for methodological details. The SHARE data collection has been primarily funded by the European Commission through FP5 (QLK6-CT-2001-00360), FP6 (SHARE-I3: RII-CT-2006-062193, COMPARE: CIT5-CT-2005-028857, SHARELIFE: CIT4-CT-2006-028812) and FP7 (SHARE-PREP: №211909, SHARE-LEAP: №227822, SHARE M4: №261982). Additional funding from the German Ministry of Education and Research, the U.S. National Institute on Aging (U01_AG09740-13S2, P01_AG005842, P01_AG08291, P30_AG12815, R21_AG025169, Y1-AG-4553-01, IAG_BSR06-11, OGHA_04-064) and from various national funding sources is gratefully acknowledged (see www.share-project.org). This work has been carried out thanks to the support of the A*MIDEX project (no. ANR-11-IDEX-0001-02) funded by the “Investissements d’Avenir” French Government program, managed by the French National Research Agency (ANR).

Ethics approval

Our study is based on data from SHARE Waves 1, 2, 3 (SHARELIFE), 4 and 5 (DOIs: 10.6103/SHARE.w1.500, 10.6103/SHARE.w2.500, 10.6103/SHARE.w3.500, 10.6103/SHARE.w4.500, 10.6103/SHARE.w5.500). The SHARE study is subject to continuous ethics review. During Waves 1 to 4, SHARE was reviewed and approved by the Ethics Committee of the University of Mannheim. Wave 4 of SHARE and the continuation of the project were reviewed and approved by the Ethics Council of the Max Planck Society.

Highlights:

- We investigate differences in health patterns between immigrants and citizens in Europe.
- We use panel data on more than 100,000 older adults.
- Results show a reversal of health differentials between citizens and immigrants.
- There is also substantial heterogeneity across immigrant groups.
- The importance is stressed on the wealth of the country of origin.

1
2
3
4 **Health differentials between citizens and immigrants in Europe:**
5
6 **A heterogeneous convergence**
7

8
9
10
11 **Abstract**
12

13
14 The literature on immigration and health has provided mixed evidence on
15 the health differentials between immigrants and citizens, while a growing body
16 of evidence alludes to the unhealthy assimilation of immigrants. Relying on
17 five different health measures, the present paper investigates the heterogeneity
18 in health patterns between immigrants and citizens, and also between immi-
19 grants depending on their country of origin. We use panel data on more than
20 100,000 older adults living in nineteen European countries. Our panel data
21 methodology allows for unobserved heterogeneity. We document the existence
22 of a healthy immigrant effect, of an unhealthy convergence, and of a rever-
23 sal of the health differentials between citizens and immigrants over time. We
24 are able to estimate the time threshold after which immigrants' health be-
25 comes worse than that of citizens. We further document some heterogeneity
26 in the convergence of health differentials between immigrants and citizens in
27 Europe. Namely, the unhealthy convergence is more pronounced in terms of
28 chronic conditions for immigrants from low-HDI countries, and in terms of
29 self-assessed health and body-mass index for immigrants from medium- and
30 high-HDI countries.
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

50 **Keywords:** Healthy immigrant effect · Europe · Health differentials · Health
51 convergence · Immigration · Health economics.
52
53

54
55 **JEL classification:** F22 · I14 · J15.
56
57
58

1
2
3
4 **1. Introduction**
5
6

7 The literature on immigration and health has provided mixed evidence on the
8 health differentials between citizens and immigrants in developed countries. While
9 some studies report that immigrants are in better health than citizens [1, 2, 3, 4, 5] -
10 providing support for the so-called *healthy immigrant effect* - others find the opposite
11 [6, 7]. A growing body of evidence further shows that immigrants' health deteriorates
12 with the length of residence in the host country [8, 9, 10, 11]. This phenomenon,
13 sometimes called *unhealthy assimilation*, implies that the healthy immigrant effect,
14 if any, is transitory. Describing and explaining the evolution of immigrants' health is
15 of particular importance from a research and policy perspective. Once settled in the
16 host country, immigrants would tend to have different health trajectories, not only
17 relative to their native counterparts, but also among subgroups of immigrants.
18
19
20
21
22
23
24
25
26
27
28
29

30 The present paper attempts to assess whether the healthy immigrant effect, if
31 any, is followed by a convergence of immigrants' health status toward that of citizens
32 in Europe. We use panel data on more than 100,000 older adults living in nineteen
33 European countries. The panel data methodology employed allows for unobserved
34 heterogeneity. We document the existence of a healthy immigrant effect, of an un-
35 healthy convergence, and of a reversal of the health differentials between citizens and
36 immigrants over time. We are able to estimate the time threshold after which immi-
37 grant's health becomes worse than that of citizens. We further show that the effect
38 on health of the immigrants' length of residence in the host country differs depending
39 on the wealth of the country of origin and on the health measure considered. The
40 present paper fits in the literature investigating the heterogeneity in health patterns
41 across immigrant groups, with a focus on the wealth of the immigrants' country
42 of origin. Once settled in the European countries, immigrants from poor countries
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 tend to work in the so-called 3D jobs (dirty, dangerous and demeaning) [12]. In this
5 regard, the positive health selection (the healthy immigrant effect) would be higher
6 for immigrants from poorer countries because of the physically demanding nature of
7 their work [13]. This would also be a factor of faster health deterioration.
8
9

10
11
12 As we investigate the effect on health of the immigrants' length of stay in Europe,
13 we focus on older immigrants. In North-West European countries, the three decades
14 after 1945 were characterized by large-scale labour immigration. Immigration into
15 southern European destinations became significant from the 1970s [14]. We are
16 now able to investigate the evolution of immigrants' health over a long time span,
17 which explains the recent development of studies on older immigrants in Europe
18 [14, 8, 9, 7, 15].
19
20
21
22
23
24
25
26

27 The remaining of the paper is organized as follows. Section 2 reviews the liter-
28 ature on immigrants' health in developed countries. The data and the econometric
29 methodology are detailed in Sections 3 and 4, respectively. Section 5 presents the
30 results, and Section 6 discusses the implications of the findings and concludes.
31
32
33
34
35
36

37 **2. Literature review: immigrant's health in developed countries**

38
39

40 In developed countries, health differentials between natives and immigrants and
41 their evolution have been linked to the interplay of several factors. These include
42 socio-economic conditions, individual factors (immigrants' behavior, health-related
43 selection), and characteristics of both the country of origin (social norms, wealth)
44 and of destination (social norms, integration policies towards immigrants). These
45 health differentials are often mirrored by differences in access to health care.
46
47
48
49
50
51

52 McDonald and Kennedy [1] document a healthy immigrant effect and an un-
53 healthy convergence in Canada regarding chronic conditions, but not self-assessed
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 health, indicating that this phenomenon would be more pronounced in terms of
5
6 physical health than health perceptions. Also in Canada, while showing the exist-
7
8 tence of a healthy immigrant effect in the case of depression, Wu and Schimmele [2]
9
10 find that depression among immigrants increases with their length of stay.
11

12 In the United States, immigrants are less likely to have health insurance coverage
13
14 and have lower health care use and spending than their native counterparts [16, 17,
15
16 18, 19, 20]. Observed differences in the access to health insurance and health care
17
18 have been seen as one of the interplaying factors behind citizen-immigrant health
19
20 differentials. Antecol and Bedard [3] show that immigrants have a lower body-
21
22 mass index (BMI) upon arrival, but that they converge to that of the Americans
23
24 as the length of stay increases. This relationship between BMI and duration of
25
26 residence varies by gender and ethnicity [21], and by arrival cohorts, with more recent
27
28 immigrant cohorts being more likely to be obese upon arrival and to experience
29
30 a faster unhealthy convergence [22]. Other recent studies show that social norms
31
32 in both the host country and the country of origin influence immigrants' health
33
34 behaviors. Focusing on smoking behaviors, Leung [23] finds that assimilation in
35
36 the United States is positively (negatively) associated with the likelihood of being a
37
38 smoker for immigrants from low-smoking (high-smoking) countries.
39
40

41 Analyses of the relationship between immigrants' health and duration of residence
42
43 in Europe have yielded mixed and sometimes conflicting evidence. Lindert et al. [24]
44
45 report disparities across migrant groups in the prevalence of mental disorders and in
46
47 the access and utilization of mental health services. Solé-Auró et al. [7] investigate the
48
49 differences in health care utilization between older natives and immigrants. They find
50
51 that older immigrants, particularly those who arrived recently, have a higher use of
52
53 health care services (physician visits and hospital stays), due to differences in health,
54
55 health behaviors, socio-economic status or countries' health system characteristics.
56
57
58

1
2
3
4 Moullan and Jusot [25] show the existence of a North-South gradient in immigrants'
5 health status, with immigrants in Italy and Spain having a better health status than
6 in Belgium and France. Using three different health measures (self-reported health
7 status, limiting long-standing illnesses, and self-reported chronic illness), Giannoni
8 et al. [5] show that non-European immigrants tend to have better health in countries
9 with pro-immigrant integration policies.

10
11
12
13
14
15
16 Using Swedish data, Pudaric et al. [26] show that, compared with Swedish-born
17 individuals, country of birth is associated with poor health status (for immigrants
18 from Southern Europe, Eastern Europe and Finland) and impaired instrumental ac-
19 tivities of daily living (for Southern Europeans, Finns and refugees from developing
20 countries). The authors also provide a theoretical framework for the link between
21 immigrants' health deterioration and acculturation. They argue that poorly ac-
22 culturated immigrants are exposed to long-term stress reactions due to a series of
23 individual, structural and cultural factors, which may in turn harm their health.
24 Bengtsson and Scott [27] highlight a gap in sickness benefit consumption between
25 Swedish natives and immigrants, and also between immigrants depending on the
26 country of birth. The authors show that while western immigrants and natives dis-
27 play similar patterns, immigrants from labor-sending countries, whose occupations
28 are more physically demanding, have a higher average number of sick days.
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43

44 In Spain, Malmusi et al. [8] document large migration-related health inequalities,
45 with immigrants from poor areas being the worse-off group, which can be related
46 to immigrants' socio-economic deprivation compared with natives. Evidence that
47 immigrants' health deteriorates over time at a faster rate for immigrants working in
48 high-risk jobs have been found in Spain [28] and in Germany [11].
49
50
51
52
53

54 In France, Vignier et al. [29] report that individuals in threat-related exile have
55 a lower probability of accessing health care. Furthermore, it has been shown that
56
57
58

1
2
3
4 undocumented immigrants experience great difficulty in exercising their rights to
5 health care and tend to underutilize the *State Medical Assistance* [30]. The settlement
6 process of recently arrived immigrants from sub-Saharan Africa has been shown to
7 be long and precarious [31]. This long period of insecurity is mirrored by a lower
8 health care utilization of recently arrived immigrants compared to the French-born
9 population with equivalent health needs. This is reported in Berchet and Jusot
10 [32], who also indicate that the healthy immigrant effect was mainly observed for
11 immigrants who settled in France before the economic crisis of the mid-1970s. And
12 even for these first- and second-generations immigrants, studies show that they suffer
13 from premature dependency in old age [33], and, more generally, that their health
14 became worse than that of the French-born population from the mid-2000s [34, 35].
15 Such health disparities are mainly due to differences in social capital, income and
16 occupation status [36], and vary depending on the region of origin [37].
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31

32 **3. Data**

33
34
35
36 We use data from the SHARE survey [38], which primarily aims at investigat-
37 ing ageing issues in Europe. The SHARE survey has also been used to investigate
38 the relationship between immigration and health [9, 7, 15]. Our analysis is based
39 on four waves conducted in 2004-2005, 2006-2007, 2010-2012 and 2013, respectively.
40 These waves share the same methodology. The survey samples individuals over 50
41 years old in European countries. Individuals of the same household are also sur-
42 veyed, regardless of their age. The 19 countries in the sample are Austria, Belgium,
43 Czech Republic, Denmark, Estonia, France, Germany, Greece, Hungary, Ireland,
44 Italy, Luxembourg, the Netherlands, Poland, Portugal, Slovenia, Spain, Sweden,
45 and Switzerland.
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 Individuals' citizenship status (citizen or immigrant) is the main variable of in-
5
6 terest. Among the alternative definitions of "immigrant", that are based on country
7
8 of birth, nationality or length of stay in the host country [39], we have opted for the
9
10 definition based on nationality, as the focus is on individuals who are legal residents,
11
12 either permanent, temporary or refugees, and who represent a distinct category of
13
14 the population. A study including three different categories - immigrants born for-
15
16 eigners in a foreign country, naturalized citizens and native citizens - would have
17
18 been of great interest, but we cannot identify naturalized citizens in our data.
19

20
21 When individuals are surveyed for the first time, major life events are collected
22
23 retrospectively. Thus, we know the immigrants' year of arrival, allowing us to com-
24
25 pute the length of stay in the host country. As it is likely to be highly correlated
26
27 with age, we divide the length of stay by the age. This measures the immigrants'
28
29 share of lifetime in the host country. Immigrants in the sample arrived in the host
30
31 country between 1916 and 2013. Immigrants' year of arrival distribution is shown in
32
33 Figure 1.
34

35
36 *[Figure 1 about here]*
37
38

39
40 We also have information on the immigrants' country of origin, from which we
41
42 construct a categorical variable based on the wealth of the country of origin. We
43
44 use the UN's Human Development Index (HDI) as a measure of wealth. The HDI
45
46 index doesn't measure economic growth alone but also considers individuals and their
47
48 capabilities. GDP is a particular measure of wealth focusing only on this part of
49
50 wealth which is measurable by a monetary value. By using HDI we intend to move
51
52 beyond GDP and account a broader societal dimension of wealth and well-being.
53
54 Based on their HDI value in 2015, countries of origin are classified as low human
55
56 development ($HDI < 0.700$), medium ($0.700 \geq HDI < 0.836$) and high human
57
58 development ($HDI \geq 0.836$).
59

1
2
3
4 development (HDI \geq 0.836). The 0.700 cut-off is commonly used to distinguish
5
6 low- from medium-HDI countries, and the 0.836 cut-off represents the lowest HDI
7
8 value among European Union countries (Hungary), as performed by Malmusi et al.
9
10 [8]. Although immigrants in the sample arrived in the host country between 1916
11
12 and 2013, we only use the 2015 HDI of the country of origin, as more than 83%
13
14 of the immigrants left their country of origin before the oldest year for which HDI
15
16 data is available. The wealth of the country of origin should be interpreted with an
17
18 appropriate degree of caution related to the fact that, although HDI data exhibit a
19
20 strong path-dependency, some countries may be categorized in a given HDI category
21
22 which was not the same at the time of migration.
23
24

25 Detailed definitions of the variables used are given in Table 1. Five different health
26
27 measures are used as dependent variables: (1) SAH (whether the individual reports a
28
29 less than very good health), as is common in the literature (see for instance Antecol
30
31 and Bedard 3 or Malmusi et al. 8), (2) the body-mass index (whether the individual
32
33 is overweight or obese), (3) chronic conditions (the number of chronic diseases), (4)
34
35 mental health (whether the individual is depressed), and (5) physical limitations
36
37 (whether the individual has one or more limitations with activities of daily living
38
39 [ADL]). An extended discussion on the relevance of the health measures considered
40
41 is provided in Appendix A1 of the electronic supplementary material. Descriptive
42
43 statistics of the variables used in the analyses are given in Table A1 of the electronic
44
45 supplementary material. Individuals in the sample are aged 22 to 103. Citizens'
46
47 and immigrants' age distributions are relatively similar, as shown in Table A2 of
48
49 the electronic supplementary material. Table A3 of the electronic supplementary
50
51 material displays the percentage of immigrants by region of origin. The majority of
52
53 immigrants come from the European Union or other European countries (48.09% and
54
55 30.48%, respectively). Stateless persons represent 8.07% of the sample of immigrants,
56
57
58
59
60
61
62
63
64
65

1
2
3
4 and immigrants coming from Africa represent 7.74%. Immigrants from the Americas
5
6 or Asia and Oceania represent slightly less than 3%, respectively.
7
8

9 *[Table 1 about here]*
10

11 12 13 **4. Econometric methodology** 14

15
16 We use a Mundlak [40] panel probit model for all binary dependent variables
17 (Equation 1), and a Mundlak [40] panel Poisson model for the discrete positive de-
18 pendent variable, that is, the number of chronic diseases (Equation 2). The Mundlak
19 [40] specification is a fixed effect for all time-varying variables and a random effect
20 for all time-constant variables. With a usual fixed effect model we would not be able
21 to estimate a parameter for time-constant variables. With a random specification
22 we would impose a certain specification to time-varying and time-constant variables.
23 The Mundlak specification allows us to take the best of the two specification by
24 transforming a random effect statistical routine in a fixed effect model for the time-
25 varying variables if and only if the intra individuals means of these variables are
26 included [41]. For individual i ($i = 1, \dots, n$) in survey year t ($t = 1, \dots, T$):
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41

$$42 \quad P(y_{it} = 1|x_{it}) = \Phi(x_{it}\beta + c_i) \quad (1)$$

43
44 where Φ is the standard normal cumulative distribution function.
45
46
47

$$48 \quad P(Y_{it} = y_{it}|x_{it}) = F(y_{it}, x_{it}\beta + c_i) \quad (2)$$

49
50 where $F(x, z) = P(X = x)$. X is Poisson distributed with mean $\exp(z)$.
51
52

53 Equation 1 and Equation 2 are estimated using maximum likelihood. In both
54 models, x_{it} is the vector of explanatory variables (some of these variables are con-
55
56
57
58

stant over time) and c_i is the time-invariant unobservable. We include the within-individual means of the time-varying regressors, \bar{x}_i , to allow for the possibility that the unobserved time-invariant individual effect and the regressors are correlated (see Wooldridge 42 for further methodological details):

$$c_i = \bar{x}_i\theta + \nu_i \quad \text{and} \quad E(c_i|x_i) = \bar{x}_i\theta \quad (3)$$

where \bar{x}_i is the panel-level mean of x_{it} , and ν_i is a time-invariant unobservable that is uncorrelated to the regressors.

We include both age and period (survey wave) effects in the model. Cohort effects are not included due to the age-period-cohort identification problem and the impossibility to disentangle these three effects, as they are mathematically confounded [43]. Period effects are likely to capture unobserved factors common to all individuals in the sample. Country fixed effects are also included in the model to control for unobserved country-specific characteristics. Robust standard errors are computed at the household level to account for intra-household correlation. By using the Mundlak specification, the model accounts for the correlation between time-varying regressors and individual effects. We nonetheless acknowledge the potential presence of time-varying sources of endogeneity (for instance an income shock affecting both the access to health care and health outcomes).

Finally, we calculate average partial effects (APEs) to discuss the results in terms of magnitude. APEs are computed by averaging the partial effects across the population distribution of individual effects, which integrates out the unobserved heterogeneity [42].

As we focus on older immigrants, we cannot rule out the potential presence of a so-called *salmon bias* in our analysis. The literature on the salmon bias - immigrants

1
2
3
4 who suffer from a long-term illness returning to their country of origin - has recently
5
6 been reviewed in a study by Wallace and Kulu [44], reporting mixed evidence. More
7
8 importantly, when there is evidence of a salmon bias, studies show that the mortality
9
10 of those who return to their country of origin should have been unrealistically high
11
12 in order to offset the migrant mortality advantage [45, 44]. From an econometric
13
14 standpoint, the salmon bias would imply that there is measurement error in the
15
16 dependent variable due to the fact that health of some immigrants would not be
17
18 accounted for. The measurement error would thus be positively associated with the
19
20 dummy variable for being an immigrant. In a similar manner, the immigrants' length
21
22 of stay in the host country would be positively associated with the measurement error,
23
24 in the sense that the parameter on length of stay would have been larger without
25
26 the presence of a salmon bias, if any.
27
28
29

30 **5. Results**

31 32 33 **5.1 Baseline model**

34
35
36
37 Maximum likelihood estimates of the model (Equation 1 and Equation 2) and
38
39 APEs are reported in Table A4 of the electronic supplementary material. The es-
40
41 timates show that immigrants are more likely to be in better health than citizens.
42
43 This is true for all dependent variables except mental health, for which there is no
44
45 significant difference between citizens and immigrants. The APE of being an im-
46
47 migrant ranges from -0.029 (for ADL limitations) to -0.067 (for chronic conditions).
48
49 The probability of reporting poor SAH is 4.2%-points lower for immigrants than for
50
51 citizens. As reported in the introduction, such healthy immigrant effect has been
52
53 highlighted previously. Nevertheless, the results reveal that the immigrants' length
54
55 of stay - measured as the share of lifetime in the host country - is significantly and
56
57
58
59
60
61
62
63
64
65

1
2
3
4 negatively associated with their health for all five health measures. A 10% increase
5
6 in the share of lifetime in the host country increases the probability of reporting poor
7
8 SAH by 1.13%-points. Immigrants who spent most of their lives in the host country
9
10 have a 11.3% higher probability of reporting poor health than those who arrived
11
12 very recently. This is illustrated in Figure 2, which shows the estimated relation-
13
14 ship between the immigrants' length of stay and their health for the five dependent
15
16 variables.
17

18
19
20 *[Figure 2 about here]*
21
22

23 The estimated health of citizens is included for comparison. We can see that the
24
25 healthy immigrant effect holds only for immigrants who arrived recently in the host
26
27 country. For all five health measures, health worsens with the share of lifetime in
28
29 the host country. Another important result is that, when the share of lifetime in
30
31 the host country increases, immigrants' health eventually becomes poorer than that
32
33 of citizens, as illustrated in Figure 2. Immigrants' self assessed and mental health
34
35 reach the level of citizens when they have spent around 30 to 35% of their life in the
36
37 host country (corresponding to, on average, approximately 18 to 21 years spent in
38
39 the host country). For obesity and overweight, immigrants reach the level of citizens
40
41 after having spent 45% of their life in the host country (on average approximately
42
43 27 years). This figure rises to 50% for chronic conditions (on average approximately
44
45 30 years) and 80% for ADL limitations (on average approximately 52 years).
46
47

48 We now interpret the coefficients for the other variables, which are common to
49
50 all individuals in the sample. Compared to men, women are less likely to report
51
52 poor health, to be overweight or obese, or to have physical limitations, but are
53
54 more likely to have poor mental health. No significant gender difference is found
55
56 for chronic conditions. Formal education is negatively associated with poor health
57
58

1
2
3
4 status, with evidence of an education gradient. Current job situation is found to have
5
6 different effects depending on the health measure considered. Note that the variable
7
8 on current job situation has a specific category for permanently sick individuals,
9
10 because their job situation is necessarily determined by their condition. This allows
11
12 to appropriately capture the effect of being unemployed. In the case of SAH and
13
14 mental health, being retired is associated with better health compared to all other
15
16 categories (employed or self-employed, unemployed and homemaker or permanently
17
18 sick). Being employed or self employed is associated with better health compared
19
20 to all other categories in the case of BMI and chronic conditions. Regarding ADL
21
22 limitations, being unemployed, homemaker or permanently sick is associated with
23
24 poor health, while no significant association is found for the other categories. Marital
25
26 status is found to have no effect on chronic conditions and ADL limitations. The
27
28 likelihood of having poor mental health is 9%-points higher for widowed than for
29
30 individuals who are married or in a registered partnership (the reference category),
31
32 while no significant association is found for the other categories (never married and
33
34 divorced). Being divorced is positively associated with being overweight or obese.
35
36 Never married and widowed individuals are more likely to have poor SAH. Drinking
37
38 is positively associated with being overweight or obese, negatively associated with
39
40 having poor mental health, and has no significant association with SAH, chronic
41
42 conditions and ADL limitations. Physical inactivity increases the probability of
43
44 having poor health for all health measures except BMI, for which it is negatively
45
46 associated with being overweight or obese. Having children is associated with better
47
48 mental and SAH, fewer chronic conditions, a higher likelihood of being overweight or
49
50 obese and appears to be not associated with ADL limitations. Having grandchildren
51
52 is associated with a higher likelihood of being overweight or obese, more chronic
53
54 conditions, poor SAH (for three grandchildren or more) and is not associated with
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 mental health and ADL limitations. For all health measures except BMI, the number
5
6 of medical consultations is strongly associated with poor health, with a marked
7
8 gradient. In the same way, the number of nights in hospital is linearly associated
9
10 with poor health except for BMI, for which individuals who spent six or more nights
11
12 in hospital in the year preceding the survey are less likely to be overweight or obese.
13
14 Households' ability to make ends meet is associated with a higher likelihood of being
15
16 overweight or obese. For all other health measures, the ability to make ends meet
17
18 is associated with better health, with evidence of a socio-economic gradient in the
19
20 case of self-assessed and mental health. The analysis of period (survey wave) effects
21
22 indicates that the likelihood of being overweight or obese, of developing chronic
23
24 diseases and of having poor mental health increased over time (from 2004-2005 to
25
26 2013). Almost no significant period effects are found for self-assessed health and
27
28 ADL limitations. Finally, age is positively associated with poor health in the case
29
30 of SAH and ADL limitations, and negatively associated with poor health in the
31
32 case of BMI, chronic conditions and mental health. The unexpected negative effect
33
34 of age for chronic conditions and mental health might due to the fact that, for an
35
36 undocumented reason, the average number of chronic conditions and mental health
37
38 are slightly decreasing between each individual's first and second interview (which
39
40 may be distinct from the first and second wave, as some individuals entered the
41
42 SHARE survey after the first survey-wave).
43
44
45
46

47 **5.2. Wealth of the country of origin**

48
49

50
51 To gain more understanding about the effects of being immigrant on health, we
52
53 also investigate whether the effects highlighted previously change depending on the
54
55 wealth of the country of origin. To do so, the immigrant status variable of the
56
57
58
59
60
61
62
63
64
65

1
2
3
4 previous model is allowed to take on several values based on the level of human
5
6 development of the country of origin. This variable is also interacted with the im-
7
8 migrants' share of lifetime in the host country. Maximum likelihood estimates and
9
10 APEs are presented in Table A5 of the electronic supplementary material. Note
11
12 that all the other results remain qualitatively the same, in terms of sign, magnitude,
13
14 and statistical significance. Figure 3 shows the estimated relationship between the
15
16 immigrants' length of stay and their probability of having poor health depending on
17
18 the wealth of the country of origin. This is done for the five health measures.
19
20

21
22 *[Figure 3 about here]*
23
24

25 The results reveal that the effect of being an immigrant on health, and also that
26
27 of the length of stay in the host country, varies depending on the wealth of the coun-
28
29 try of origin and the health measure considered. When arriving in the host country,
30
31 immigrants from high HDI-countries are less likely to have poor SAH than citizens,
32
33 while no significant association is found for the other categories. Nevertheless, when
34
35 the share of lifetime in the host country increases, immigrants from medium- and
36
37 high-HDI countries have poorer SAH, as illustrated in Figure 3a. The length of stay
38
39 does not appear to change the SAH of stateless persons or immigrants from low-HDI
40
41 countries. In the case of BMI, we found previously that immigrants are less likely
42
43 to be overweight or obese than natives, but that this likelihood increases with the
44
45 share of lifetime in the country. This result seems to hold mainly for immigrants
46
47 from high-HDI countries. Although their likelihood of being overweight or obese is
48
49 10.3%-points lower than that of citizens, immigrants from high-HDI countries who
50
51 spent most of their lives in the host country are 21.4% more likely to be overweight
52
53 or obese than those who arrived very recently, as shown in Figure 3b. The results
54
55 regarding chronic conditions reveal a marked socio-economic gradient in the effect
56
57
58

1
2
3
4 of being an immigrant and in that of their length of stay in the host country. Im-
5
6 migrants have a lower likelihood of developing chronic diseases, as shown previously.
7
8 Compared to citizens, the likelihood of developing a chronic disease is 20.1%-points
9
10 lower for immigrants from low-HDI countries, while this figure falls to 8.4% and 3%
11
12 for immigrants from medium- and high-HDI countries, respectively. We showed ear-
13
14 lier that, among immigrant, the likelihood of developing chronic diseases increases
15
16 with the share of lifetime in the country. We now show that, although immigrants
17
18 from low-HDI countries have the lowest likelihood of having chronic conditions, they
19
20 have the highest rate of health deterioration. There is a 28.2% difference between
21
22 immigrants from low-HDI countries who arrived very recently and those who spent
23
24 most of their lives in the host country. This figure falls to 14.4% and 7.7% for immi-
25
26 grants from medium- and high-HDI countries, respectively. The poorer the country
27
28 of origin, the higher the deterioration of health with increasing share of lifetime in
29
30 the host country. This is illustrated in Figure 3c. In the case of mental health, it
31
32 seems that stateless persons have a lower likelihood of being depressed, while no
33
34 significant association is found for the other groups. The results also indicate that
35
36 immigrants from medium-HDI countries and stateless persons experience a mental
37
38 health deterioration with the length of stay in the host country. Finally, the results
39
40 show that the results found previously regarding ADL limitations hold mainly for
41
42 immigrants from medium-HDI countries. They have a lower likelihood of having one
43
44 or more ADL limitations, but this likelihood increases with the length of stay in the
45
46 host country.
47
48
49
50

51 **6. Conclusion**

52
53
54
55 The question of whether there are significant health differentials between citizens
56
57
58
59

1
2
3
4 and immigrants has extensively been addressed in the literature. Nonetheless, less is
5
6 known about the evolution of immigrants' health over time. This paper attempted
7
8 therefore to explore the dynamics of citizen-immigrant health differentials as well
9
10 as the determinants of these dynamics. It employed panel models that allow for
11
12 unobserved heterogeneity amongst more than 100,000 older persons living in nine-
13
14 teen European countries. Unlike previous studies that only focused on one or two
15
16 measures of health, the present analysis has considered five different indicators of
17
18 health status. This allowed to account for the multidimensionality of health and its
19
20 evolution over time. Assessing the impact of the length of stay on immigrant health
21
22 requires adjusting for the effect of age. The latter has been accounted for by using
23
24 an age-adjusted measure of the length of stay in the host country.
25
26

27 Our paper highlighted large health differentials (1) between citizens and immi-
28
29 grants, (2) between immigrants depending on their length of stay in the host country,
30
31 and (3) between immigrants depending on the wealth of their country of origin. These
32
33 differentials are found to vary depending on the health measure considered. Some
34
35 interesting results emerging from our analysis are worth discussing. First, our study
36
37 generally, corroborates previous evidence on the healthy immigrant effect – according
38
39 to which immigrants are initially more likely to be in better health as compared to
40
41 citizens [1, 2, 3, 4, 5]. Secondly, our study also corroborates previous findings on the
42
43 unhealthy assimilation of immigrants in developed countries [8, 9, 10, 11]. For all
44
45 health measures considered, we found that, on average, immigrants' health deterio-
46
47 rates over time. In contrast to previous studies, our results not only shed light on
48
49 the convergence of immigrants' health towards that of citizens. Interestingly, using
50
51 an age-adjusted measure of the length of stay in the host country, we were able to
52
53 show that this unhealthy convergence is such that the health of immigrants eventu-
54
55 ally becomes poorer than that of citizens. Thirdly, our results clearly revealed the
56
57
58
59
60
61
62
63
64
65

1
2
3
4 presence of substantial heterogeneity in the observed unhealthy convergence, with
5
6 respect to both the wealth of the country of origin and the health measured con-
7
8 sidered. For instance, in terms of chronic conditions, the health of immigrants from
9
10 low-HDI countries tends to deteriorate faster than that of immigrants from richer
11
12 countries. This is not surprising given that immigrants from low-HDI countries tend
13
14 to work in the so-called 3D jobs (dirty, dangerous and demeaning) once settled in
15
16 the European countries [12]. By contrast, the unhealthy convergence with respect
17
18 to SAH seems to hold mainly for immigrants from medium- and high-HDI coun-
19
20 tries. In this case, the convergence of SAH towards that of citizens may well be a
21
22 result of peer effects. Self-assessments of health have been shown to be influenced
23
24 by social comparisons within age groups, especially among older people [46]. Thus,
25
26 immigrants from medium- and high-HDI countries may tend to rate their health as
27
28 similar to that of their native peers, which are more likely to belong to the same
29
30 socio-economic group.
31
32

33 Altogether, such heterogeneity in the convergence of health differentials between
34
35 immigrants and citizens in Europe seems to indicate that the unhealthy convergence
36
37 is more pronounced in terms of chronic conditions for immigrants from low-HDI
38
39 countries, and in terms of SAH and BMI for immigrants from medium- and high-
40
41 HDI countries. Future research shall attempt to identify the underlying factors of
42
43 such heterogeneous convergence.
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

References

- [1] James Ted McDonald and Steven Kennedy. Insights into the ‘healthy immigrant effect’: health status and health service use of immigrants to Canada. *Social Science & Medicine*, 59(8):1613–1627, 2004.
- [2] Zheng Wu and Christoph M Schimmele. The healthy migrant effect on depression: variation over time? *Canadian Studies in Population*, 32(2):271–295, 2005.
- [3] Heather Antecol and Kelly Bedard. Unhealthy assimilation: why do immigrants converge to American health status levels? *Demography*, 43(2):337–360, 2006.
- [4] Cristina Hernández-Quevedo and Dolores Jiménez-Rubio. A comparison of the health status and health care utilization patterns between foreigners and the national population in Spain: new evidence from the Spanish National Health Survey. *Social Science & Medicine*, 69(3):370–378, 2009.
- [5] Margherita Giannoni, Luisa Franzini, and Giuliano Masiero. Migrant integration policies and health inequalities in Europe. *BMC Public Health*, 16(1):463, 2016.
- [6] Signe Smith Nielsen and Allan Krasnik. Poorer self-perceived health among migrants and ethnic minorities versus the majority population in Europe: a systematic review. *International Journal of Public Health*, 55(5):357–371, 2010.
- [7] Aïda Solé-Auró, Montserrat Guillén, and Eileen M Crimmins. Health care usage among immigrants and native-born elderly populations in eleven European countries: results from SHARE. *European Journal of Health Economics*, 13(6):741–754, 2012.
- [8] Davide Malmusi, Carme Borrell, and Joan Benach. Migration-related health inequalities: showing the complex interactions between gender, social class and place of origin. *Social Science & Medicine*, 71(9):1610–1619, 2010.
- [9] Donatella Lanari and Odoardo Bussini. International migration and health inequalities in later life. *Ageing & Society*, 32(6):935–962, 2012.
- [10] Santosh Jatrana, Samba Siva Rao Pasupuleti, and Ken Richardson. Nativity, duration of residence and chronic health conditions in Australia: Do trends converge towards the native-born population? *Social Science & Medicine*, 119:53–63, 2014.
- [11] Osea Giuntella and Fabrizio Mazzonna. Do immigrants improve the health of natives? *Journal of Health Economics*, 43:140–153, 2015.
- [12] Peter Stalker. *The work of strangers: a survey of international labour migration*. Geneva, Switzerland: International Labour Office, 1994.
- [13] Yao Lu. Test of the “healthy migrant hypothesis”: a longitudinal analysis of health selectivity of internal migration in Indonesia. *Social Science & Medicine*, 67(8):1331–1339, 2008.
- [14] Paul White. Migrant populations approaching old age: prospects in Europe. *Journal of Ethnic and Migration Studies*, 32(8):1283–1300, 2006.
- [15] Amelie F Constant, Teresa García-Muñoz, Shoshana Neuman, and Tzahi Neuman. A “healthy immigrant effect” or a “sick immigrant effect”? Selection and policies matter. *European Journal of Health Economics*, 19(1):103–121, 2018.
- [16] Olveen Carrasquillo, Angeles I Carrasquillo, and Steven Shea. Health insurance coverage of immigrants living in the United States: differences by citizenship status and country of origin. *American Journal of Public Health*, 90(6):917, 2000.
- [17] Leighton Ku and Sheetal Matani. Left out: immigrants’ access to health care and insurance. *Health Affairs*, 20(1):247–256, 2001.

- 1
2
3
4 [18] Leighton Ku. Health insurance coverage and medical expenditures of immigrants and native-
5 born citizens in the United States. *American Journal of Public Health*, 99(7):1322–1328, 2009.
6
7 [19] Adriana M Reyes and Melissa Hardy. Another health insurance gap: gaining and losing
8 coverage among natives and immigrants at older ages. *Social Science Research*, 43:145–156,
9 2014.
- 10 [20] Wassim Tarraf, Elham Mahmoudi, Heather E Dillaway, and Hector M González. Health
11 spending among working-age immigrants with disabilities compared to those born in the US.
12 *Disability and Health Journal*, 9(3):479–490, 2016.
13
14 [21] Annie Ro and Georgiana Bostean. Duration of US stay and body mass index among Latino
15 and Asian immigrants: a test of theoretical pathways. *Social Science & Medicine*, 144:39–47,
16 2015.
- 17 [22] Osea Giuntella and Luca Stella. The acceleration of immigrant unhealthy assimilation. *Health*
18 *Economics*, 26(4):511–518, 2017.
19
20 [23] Leigh Ann Leung. Healthy and unhealthy assimilation: country of origin and smoking behavior
21 among immigrants. *Health Economics*, 23(12):1411–1429, 2014.
22
23 [24] Jutta Lindert, Meryam Schouler-Ocak, A Heinz, and Stefan Priebe. Mental health, health
24 care utilisation of migrants in Europe. *European Psychiatry*, 23:14–20, 2008.
- 25 [25] Yasser Moullan and Florence Jusot. Why is the ‘healthy immigrant effect’ different between
26 European countries? *European Journal of Public Health*, 24(Supplement 1):80–86, 2014.
27
28 [26] Sonja Pudaric, Jan Sundquist, and Sven-Erik Johansson. Country of birth, instrumental
29 activities of daily living, self-rated health and mortality: a Swedish population-based survey
30 of people aged 55–74. *Social Science & Medicine*, 56(12):2493–2503, 2003.
- 31 [27] Tommy Bengtsson and Kirk Scott. Immigrant consumption of sickness benefits in Sweden,
32 1982–1991. *The Journal of Socio-Economics*, 35(3):440–457, 2006.
33
34 [28] Meritxell Solé, Luis Diaz-Serrano, and Marisol Rodríguez. Disparities in work, risk and health
35 between immigrants and native-born Spaniards. *Social Science & Medicine*, 76:179–187, 2013.
- 36 [29] Nicolas Vignier, Pierre Chauvin, and Rosemary Dray-Spira. Un système de protection sociale
37 universaliste, mais des barrières à l’accès aux soins encore trop nombreuses. In Annabel
38 Desgrées du Loû and France Lert, editors, *Parcours de vie et santé des Africains immigrés en*
39 *France*, pages 113–135. Paris, France: La Découverte, 2017.
40
41 [30] Jean-Marie André and Fabienne Azzedine. Access to healthcare for undocumented migrants
42 in France: a critical examination of State Medical Assistance. *Public Health Reviews*, 37(1):5,
43 2016.
- 44 [31] A Gosselin, Annabel Desgrées du Loû, E Lelièvre, F Lert, R Dray-Spira, and N Lydié. How
45 long do sub-Saharan migrants take to settle in France? *Population et Sociétés*, 533(May 2016),
46 2016.
47
48 [32] Caroline Berchet and Florence Jusot. État de santé et recours aux soins des immigrés en
49 france: une revue de la littérature. *Journal de Pédiatrie et de Puériculture*, 25(2):120–125,
50 2012.
51
52 [33] Mathilde Plard, Aurélien Martineau, and Sébastien Fleuret. Les immigrés au seuil du grand
53 âge. *Hommes & Migrations*, (1):31–37, 2015.
- 54 [34] Claudine Attias-Donfut and Philippe Tessier. Santé et vieillissement des immigrés. *Retraite*
55 *et Société*, (3):89–129, 2005.
56
57 [35] Florence Jusot, Jérôme Silva, Paul Dourgnon, and Catherine Sermet. Inégalités de santé liées
58

- 1
2
3
4 à l’immigration en France. *Revue Économique*, 60(2):385–411, 2009.
- 5 [36] Caroline Berchet and Florence Jusot. Inégalités de santé liées à l’immigration et capital social:
6 une analyse en décomposition. *Économie Publique/Public Economics*, 24-25:73–100, 2012.
- 7 [37] Nicolas Vaillant and François-Charles Wolff. Origin differences in self-reported health among
8 older migrants living in France. *Public Health*, 124(2):90–98, 2010.
- 9 [38] Axel Börsch-Supan, Martina Brandt, Christian Hunkler, Thorsten Kneip, Julie Korbmacher,
10 Frederic Malter, Barbara Schaan, Stephanie Stuck, and Sabrina Zuber. Data resource profile:
11 the Survey of Health, Ageing and Retirement in Europe (SHARE). *International Journal of*
12 *Epidemiology*, 42(4):992–1001, 2013.
- 13 [39] Bridget Anderson and Scott Blinder. Who counts as a migrant? Definitions
14 and their consequences. *Briefing, The Migration Observatory at the University*
15 *of Oxford*, [http://www.migrationobservatory.ox.ac.uk/resources/briefings/who-counts-as-a-](http://www.migrationobservatory.ox.ac.uk/resources/briefings/who-counts-as-a-migrant-definitions-and-their-consequences/)
16 [migrant-definitions-and-their-consequences/](http://www.migrationobservatory.ox.ac.uk/resources/briefings/who-counts-as-a-migrant-definitions-and-their-consequences/), 2017.
- 17 [40] Yair Mundlak. On the pooling of time series and cross section data. *Econometrica*, 46(1):
18 69–85, 1978.
- 19 [41] Jean-Baptiste Combes, Ulf-Göran Gerdtham, and Johan Jarl. Equalisation of alcohol par-
20 ticipation among socioeconomic groups over time: an analysis based on the total differential
21 approach and longitudinal data from Sweden. *International Journal for Equity in Health*, 10
22 (1):10, 2011.
- 23 [42] Jeffrey M Wooldridge. *Econometric analysis of cross section and panel data*. Cambridge, MA:
24 MIT Press, 2002.
- 25 [43] Andrew Bell and Kelvyn Jones. The impossibility of separating age, period and cohort effects.
26 *Social Science & Medicine*, 93:163–165, 2013.
- 27 [44] Matthew Wallace and Hill Kulu. Can the salmon bias effect explain the migrant mortality
28 advantage in England and Wales? *Population, Space and Place*, e2146, 2018.
- 29 [45] Hadewijch Vandenheede, Didier Willaert, Hannelore De Grande, Steven Simoens, and
30 Christophe Vanroelen. Mortality in adult immigrants in the 2000s in Belgium: a test of
31 the “healthy-migrant” and the “migration-as-rapid-health-transition” hypotheses. *Tropical*
32 *Medicine & International Health*, 20(12):1832–1845, 2015.
- 33 [46] Marja Jylhä. What is self-rated health and why does it predict mortality? Towards a unified
34 conceptual model. *Social Science & Medicine*, 69(3):307–316, 2009.
- 35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Table 1: Definitions of the dependent and independent variables

Variable	Type	Definition
Health measures (dependent variables)		
Self-assessed health	Binary	1 if the individual i reports a less than very good health in survey wave t ; 0 otherwise
BMI	Binary	1 if the individual i is overweight or obese in survey wave t ; 0 otherwise
Chronic conditions	Discrete	Individual i 's number of chronic diseases in survey wave t
Mental health	Binary	1 if the individual i is depressed in survey wave t ; 0 otherwise
ADL limitations	Binary	1 if the individual i has one or more limitations with activities of daily living in survey wave t ; 0 otherwise
Citizenship status	Binary	Citizen (=base category); Immigrant (legal residents, either permanent, temporary or refugee)
Immigrant's share of lifetime in the host country	Continuous	Immigrant i 's length of stay in the host country divided by his/her age
Wealth of the country of origin	Categorical	Citizen (=base category); Low HDI (<0.700); Medium HDI (0.700>=HDI<0.836); High HDI (>=0.836); Stateless
Gender	Binary	Male (=base category); Female
Formal education	Categorical	None or Primary (=base category); Secondary; Tertiary
Current job situation	Categorical	Retired (=base category); Employed or Self-employed; Unemployed; Homemaker or Permanently sick [in survey wave t]
Marital status	Categorical	Married or Registered partnership (=base category); Never married; Divorced; Widowed [in survey wave t]
Drinking	Binary	1 if the individual i was drinking more than 2 glasses of alcohol almost everyday in survey wave t ; 0 otherwise
Physical inactivity	Binary	1 if the individual i was physically inactive in survey wave t ; 0 otherwise
Number of children	Categorical	0 (=base category); 1; 2; 3 or more [in survey wave t]
Number of grandchildren	Categorical	0 (=base category); 1; 2; 3 or more [in survey wave t]
Doctor consultations	Categorical	Number of doctor consultations: 1st quartile (=base category); 2nd; 3rd; 4th [in survey wave t]
Nights in hospital	Categorical	0 (=base category); Between 1 and 5; 6 or more [in survey wave t]
Household able to make ends meet	Categorical	With great difficulty (=base category); With some difficulty; Fairly easily; Easily [in survey wave t]
Country	Categorical	Austria (=base category); Belgium; Czech Republic; Denmark; Estonia; France; Germany; Greece; Hungary; Ireland; Italy; Luxembourg; the Netherlands; Poland; Portugal; Slovenia; Spain; Sweden; Switzerland
Survey wave	Categorical	2004-2005 (=base category); 2006-2007; 2010-2012; 2013
Age	Continuous	In years

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure 1: Immigrants' year of arrival distribution






Figure 2: Effect of the immigrants' share of lifetime in the host country on health


(a) Self-assessed health




(d) Mental health



(b) BMI



(e) ADL limitations



(c) Chronic conditions






Figure 3: Effect of the immigrants' share of lifetime in the host country on health given the wealth of the country of origin


(a) Self-assessed health




(d) Mental health



(b) BMI



(e) ADL limitations



(c) Chronic conditions

