

Cytosine hydroxymethylation by TET enzymes: From the control of gene expression to the regulation of DNA repair mechanisms, and back

Audrey Lejart, Gilles Salbert, Sébastien Huet

► **To cite this version:**

Audrey Lejart, Gilles Salbert, Sébastien Huet. Cytosine hydroxymethylation by TET enzymes: From the control of gene expression to the regulation of DNA repair mechanisms, and back. *AIMS Biophysics*, AIMS Press, 2018, 5 (3), pp.182-193. 10.3934/biophy.2018.3.182 . hal-01993674

HAL Id: hal-01993674

<https://hal-univ-rennes1.archives-ouvertes.fr/hal-01993674>

Submitted on 18 Jul 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.





Review

Cytosine hydroxymethylation by TET enzymes: From the control of gene expression to the regulation of DNA repair mechanisms, and back

Audrey Lejart, Gilles Salbert and Sébastien Huet*

Univ Rennes, CNRS, IGDR-UMR 6290, F35000 Rennes, France

* **Correspondence:** Email: sebastien.huet@univ-rennes1.fr; Tel: +330223234557.

Abstract: Chromatin is a complex multi-scale structure composed of DNA wrapped around nucleosomes. The compaction state is finely regulated mainly by epigenetic marks present not only on nucleosomes but also on the DNA itself. The most studied DNA post-transcriptional modification is 5-methylcytosine (5-mC). Methylation of the cytosine at CpG islands localized at the promoter is associated with repression of transcription. On the contrary, enrichment of 5-hydroxymethylcytosine (5-hmC), one of the oxidation products of 5-mC by TET (ten-eleven translocation) enzymes, on promoters and enhancers promotes transcription activation. Recently, a new role of 5-hmC has been proposed in the context of DNA repair. 5-hmC was found to be enriched at DNA lesions and knockdown of TET led to impaired repair efficiency. Here, we review our current knowledge regarding the role of the regulation of the 5-mC/5-hmC balance by TET enzymes in the context of transcription modulation as well as DNA repair processes. In a final section, we speculate on the potential involvement of TET proteins in DNA repair mechanisms associated with transcription activation.

Keywords: chromatin; DNA repair; transcription; ten eleven translocation; 5-hydroxymethylcytosine

1. Introduction

Research work performed over the last fifty years have shown that our genetic material is not simply a linear sequence of 3 billion base pairs coding for all the proteins required by the cell. An additional layer of complexity, which allows to reach different outcomes while starting from the same DNA sequence, is provided by epigenetic regulatory mechanisms. These processes rely on labile chemical tag, such as phosphorylation or methylation, tagging the DNA double-helix or the

nucleosomes. Epigenetic marks serve two major functions: (i) they participate to signaling pathways; and (ii) they regulate chromatin architecture [1]. The principle of epigenetic signaling follows a quite simple rule: A given mark will attract or repel specific effector proteins participating in physiological mechanisms such as DNA transcription, replication or repair. The structural role of the epigenetic code remains more fuzzy but might rely on a modification of the physico-chemical properties of the chromatin fiber, thus impacting its folding state inside the nucleus. Whether epigenetic marks are able to regulate all scales of multi-step 3D chromatin organization or only specific ones remains unclear [2].

The most studied epigenetic marks are those found along the N-terminal tails of the core histones. These tails, which are localized at the surface of the compact nucleosome particle, are readily accessible to nuclear proteins among which specific enzymes responsible for writing or erasing epigenetic marks, as well as effector proteins involved in cellular processes using DNA as a template [3]. Methylation or acetylation of the histone tails are also known to regulate the interaction between the histones and the DNA, thus affecting the stability of the nucleosome. At higher folding scales, these epigenetic marks probably also regulate nucleosome/nucleosome interactions, which in turn could impact higher folding-levels of the chromatin as well as its compaction state [4].

Besides histones, the DNA molecule itself is also subject to epigenetic modifications. In contrast to histone marks which are highly diverse, only one major epigenetic modification is found on DNA and corresponds to cytosine methylation on carbon 5, mainly in a CpG dinucleotide context. With the exception of CpG islands, CpGs are highly methylated throughout the genome of somatic cells and changes in their level of methylation have been associated to differentiation and tumorigenesis [5]. While the direct impact of methylated cytosines (5-mC) on chromatin structure remains unclear, this mark shows key signaling functions in relation to transcription regulation. Indeed, depending on their CpG density, promoters enriched in 5-mC at CpG sites tend to have a lower transcription rate, a fact that has been attributed to a lack of transcription factor binding [6]. This mark is also involved in X-chromosome inactivation [7] as well as genome imprinting [8].

Cytosine methylation, in line with all other epigenetic modifications, is a dynamic mark. DNMT (DNA methyl transferase) enzymes are in charge of adding this mark along the DNA helix [9]. 5-mC removal has been proposed to occur by passive dilution [10] but also involve two different active multi-step processes (Figure 1). First, activation induced deaminases (AID), which deaminate cytosine into uracil, are also able to convert 5mC into thymidine, yet with a lower efficiency [11]. The relevance of this 5-mC clearance pathway, which later involves the Base Excision Repair (BER) machinery to replace the thymidine by a cytosine, remains debated [12]. Alternatively, 5-mC undergoes successive oxydation steps that lead to first 5-hmC (5-hydroxymethylcytosine), then 5-fC (5-formylcytosine) and finally 5-caC (5-carboxylcytosine). These oxydation steps are carried out by the TET (Ten-eleven translocation) enzymes which are 2-oxoglutarate and Fe(II)-dependent dioxygenases [13,14]. 5-fC and 5-caC are then excised and repaired by the BER machinery to ultimately return to the unmethylated cytosine state [15,16]. Interestingly, mutual regulations have been reported between enzymes controlling DNA methylation and those in charge of adding and erasing methylation at histone tails, suggesting dynamic crosstalks between DNA and histone epigenetic marks [17]. The TET family is composed of three members: TET1, TET2 and TET3. Despite sharing the same catalytic activity, these proteins are expressed differentially during development and are also present in different cell types, suggesting that they may fulfill different functions [18].




Figure 1. Cytosine methylation/demethylation cycle by DNMTs and TET enzymes. Cytosine can be methylated by DNA methyl transferase 1 (DNMT1) for methylation maintenance or by DNMT3a and DNMT3b for de novo methylation. TET (Ten-eleven translocation) enzymes oxidize 5-mC into 5-hmC (5-hydroxymethylcytosine), then 5-fC (5-formylcytosine) and finally 5-caC (5-carboxylcytosine). Both 5-fC and 5-caC are recognized and repaired by the base excision repair (BER) system, leading the restoration of an unmodified cytosine.

In this dynamic regulation of the cytosine methylation illustrated on Figure 1, 5-hmC can be seen as a transient demethylation intermediate similar to 5-fC and 5-caC. However, several observations suggest that 5-hmC might be more than such transient intermediate. First, 5-hmC is much more stable than the two other oxidized forms of 5-mC. Second, specific readers of 5-hmC have been identified such as the DNMT1-interacting protein UHRF2 or the homeobox protein Zfx1 [19]. Conversely, 5-hmC may also repel proteins that bind to 5-mC [19]. Altogether, these findings led to the idea that 5hmC could be a fully-fledged epigenetic mark and serve signaling functions by attracting or repelling specific factors [20]. In fact, data accumulate in favor of specific functional roles for 5-hmC and for the regulation of the 5-mC/5-hmC balance via TET enzymes. High levels of 5-hmC, associated with strong expression of TET enzymes, correlate with cell pluripotency [21] and are important for embryonic stem cell maintenance since impairing 5-hmC by knocking-out TET1 induces abnormal ESC differentiation [21,22]. More recently, it was also shown that hydroxymethylation of specific loci along the genome is important at late differentiation stages such as the transition from B cells to plasma cells [23] and during cell reprogramming towards pluripotency [24]. In adult organisms, 5-hmC shows tissue specificity with high levels found in the brain in contrast to other organs [25–27]. While the exact function of 5-hmC in the brain remains unclear [28], decreased level of this epigenetic mark has been observed in several neurodegenerative diseases [29]. Finally, reduced 5-hmC levels is a feature shared by multiple cancers [30]. Importantly, decreased 5-hmC levels do not depend on tumor stage, indicating that loss of 5-hmC is an early event in cancer development [31]. These various results suggest that, while 5-hmC is definitely a central player in transcription regulation, it may also participate in other cellular

functions involving DNA transactions. In this review, we present the findings that recently improved our understanding of the exact role of 5-hmC and TET enzymes in two different processes: gene transcription and DNA repair.

2. Cytosine hydroxy-methylation by TET enzymes: A central regulator of gene transcription

Before describing the current knowledge regarding the function of cytosine hydroxymethylation, it is useful to first briefly recall the function of the best known modified state of the cytosine: The 5-methylation. This epigenetic mark is a well-known negative regulator of transcription. It can regulate the binding of transcription factors at CpC islands located in regulatory regions [6,32], but it also recruits methyl-CpG binding proteins including MeCP2, MBP2 which can in turn attract co-repressor complexes displaying histone deacetylase activities such as SIN3A [33]. The cooperative action of these different players is thought to create a closed chromatin state repressive to transcription [34]. Outside the areas enriched in CpGs, the exact function of 5-mC is less clear [35]. Mirroring the inhibitory role of cytosine methylation, 5-hmC is found at genomic areas associated with active transcription. High levels of 5-hmC are found at active promoters, enhancers, transcription start sites (TSS) and gene bodies [36,37]. Since 5-hmC is a demethylation intermediate of 5-mC, these increased levels of hydroxymethylation at active transcription sites may just reflect an increased demethylation activity in these areas to prevent silencing [38]. However, several lines of evidence suggest a specific role for hydroxymethylation by TET enzymes in both the recruitment of early components of the transcription machinery and in the establishment of a chromatin landscape favorable for transcription.

First, 5-hmC is important at initial stages of transcription activation during the recruitment of pioneering factors responsible for chromatin opening at enhancer elements to facilitate the binding of subsequent transcription factors [32]. In the context of P19 cell differentiation induced by retinoic acid, two pioneering factors, MEIS1 and PBX1 are more retained to binding sites enriched in 5-hmC [32]. At enhancers bound by these factors, 5-hmC is also required for chromatin opening and for the first step of enhancer activation named enhancer priming. The mechanism underlying this positive impact of 5-hmC on the recruitment of pioneering factors remains nevertheless unclear. One hypothesis is that 5-hmC may recruit remodeling factors leading to nucleosome eviction or destabilization, thus favoring the binding of pioneering factors [32]. In line with this model, TET enzymes were shown to associate with several chromatin remodeling complexes. During osteogenesis, the activity of TET1/2 at the promoter of the bone master transcription factor Sp7 promotes the recruitment of the catalytic subunits BRG1 and BRM of the SWI/SNF chromatin remodeler, which leads to the eviction of histone H3 [39]. Surprisingly, TET1 was shown to associate not only with remodelers promoting transcription such as the SWI/SNF, but also with repressive remodeling factors such as the NURD (Nucleosome Remodeling Deacetylase) or the Polycomb complexes [40,41]. These interactions with remodelers displaying opposite functions led to propose that TET1 might balance the activity of both remodelers to allow the fine-tuning of the gene expression levels [40].

A second important function of cytosine hydroxymethylation via TET enzymes is the establishment of a chromatin landscape favorable for transcription, both in terms of epigenetic marks and 3-dimensional structure. At promoters, TET2 enzymes participate in the enrichment in the active H3K4me3 mark (tri-methylation of lysine 4 of histone H3) [42] and might also help to erase the repressive marks H3K9me3 and H3K27me3 [39]. TET enzymes also promote the mono-methylation

of lysine 4 on histone H3 (H3K4me1) at enhancers, which is important for the activation of these regulatory elements [32]. However, cytosine hydroxymethylation is not only associated with active epigenetic histone marks. Similar to the association of TET proteins with antagonist remodeling complexes discussed in the previous paragraph, these enzymes also contribute to the formation of dual epigenetic patterns. TET2 activity appears crucial for the establishment of bivalent chromatin domains enriched in both the active H3K4me3 and the repressive H3K27me3 marks at CpG islands [43]. All these different impacts of the 5-mC/5-hmC balance on the epigenetic histone code are mediated by complex cross-regulations between TET enzymes and histone methyltransferases and demethylases [39,43]. For example, the activity of TET1/2 was shown to be essential for the recruitment of Wdr5 and Set1b, two subunits of the histone methyltransferase complex COMPASS, and of the histone demethylases Jmjd2a and Jmjd3 at the Sp7 promoter upon activation of this gene [39]. This composite epigenetic network does not only signal genetic area for the transcription machinery, it also modulates the chromatin compaction state, a process which could regulate access to the target DNA sequence. Evidences for a specific effect of 5-hmC on the chromatin structure remains nevertheless relatively sparse. Results obtained on in-vitro reconstituted nucleosomes show that 5-hmC is able to modulate the interactions between the nucleosome core particle and the DNA thus affecting both nucleosome stability and compactness [44]. In cells, high 5-hmC levels are associated with loose chromatin packing [32] but it is difficult to establish a direct causal link between these two features.

The data presented above clearly show that the 5-mC to 5-hmC oxidation dynamics controlled by TET enzymes is involved at multiple stages in transcription regulation, allowing to fine-tune the expression of our genome. Nevertheless, for many of the results mentioned in this section, it is important to point out that it is difficult to discriminate a specific effect of the 5-hmC mark from the demethylation activity of the TET enzymes. Moreover, it has also been proposed that TET enzymes may act as scaffolding proteins for the recruitment of members of the transcription machinery independently of their enzymatic activity. Deplus et al. showed that, while TET2/3 participate in the recruitment of the SET1/COMPASS methyltransferase complex as well as the O-GlcNAc transferase at TSS and CpG-rich sites, no enrichment of 5-hmC is observed in these genomic regions [45]. Furthermore, TET3 isoform seems to participate in the stabilization of several nuclear receptors onto their binding sites on the chromatin independently of its dioxygenase activity [46]. Future work should help to better understand these intricate functions of the TET enzymes.

3. An emerging role for TET enzymes in DNA repair mechanisms

Recent results suggest that, besides its role in the regulation of transcription, the 5mC to 5hmC transition mediated by TET enzymes may also be involved in the DNA damage response (DDR). First, TET2 was recently shown to be crucial for the clearance of aberrant DNA methylation associated with oxidative stress [47]. Furthermore, in mouse cells of the haematopoietic lineage, simple TET1 knockout or double knockout of TET2 and TET3 led to DNA repair defects as shown by increased phosphorylation of histone variant H2AX (γ H2AX) both in the absence of exogenous DNA damage and after X-ray irradiation [48,49]. In line with this impaired DDR, removal of TET enzymes induces chromosome segregation abnormalities during mitosis [50] and unbalanced chromosome translocations [49], two mechanisms leading to genomic instability [51] and, ultimately, to tumorigenesis [49]. This involvement of TET enzymes during the DDR might be simply explained

by the impairment of the expression of many key repair proteins in TET knockout cells [48,49]. However, several lines of evidences also suggest a more direct involvement of the TET enzymes during DDR.

An important finding suggesting a direct function of TET enzymes during DNA repair is the recruitment of these enzymes at DNA lesions induced by laser irradiation, leading to a local increase of the 5-hmC mark [50]. Depending on the DNA damaging conditions, such gain in 5-hmC levels seems to require the activation of the DNA-damage-related PI3K kinases ATM, which is involved mainly into double-strand break repair mechanisms, or ATR, which was found activated for a large spectrum of DNA insults [52,53]. Both kinases participate in DNA damage signaling, for example via H2AX phosphorylation, and are also targeting multiple repair proteins to modulate their activity during the DDR [54]. More specifically, TET1 is a substrate of ATM and TET3 interacts with ATR and is phosphorylated by this enzyme [52,53]. In addition to this regulation via the ATM/ATR kinases, a complex interplay has also been reported between TET enzymes and the poly-ADP-ribose polymerase 1 (PARP1). PARP1-dependent signaling plays multiple roles during the DDR from the recruitment of early repair factors to chromatin remodeling in the vicinity of the DNA lesions [55,56]. In vitro, PARP1 and TET1 are able to modulate each other's activities [57]. Moreover, PARP1 activity regulates the transcription of the TET1 gene [58]. Altogether, the recruitment of TET enzymes at DNA lesions and their relationships with key components of the DNA repair machinery point towards a specific function of 5-hmC during DDR. Nevertheless, more work is needed to delineate the exact function of this mark at DNA breaks: Could it be the equivalent, along the DNA double-helix, to ATM-dependent H2AX phosphorylation, which signals the presence of DNA breaks, or could it promote, together with PARP1 the formation of a loose chromatin architecture facilitating access for the DNA repair machinery?

4. Coupling DNA repair mechanisms and transcription modulation: A new job for TET enzymes?

DNA transcription and DNA repair are not independent mechanisms in the cell nucleus but instead display complex inter-dependencies. Multiple factors involved in transcription inhibition, including subunits of the polycomb and NURD complexes [59] or the heterochromatin protein 1 [60], are recruited at DNA breaks, where they add repressive epigenetic marks such as trimethylation of the lysine 27 of histone 3 [61]. Thus, it seems important to shut-down transcription in the vicinity of DNA damage sites to avoid interference between the transcription machinery and the recruitment of DNA repair proteins [62]. However, the reverse does not seem to hold true and several recent experimental evidences suggest that activating the transcription of certain genes actually requires DNA breaks induction and repair [63].

In 2006, Ju et al. were the first to report that efficient estrogen-dependent transcription of the TFF1 gene requires the occurrence of double-strand breaks (DSBs) at the promoter via a topoisomerase II β activity [64]. Intriguingly, it was later reported that inducing DSBs at promoters of neuronal-activity regulated genes by chemical agents or the CRISPR-Cas9 endonuclease system was sufficient to activate the transcription of these genes [65]. Other types of DNA lesions were also observed in relation to transcription activation such as base oxydation [66] or DNA nicks [67] and these lesions were not only found at gene promoters but also within gene bodies [68] and at enhancers [67]. Importantly, transcriptional activation did not only required the induction of DNA

lesions but also their resolution via multiple repair factors including PARP1 and the DDR-dependent PIKK kinase DNA-PK [64].

The fact that many of the transcription-related DNA lesions are induced by topoisomerases and that DNA break induction seems tightly linked to transcription elongation both suggest that the formation of these breaks is necessary for the release of the topological constraints associated with RNA polymerase II progression [69]. Besides this topological effect, the presence of DNA damage at transcriptionally-active genes might also be related to the formation of a non-canonical DNA structure: the R-loop [70]. R-loops are three-stranded structures composed of an hybrid duplex associating nascent RNA with the template DNA strand [71], and the single complementary DNA strand. R-loops, which are favored by CG enrichment, might help to stabilize the transcription bubble [70] but their resolution is associated with the formation of DNA lesions that might result in genomic instability if incorrectly processed [72].

Since TET enzymes have been involved in both early stages of transcription activation and repair mechanisms, it is tempting to speculate that they may participate in this complex interplay between the repair and transcription machineries observed upon activation of the transcription process. In favor of this hypothesis, the TFF1 promoter, at which transcription-related DNA lesions were initially observed [64], is also known to undergo cycles of methylation/demethylation [73]. Furthermore, it was also recently reported that the pioneering factor FOXA1, which is found preferentially at hydroxymethylated enhancers [74], is able to nucleate numerous repair factors [75]. It remains nevertheless unclear whether TET enzymes directly participate in these different processes.

Affecting the 5mC/5hmC balance via TET activity in the context of the transcription-related DNA lesions may fulfill several functions. Because transcription requires nucleosome eviction, epigenetic histone modifications classically used by the cell as signaling marks are transiently absent from the transcribed area. Cytosine hydroxymethylation via TET activity at nucleosome-depleted areas such as R-loops [76] or topologically constrained areas, may help to signal DNA lesions for the repair machinery or orient the repair process towards the most appropriate pathway to avoid deleterious effects [72]. Independently of their activity, TET enzymes may also serve as recruitment platform for repair factors similar to what was reported for the pioneering factor FOXA1 [75]. Repair factors found at active transcription sites not only allow DNA break resolution, they also seem to contribute to the establishment of a transcription-competent chromatin structure. For example, upon activation of the TFF1 gene, topoisomerase II β and PARP1 were shown to promote exchange of histone linker H1, involved in transcription repression, for the high mobility group B proteins [64]. This chromatin remodeling activity may be concerted with the one reported for TET enzymes via cross-regulations such as the one reported between PARP1 and TET1 [57]. These different roles of the TET enzymes are nevertheless currently mostly speculative and more work is needed to delineate more precisely the exact implication of the TET enzymes during transcription-induced DNA repair mechanisms.

In this short review, we have highlighted how recent findings extended the spectrum of action of the TET enzymes from the modulation of transcription to a contribution to DNA repair. The involvement of TET enzymes in both mechanisms might reflect the fact that repair and transcription are not independent from each other and appear highly entangled.

Acknowledgments

Audrey Lejart was supported by a PhD fellowship from the Région Bretagne and the Ligue Nationale contre le Cancer.

Conflict of interest

All authors declare no conflicts of interest in this paper.

References

1. Geiman TM, Robertson KD (2002) Chromatin remodeling, histone modifications, and DNA methylation—how does it all fit together? *J Cell Biochem* 87: 117–125.
2. Bonev B, Cavalli G (2016) Organization and function of the 3D genome. *Nat Rev Genet* 17: 661–678.
3. Bannister AJ, Kouzarides T (2011) Regulation of chromatin by histone modifications. *Cell Res* 21: 381–395.
4. Tessarz P, Kouzarides T (2014) Histone core modifications regulating nucleosome structure and dynamics. *Nat Rev Mol Cell Bio* 15: 703–708.
5. Koch A, Joosten SC, Feng Z, et al. (2018) Analysis of DNA methylation in cancer: Location revisited. *Nat Rev Clin Oncol* 15: 459–466.
6. Watt F, Molloy PL (1988) Cytosine methylation prevents binding to DNA of a HeLa cell transcription factor required for optimal expression of the adenovirus major late promoter. *Gene Dev* 2: 1136–1143.
7. Csankovszki G, Nagy A, Jaenisch R (2001) Synergism of Xist RNA, DNA methylation, and histone hypoacetylation in maintaining X chromosome inactivation. *J Cell Biol* 153: 773–784.
8. Li E, Beard C, Jaenisch R (1993) Role for DNA methylation in genomic imprinting. *Nature* 366: 362–365.
9. Okano M, Bell DW, Haber DA, et al. (1999) DNA methyltransferases Dnmt3a and Dnmt3b are essential for de novo methylation and mammalian development. *Cell* 99: 247–257.
10. Hashimoto H, Liu Y, Upadhyay AK, et al. (2012) Recognition and potential mechanisms for replication and erasure of cytosine hydroxymethylation. *Nucleic Acids Res* 40: 4841–4849.
11. Nabel CS, Jia H, Ye Y, et al. (2012) AID/APOBEC deaminases disfavor modified cytosines implicated in DNA demethylation. *Nat Chem Biol* 8: 751–758.
12. Bochtler M, Kolano A, Xu GL (2017) DNA demethylation pathways: Additional players and regulators. *Bioessays* 39: 1–13.
13. Tahiliani M, Koh KP, Shen Y, et al. (2009) Conversion of 5-methylcytosine to 5-hydroxymethylcytosine in mammalian DNA by MLL partner TET1. *Science* 324: 930–935.
14. Tamanaha E, Guan S, Marks K, et al. (2016) Distributive processing by the iron(II)/ α -ketoglutarate-dependent catalytic domains of the TET enzymes is consistent with epigenetic roles for oxidized 5-methylcytosine bases. *J Am Chem Soc* 138: 9345–9348.
15. Müller U, Bauer C, Siegl M, et al. (2014) TET-mediated oxidation of methylcytosine causes TDG or NEIL glycosylase dependent gene reactivation. *Nucleic Acids Res* 42: 8592–8604.

16. Weber AR, Krawczyk C, Robertson AB, et al. (2016) Biochemical reconstitution of TET1-TDG-BER-dependent active DNA demethylation reveals a highly coordinated mechanism. *Nat Commun* 7: 10806.
17. Wang J, Hevi S, Kurash JK, et al. (2009) The lysine demethylase LSD1 (KDM1) is required for maintenance of global DNA methylation. *Nat Genet* 41: 125–129.
18. Szwagierczak A, Bultmann S, Schmidt CS, et al. (2010) Sensitive enzymatic quantification of 5-hydroxymethylcytosine in genomic DNA. *Nucleic Acids Res* 38: e181.
19. Spruijt CG, Gnerlich F, Smits AH, et al. (2013) Dynamic readers for 5-(hydroxy)methylcytosine and its oxidized derivatives. *Cell* 152: 1146–1159.
20. Matarese F, Carrillode SPE, Stunnenberg HG (2011) 5-Hydroxymethylcytosine: A new kid on the epigenetic block? *Mol Syst Biol* 7: 562.
21. Koh KP, Yabuuchi A, Rao S, et al. (2011) Tet1 and Tet2 regulate 5-hydroxymethylcytosine production and cell lineage specification in mouse embryonic stem cells. *Cell Stem Cell* 8: 200–213.
22. Ito S, D'Alessio AC, Taranova OV, et al. (2010) Role of Tet proteins in 5mC to 5hmC conversion, ES-cell self-renewal and inner cell mass specification. *Nature* 466: 1129–1133.
23. Caron G, Hussein M, Kulis M, et al. (2015) Cell-cycle-dependent reconfiguration of the DNA methylome during terminal differentiation of human B cells into plasma cells. *Cell Rep* 13: 1059–1071.
24. Costa Y, Ding J, Theunissen TW, et al. (2013) NANOG-dependent function of TET1 and TET2 in establishment of pluripotency. *Nature* 495: 370–374.
25. Kriaucionis S, Heintz N (2009) The nuclear DNA base 5-hydroxymethylcytosine is present in Purkinje neurons and the brain. *Science* 324: 929–930.
26. Jin SG, Wu X, Li AX, et al. (2011) Genomic mapping of 5-hydroxymethylcytosine in the human brain. *Nucleic Acids Res* 39: 5015–5024.
27. Nestor CE, Ottaviano R, Reddington J, et al. (2012) Tissue type is a major modifier of the 5-hydroxymethylcytosine content of human genes. *Genome Res* 22: 467–477.
28. Szulwach KE, Li X, Li Y, et al. (2011) 5-hmC-mediated epigenetic dynamics during postnatal neurodevelopment and aging. *Nat Neurosci* 14: 1607–1616.
29. Sherwani SI, Khan HA (2015) Role of 5-hydroxymethylcytosine in neurodegeneration. *Gene* 570: 17–24.
30. Jeschke J, Collignon E, Fuks F (2016) Portraits of TET-mediated DNA hydroxymethylation in cancer. *Curr Opin Genet Dev* 36: 16–26.
31. Haffner MC, Chaux A, Meeker AK, et al. (2011) Global 5-hydroxymethylcytosine content is significantly reduced in tissue stem/progenitor cell compartments and in human cancers. *Oncotarget* 2: 627–637.
32. Mahé EA, Madigou T, S'érandour AA, et al. (2017) Cytosine modifications modulate the chromatin architecture of transcriptional enhancers. *Genome Res* 27: 947–958.
33. Nan X, Ng HH, Johnson CA, et al. (1998) Transcriptional repression by the methyl-CpG-binding protein MeCP2 involves a histone deacetylase complex. *Nature* 393: 386–389.
34. Ng HH, Bird A (1999) DNA methylation and chromatin modification. *Curr Opin Genet Dev* 9: 158–163.
35. Schübeler D (2015) Function and information content of DNA methylation. *Nature* 517: 321–326.

36. Williams K, Christensen J, Pedersen MT, et al. (2011) TET1 and hydroxymethylcytosine in transcription and DNA methylation fidelity. *Nature* 473: 343–348.
37. S érandour AA, Avner S, Oger F, et al. (2012) Dynamic hydroxymethylation of deoxyribonucleic acid marks differentiation-associated enhancers. *Nucleic Acids Res* 40: 8255–8265.
38. Song CX, He C (2013) Potential functional roles of DNA demethylation intermediates. *Trends Biochem Sci* 38: 480–484.
39. Sepulveda H, Villagra A, Montecino M (2017) Tet-mediated DNA demethylation is required for SWI/SNF-dependent chromatin remodeling and histone-modifying activities that trigger expression of the Sp7 osteoblast master gene during mesenchymal lineage commitment. *Mol Cell Biol* 37.
40. Yildirim O, Li R, Hung JH, et al. (2011) Mbd3/NURD complex regulates expression of 5-hydroxymethylcytosine marked genes in embryonic stem cells. *Cell* 147: 1498–1510.
41. Neri F, Incarnato D, Krepelova A, et al. (2013) Genome-wide analysis identifies a functional association of Tet1 and Polycomb repressive complex 2 in mouse embryonic stem cells. *Genome Biol* 14: R91.
42. Deplus R, Delatte B, Schwinn MK, et al. (2013) TET2 and TET3 regulate GlcNAcylation and H3K4 methylation through OGT and SET1/COMPASS. *EMBO J* 32: 645–655.
43. Kong L, Tan L, Lv R, et al. (2016) A primary role of TET proteins in establishment and maintenance of De Novo bivalency at CpG islands. *Nucleic Acids Res* 44: 8682–8692.
44. Mendonca A, Chang EH, Liu W, et al. (2014) Hydroxymethylation of DNA influences nucleosomal conformation and stability in vitro. *BBA-Gene Regul Mech* 1839: 1323–1329.
45. Deplus R, Delatte B, Schwinn MK, et al. (2013) TET2 and TET3 regulate GlcNAcylation and H3K4 methylation through OGT and SET1/COMPASS. *EMBO J* 32: 645–655.
46. Guan W, Guyot R, Samarut J, et al. (2017) Methylcytosine dioxygenase TET3 interacts with thyroid hormone nuclear receptors and stabilizes their association to chromatin. *P Natl Acad Sci USA* 114: 8229–8234.
47. Zhang YW, Wang Z, Xie W, et al. (2017) Acetylation enhances TET2 function in protecting against abnormal DNA methylation during oxidative stress. *Mol Cell* 65: 323–335.
48. Cimmino L, Dawlaty MM, Ndiaye-Lobry D, et al. (2015) TET1 is a tumor suppressor of hematopoietic malignancy. *Nat Immunol* 16: 653–662.
49. An J, González-Avalos E, Chawla A, et al. (2015) Acute loss of TET function results in aggressive myeloid cancer in mice. *Nat Commun* 6: 10071.
50. Kafer GR, Li X, Horii T, et al. (2016) 5-Hydroxymethylcytosine marks sites of DNA damage and promotes genome stability. *Cell Rep* 14: 1283–1292.
51. Mahfoudhi E, Talhaoui I, Cabagnols X, et al. (2016) TET2-mediated 5-hydroxymethylcytosine induces genetic instability and mutagenesis. *DNA Rep* 43: 78–88.
52. Jiang D, Zhang Y, Hart RP, et al. (2015) Alteration in 5-hydroxymethylcytosine-mediated epigenetic regulation leads to Purkinje cell vulnerability in ATM deficiency. *Brain* 138: 3520–3536.
53. Jiang D, Wei S, Chen F, et al. (2017) TET3-mediated DNA oxidation promotes ATR-dependent DNA damage response. *EMBO Rep* 18: 781–796.
54. Blackford AN, Jackson SP (2017) ATM, ATR, and DNA-PK: the trinity at the heart of the DNA damage response. *Mol Cell* 66: 801–817.

55. Sellou H, Lebeaupein T, Chapuis C, et al. (2016) The poly(ADP-ribose)-dependent chromatin remodeler Alc1 induces local chromatin relaxation upon DNA damage. *Mol Biol Cell* 27: 3791–3799.
56. Smith R, Sellou H, Chapuis C, et al. (2018) CHD3 and CHD4 recruitment and chromatin remodeling activity at DNA breaks is promoted by early poly(ADP-ribose)-dependent chromatin relaxation. *Nucleic Acids Res* 46: 6087.
57. Ciccarone F, Valentini E, Zampieri M, et al. (2015) 5mC-hydroxylase activity is influenced by the PARylation of TET1 enzyme. *Oncotarget* 6: 24333–24347.
58. Ciccarone F, Valentini E, Bacalini MG, et al. (2014) Poly(ADP-ribosylation) is involved in the epigenetic control of TET1 gene transcription. *Oncotarget* 5: 10356–10367.
59. Chou DM, Adamson B, Dephoure NE, et al. (2010) A chromatin localization screen reveals poly (ADP ribose)-regulated recruitment of the repressive polycomb and NuRD complexes to sites of DNA damage. *P Natl Acad Sci USA* 107: 18475–18480.
60. Luijsterburg MS, Dinant C, Lans H, et al. (2009) Heterochromatin protein 1 is recruited to various types of DNA damage. *J Cell Biol* 185: 577–586.
61. Abu-Zhayia ER, Awwad SW, Ben-Oz B, et al. (2017) CDYL1 fosters double-strand break-induced transcription silencing and promotes homology-directed repair. *J Mol Cell Biol* 1: 1.
62. D'Alessandro G, Fagagna FDD (2017) Transcription and DNA damage: holding hands or crossing swords? *J Mol Biol* 429: 3215–3229.
63. Puc J, Aggarwal AK, Rosenfeld MG (2017) Physiological functions of programmed DNA breaks in signal-induced transcription. *Nat Rev Mol Cell Bio* 18: 471–476.
64. Ju BG, Lunyak VV, Perissi V, et al. (2006) A topoisomerase IIbeta-mediated dsDNA break required for regulated transcription. *Science* 312: 1798–1802.
65. Madabhushi R, Gao F, Pfenning AR, et al. (2015) Activity-induced DNA breaks govern the expression of neuronal early-response genes. *Cell* 161: 1592–1605.
66. Perillo B, Ombra MN, Bertoni A, et al. (2008) DNA oxidation as triggered by H3K9me2 demethylation drives estrogen-induced gene expression. *Science* 319: 202–206.
67. Puc J, Kozbial P, Li W, et al. (2015) Ligand-dependent enhancer activation regulated by topoisomerase-I activity. *Cell* 160: 367–380.
68. Baranello L, Wojtowicz D, Cui K, et al. (2016) RNA polymerase II regulates topoisomerase 1 activity to favor efficient transcription. *Cell* 165: 357–371.
69. Bunch H, Lawney BP, Lin YF, et al. (2015) Transcriptional elongation requires DNA break-induced signalling. *Nat Commun* 6: 10191.
70. Marnef A, Cohen S, Legube G (2017) Transcription-coupled DNA double-strand break repair: active genes need special care. *J Mol Biol* 429: 1277–1288.
71. Huertas P, Aguilera A (2003) Cotranscriptionally formed DNA:RNA hybrids mediate transcription elongation impairment and transcription-associated recombination. *Mol Cell* 12: 711–721.
72. Sollier J, Stork CT, Garc á-Rubio ML, et al. (2014) Transcription-coupled nucleotide excision repair factors promote R-loop-induced genome instability. *Mol Cell* 56: 777–785.
73. Méivier R, Gallais R, Tiffoche C, et al. (2008) Cyclical DNA methylation of a transcriptionally active promoter. *Nature* 452: 45–50.

74. Li J, Wu X, Zhou Y, et al. (2018) Decoding the dynamic DNA methylation and hydroxymethylation landscapes in endodermal lineage intermediates during pancreatic differentiation of hESC. *Nucleic Acids Res* 46: 2883–2900.
75. Zhang Y, Zhang D, Li Q, et al. (2016) Nucleation of DNA repair factors by FOXA1 links DNA demethylation to transcriptional pioneering. *Nat Genet* 48: 1003–1013.
76. Boque-Sastre R, Soler M, Oliveira-Mateos C, et al. (2015) Head-to-head antisense transcription and R-loop formation promotes transcriptional activation. *P Natl Acad Sci USA* 112: 5785–5790.



AIMS Press

© 2018 the Author(s), licensee AIMS Press. This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>)