

Landfills and preterm birth in the Guadeloupe archipelago (French West Indies) a spatial cluster analysis

Marion Istvan, Florence Rouget, Leah Michineau, Christine Monfort, Luc Multigner, Jean-François Viel

► To cite this version:

Marion Istvan, Florence Rouget, Leah Michineau, Christine Monfort, Luc Multigner, et al.. Landfills and preterm birth in the Guadeloupe archipelago (French West Indies) a spatial cluster analysis. Tropical Medicine and Health, Japanese Society of Tropical Medicine, 2019, 47, pp.4. 10.1186/s41182-018-0130-9 . hal-02019226

HAL Id: hal-02019226

<https://hal-univ-rennes1.archives-ouvertes.fr/hal-02019226>

Submitted on 30 Sep 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

SHORT REPORT

Open Access



Landfills and preterm birth in the Guadeloupe archipelago (French West Indies): a spatial cluster analysis

Marion Istvan¹, Florence Rouget¹, Léah Michineau², Christine Monfort³, Luc Multigner³ and Jean-François Viel^{1*} 

Abstract

Background: A high rate of preterm birth is observed in the Guadeloupe archipelago (French West Indies), raising the hypothesis of harmful environmental exposures, including landfilling. Our aim was to evaluate whether preterm births cluster around the three main open landfills located in this area.

Methods: The study population consisted of 911 women enrolled in the Timoun mother-child cohort (2004–2007). Home addresses during pregnancy and locations of landfills were geocoded. To test for the presence of preterm birth clusters around each dumpsite, we used a focused cluster test specifically designed to detect spatial clustering around point sources.

Results: A total of 144 (15.8%) preterm births were observed among 911 births. Using the term births ($n = 767$) as controls, a significant cluster was identified within 2 km around the Saint-François landfill with a relative risk (RR) of 4.82 ($p = 0.04$). No clusters were found around the other two landfills (RR = 2.01, $p = 0.26$ and RR = 1.06, $p = 0.64$, for La Gabarre and Baillif, respectively).

Conclusion: The paucity of data available on open landfill sites regarding waste quantities, composition, and changes over time precludes any site-specific interpretation because of the variable degree of possible emissions. This result has to be confirmed in other tropical island environments where waste management has become a major concern with the potential to negatively impact the environment and public health.

Keywords: Preterm birth, Environmental exposure, Landfill, Spatial clustering, French West Indies

Background

A high rate of preterm birth is observed in the Guadeloupe archipelago (part of the French West Indies): 15.8% (2005–2008) versus 5.5% in mainland France (2010) [1, 2]. This high preterm birth rate cannot be fully explained by the African origin of its population, leaving room for unknown or unstudied risk factors, including environmental exposures.

Waste management has always been a crucial issue in island environments due to limited land space and increased volumes and diversity of solid waste (due to rapid population growth, heavy reliance on imported goods, lack of recycling initiatives, or poor waste collection systems).

In this context, landfilling is the major (if not the only) option for waste management. However, landfills can represent a health risk for residents because of exposure to pollutants through different pathways: inhalation of substances emitted by the site, contact with water or polluted soil, and consumption of contaminated foodstuffs or drinking water. In this respect, residential proximity to landfills has been previously linked to preterm births or congenital malformations, although several alternative explanations (including ascertainment bias and residual confounding) cannot be excluded [3, 4].

Spatial epidemiology can give important clues when searching for putative environmental causes of a disease [5, 6]. In particular, cluster analysis allows researchers to assess whether people with the disease have lived closer to one another than would be expected by chance during a specific time period. Local clustering methods can

* Correspondence: jean-francois.viel@univ-rennes1.fr

¹Univ Rennes, CHU Rennes, Inserm, EHESP, Irset (Institut de recherche en santé, environnement et travail) - UMR_S 1085, 2 rue Henri Le Guilloux, F-35000 Rennes, France

Full list of author information is available at the end of the article



be differentiated as either non-focused or focused tests. Non-focused methods aim at detecting clusters without a priori knowledge of their location. Subsequently, hypotheses about exposures potentially responsible for such clusters can be made. Focused cluster tests are used when there is a priori knowledge about the location of hypothesized clusters (such as suspected or known sources of environmental contaminants) assuming that disease clusters follow the spatial patterns of environmental contaminant dispersion [7].

Because three main open landfills were used for solid waste disposal in the Guadeloupe archipelago, our aim was to evaluate whether preterm births cluster around these predefined locations.

Methods

Guadeloupe is an archipelago located in the Leeward Islands in the Caribbean. It covers an area of 1628 km² with a population of 450,000 inhabitants. This study relies on the Timoun mother-child cohort fully described elsewhere [1]. Between 2004 and 2007, 1068 women residing in Guadeloupe for more than 3 years were enrolled during the third trimester of pregnancy. In the current study, we excluded women not born in the Caribbean ($n = 110$) to reduce heterogeneity as African ancestry is a strong risk factor for preterm birth [8], cases involving multiple births ($n = 25$), severe birth defects ($n = 8$), and induced pregnancies after fertility treatment ($n = 15$), resulting in a study population of 911 women (one case involved both fertility treatment and multiple births). Gestational age in weeks was estimated by the obstetricians in charge of follow-ups. It was based on the first day of the last menstrual period and was confirmed or corrected by ultrasound. Preterm birth was defined as a birth before 37 completed weeks of gestational age.

The three main (and official) open landfills in Guadeloupe (namely, La Gabarre, Saint-François, and Baillif) opened as temporary structures in the 1970s (Table 1, Fig. 1). They then received increasing amounts of both municipal waste and hazardous waste (scrap cars, chemical waste, hospital waste, metal waste, industrial waste) in an uncontrolled and uncovered manner [9].

Home addresses during pregnancy and locations of dumpsites were geocoded. To test for the presence of

preterm birth clusters around each landfill, we used a (purely spatial) focused cluster test specifically designed to detect spatial clustering around point sources, assuming a Bernoulli distribution and using a grid file with only a single grid point reflecting the coordinates of the focus of interest [10]. The program scans for clusters of geographic size between zero and some upper limit. As a compromise between the need for spatial precision and the limited data available on landfill sites, the maximum radius was set as 2 km around each site, at the likely limit of dispersion for landfill emissions to include both air and water pathways according to a WHO report [11].

For hypothesis testing, we generated 9999 Monte Carlo replications to ensure good statistical power. Focused cluster tests were performed around each landfill in turn using the SaTScan software program (Kulldorff M., Boston, and Information Management Services, Inc., Calverton, Maryland) [12].

Results

In the study population ($n = 911$), a total of 144 (15.8%) preterm births were observed in as many different locations. Selected socio-demographic and lifestyle characteristics of the study population are reported in Table 2. No significant differences were found between mothers of preterm infants and mothers of full-term infants.


Using the term births ($n = 767$) as controls, a significant cluster was identified within 2 km around the Saint-François landfill with a relative risk (RR) of 4.82 ($p = 0.04$) (Table 3). The number of observed preterm births was twice the expected number around the La Gabarre dumpsite (RR = 2.01), yet this difference was not statistically significant ($p = 0.26$). No risk was highlighted in the vicinity of the Baillif landfill ($p = 0.64$).

Discussion

We found a significant cluster of preterm birth in the vicinity of the Saint-François landfill, which was located in a less densely populated area than the other two landfills (as reflected by the numbers of Timoun newborns and expected cases). Potential socio-demographic confounding factors did not appear to account for this observation.

Table 1 Description of the three main open landfills in the Guadeloupe archipelago

	La Gabarre	Saint-François	Baillif
Geographic coordinates	16.25956 N 61.54151 W	16.27477 N 61.28276 W	16.03139 N 61.74791 W
Municipality	Les Abymes	Saint François	Baillif
Operation	Opened in 1973 Rehabilitation in 2009	Opened in 1974 Closed in 2010 Rehabilitation in 2012	Opened in 1974 Closed in 2008 Rehabilitation in 2008
Acreeage	37 ha	8.6 ha	7 ha



The present study has several strengths. First, because of the absence of heavy industries in the Guadeloupe archipelago, no confounding by their effluents is to be predicted. Moreover, no other pollution sources are documented in the areas that surround the three dumpsites. Second, preterm birth was carefully and uniformly characterized within a cohort framework. Both the residential addresses of the pregnant women and the landfill locations were known with sufficient precision to allow appropriate geocoding. Third, this study benefited from the well-established advantages of the focused cluster test to assess whether cases are closer to the sources than expected. Addressing a specific hypothesis of concern, this inferential method has a higher power compared to other spatial methods [13].

This study is not without limitations. First, only authorized landfills were considered; however, illegal, uncontrolled landfills that receive waste without any selection at the origin may be an additional health concern. Second, the distance from the dumpsite was used as a proxy for exposure as it may reflect and integrate a broad range of contaminants and different routes of

exposure (not only air but also the contamination from soil or groundwater in the vicinity of the landfill). This buffer-based approach considers that emissions from a landfill are uniformly dispersed in all directions. It is worth noting that a 2-km radius, which is within plausible estimates of the range of chemicals dispersed from a site, has been often used to estimate exposure [13]. However, assuming that environmental exposure is equally distributed around the polluted site might be questionable for specific pollutants. Third, because of the limited number of preterm births, the possibility of insufficient statistical power to indicate clusters around the Gabarre dumpsite cannot be entirely dismissed. Unfortunately, no national birth register is available in France, and we had to rely upon a locally run mother-child cohort. Fourth, a limitation inherent to any landfill is the wide variety of pollutants, exposure pathways and exposure scenarios, entailing a large complexity and difficulty in estimating the health risks possibly involved in its vicinity. Unfortunately, specific characteristics of the dumpsites (e.g., age, waste streams

Table 2 Socio-demographic characteristics of the study population according to the pregnancy outcome (Timoun cohort, $n = 911$, Guadeloupe archipelago, 2004–2007) (from Rouget et al. 2013)

	Term births $n = 767$	Preterm births $n = 144$	HR (95% CI)
Maternal place of birth			
Guadeloupe or Martinique	669	128	Reference
Other Caribbean islands	98	16	0.9 (0.5–1.5)
Maternal age (years)			
< 20	63	11	1.0 (0.5–1.9)
20–34	437	80	Reference
≥ 35	247	53	1.3 (0.9–1.9)
Years of education			
< 5	49	13	1.7 (0.9–3.0)
5–12	564	101	Reference
> 12	154	30	1.1 (0.7–1.6)
Mother's employment during pregnancy			
No	449	83	Reference
Yes	318	61	1.0 (0.7–1.4)
Tobacco smoking during pregnancy			
No	737	138	Reference
Yes	30	6	0.9 (0.4–2.2)
Alcohol drinking during pregnancy			
No	745	141	Reference
Yes	22	3	0.8 (0.2–2.4)

HR hazard ratio, CI confidence interval

disposed of, waste quantities, operating practices, changes over time, conditions at the bottom layer) are generally not known and relevant data sets are simply not available. In this regard, further elaborating on the apparently discrepant results (the significant cluster is not observed in the vicinity of the largest landfill, i.e., La Gabarre) would be unwise.

To the best of our knowledge, the current study is the first to report on this topic in a tropical island environment. The available scientific evidence has been recently reviewed by Kihal-Talantikite et al. [13], revealing scarce literature for preterm birth. An increased risk was found with residential proximity to a hazardous waste landfill

Table 3 Clusters of preterm births in the vicinity of open landfills (Timoun cohort, Guadeloupe archipelago, 2004–2007, 2-km radius, focused tests)

	La Gabarre	Saint-François	Baillif
Timoun newborns	16	4	18
Observed cases	5	3	3
Expected cases	2.53	0.63	2.85
Relative risk	2.01	4.82	1.06
P value	0.26	0.04	0.64

in New Jersey (USA) (OR [odds ratio] = 2.10; 95% CI [confidence interval] 1.01–4.36) [14] and to a hazardous waste site in Nova Scotia (Canada) (RR [relative risk] = 1.13; 95% CI 1.04–1.22) [15]. Conversely, no association was found in the vicinity of a municipal solid waste landfill in Quebec (Canada) (OR = 1.00; 95% CI 0.88–1.13) [16] or around open dumpsites in Alaska (USA) (OR = 1.09; 95% CI 0.78–1.51) [17]. One must, however, keep in mind that the complex nature of the landfill ecosystem makes it difficult to compare studies and draw lessons from other locations.

Conclusion

Waste management in island environments has become a major concern with the potential to negatively impact development activities (including tourism and trade), food supplies, the environment, and public health. The excess risk of preterm birth observed around one Guadeloupien dumpsite supports these concerns but will need to be confirmed in other tropical island environments. In the meantime, preterm births might be considered in any risk assessment of landfill disposal sites in tropical belt.

Abbreviations

OR: Odds ratio; RR: Relative risk

Acknowledgements

Authors are grateful to Sylvaine Cordier for implementing the TIMOUN cohort.

Funding

The Timoun mother-child cohort is supported by grant from the General Health Directorate (DGS RMC11129NNA). This study was conducted in the framework of the COST Action IS1408 "Industrially Contaminated Sites and Health Network (ICSHNet)".

Availability of data and materials

Data are available from the corresponding author on reasonable request.

Authors' contributions

JFV generated the idea for the paper and prepared an analytical plan. CM and LM were responsible for data management and geocoding. MI and JFV conducted all data analyses. JFV prepared the first draft, and FR and LM critically revised it. All authors read and approved the final manuscript.

Ethics approval and consent to participate

The Guadeloupien Ethics committee for studies involving human subjects approved the study. Written informed consent was provided by each participant.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹Univ Rennes, CHU Rennes, Inserm, EHESP, Irset (Institut de recherche en santé, environnement et travail) - UMR_S 1085, 2 rue Henri Le Guilloux, F-35000 Rennes, France. ²Univ Rennes, Inserm, EHESP, Irset (Institut de recherche en santé, environnement et travail) - UMR_S 1085, F-97000

Pointe-à-Pitre, France. ³Univ Rennes, Inserm, EHESP, Irset (Institut de recherche en santé, environnement et travail) - UMR_S 1085, F-35000 Rennes, France.

Received: 6 November 2018 Accepted: 18 December 2018

Published online: 09 January 2019

References

- Rouget F, Lebreton J, Kadhel P, Monfort C, Bodeau-Livinec F, Janky E, Multigner L, Cordier S. Medical and sociodemographic risk factors for preterm birth in a French Caribbean population of African descent. *Matern Child Health J*. 2013;17:1103–11.
- Blondel B, Lelong N, Kermarrec M, Goffinet F. National Coordination Group of the National Perinatal Surveys. Trends in perinatal health in France from 1995 to 2010. Results from the French National Perinatal Surveys. *J Gynecol Obstet Biol Reprod*. 2012;41:e1–e15.
- Mattiello A, Chiadini P, Bianco E, Forgione N, Flammia I, Gallo C, Pizzuti R, Panico S. Health effects associated with the disposal of solid waste in landfills and incinerators in populations living in surrounding areas: a systematic review. *Int J Public Health*. 2013;58:725–35.
- World Health Organization, Regional office for Europe, 2015. Waste and human health: Evidence and needs - WHO Meeting Report, 2015. http://www.euro.who.int/__data/assets/pdf_file/0003/317226/Waste-human-health-Evidence-needs-mtg-report.pdf. Accessed 14 Dec 2018.
- Pfeiffer D, Robisons T, Stevenson M, Stevens K, Rogers D, Clements A. *Spatial analysis in epidemiology*. Oxford: Oxford University Press; 2008.
- Lawson AB, Banerjee S, Haining RP, Ugarte MD. *Handbook of spatial epidemiology*. Chapman & Hall/CRC handbooks of modern statistical methods. Boca Raton: CRC Press/Taylor & Francis; 2016.
- Lawson AB, Biggeri A, Williams FLR. A review of modelling approaches in health risk assessment around putative sources. In: Lawson AB, Böhning D, Lesaffre E, Biggeri A, Viel JF, Bertollini R, editors. *Disease mapping and risk assessment for public health*. Chichester: Wiley; 1999. p. 231–245.
- Muglia LJ, Katz M. The enigma of spontaneous preterm birth. *NEJM*. 2010;362:529–35.
- Plocoste T, Jacoby KS. Impact of a new legislation on volatile organic compounds emissions in an open landfill in tropical insular climate. *Int J Waste Resour*. 2016;6:236.
- Kulldorff M, Nagarwalla N. Spatial disease clusters: detection and inference. *Stat Med*. 1995;14:799–810.
- World Health Organization. Methods of assessing risk to health from exposure to hazards released from waste landfills. Report from a WHO meeting Lodz, Poland, 10–12 April 2000. Bilthoven: WHO Regional Office for Europe, European Centre for Environment and Health; 2001. <http://apps.who.int/iris/bitstream/handle/10665/108362/E71393.pdf?sequence=1&isAllowed=y>. Accessed 14 Dec 2018
- Kulldorff M. SaTScan v9.6: software for the spatial and space-time statistics. 2018. <https://www.satscan.org>. Accessed 14 Dec 2018.
- Kihal-Talantikite W, Zmirou-Navier D, Padilla C, Deguen S. Systematic literature review of reproductive outcome associated with residential proximity to polluted sites. *Int J Health Geogr*. 2017;16:20.
- Berry M, Bove F. Birth weight reduction associated with residence near a hazardous waste landfill. *Environ Health Perspect*. 1997;105:856–61.
- Dodds L, Seviour R. Congenital anomalies and other birth outcomes among infants born to women living near a hazardous waste site in Sydney. *Nova Scotia Can J Public Health*. 2001;92:331–4.
- Goldberg MS, Goulet L, Riberdy H, Bonvalot Y. Low birth weight and preterm births among infants born to women living near a municipal solid waste landfill site in Montreal, Quebec. *Environ Res*. 1995;69:37–50.
- Gilbreath S, Kass PH. Adverse birth outcomes associated with open dumpsites in Alaska Native Villages. *Am J Epidemiol*. 2006;164:518–28.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

