

Women and COPD do we need more evidence?

Christophe Gut-Gobert, Arnaud Cavallès, Adrien Dixmier, Stéphanie Guillot, Stéphane Jouneau, Christophe Leroyer, Sylvain Marchand-Adam, David Marquette, Jean-Claude Meurice, Nicolas Desvigne, et al.

► **To cite this version:**

Christophe Gut-Gobert, Arnaud Cavallès, Adrien Dixmier, Stéphanie Guillot, Stéphane Jouneau, et al.. Women and COPD do we need more evidence?. European Respiratory Review, European Respiratory Society, 2019, 28 (151), 10.1183/16000617.0055-2018 . hal-02061961

HAL Id: hal-02061961

<https://hal-univ-rennes1.archives-ouvertes.fr/hal-02061961>

Submitted on 8 Jul 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Women and COPD: do we need more evidence?

Christophe Gut-Gobert¹, Arnaud Cavallès², Adrien Dixmier³, Stéphanie Guillot⁴, Stéphane Jouneau^{5,6}, Christophe Leroyer¹, Sylvain Marchand-Adam⁷, David Marquette⁸, Jean-Claude Meurice⁹, Nicolas Desvigne¹⁰, Hugues Morel³, Christine Person-Tacnet¹¹ and Chantal Raheison¹²

Affiliations: ¹G.E.T.B.O. (Groupe d'Etude de la Thrombose de Bretagne Occidentale), Université Européenne de Bretagne, Université de Brest, EA3878, IFR148, Hôpital La Cavale Blanche, Département de Médecine Interne et Pneumologie, Brest, France. ²Institut du Thorax, CHU de Nantes, Dept of Pulmonology, Nantes, France. ³Dept of Pulmonology, Orléans Regional Hospital, Orléans, France. ⁴Unité d'Explorations Fonctionnelles Respiratoires, CHRU Rennes, Rennes, France. ⁵Service de Pneumologie, Hôpital Pontchaillou, Rennes, France. ⁶IRSET UMR 1085, Université de Rennes 1, Rennes, France. ⁷Université François Rabelais Faculté de Médecine de Tours, Inserm 1100, CHRU de Tours Service de Pneumologie, Tours, France. ⁸Dept of Pulmonary Medicine, Centre Hospitalier Bretagne Atlantique, Vannes, France. ⁹Dept of Pulmonology Centre Hospitalier de l'Université de Poitiers, Poitiers, France. ¹⁰Novartis Pharma, Scientific Operations, Rueil-Malmaison, France. ¹¹Dept of Pulmonology, CHU Angers, Angers, France. ¹²Service des Maladies Respiratoires, CHU Bordeaux, Epicene U1219 Université de Bordeaux, Bordeaux, France.

Correspondence: Christophe Gut-Gobert, Hôpital La Cavale Blanche, Département de Médecine Interne et Pneumologie, Boulevard Tanguy Prigent, 29609 Brest Cedex 2, France.
E-mail: Christophe.gut-gobert@chu-brest.fr



@ERSpublications

The face of COPD is increasingly female. We need more evidence and a change in how the disease is managed. <http://ow.ly/zueL30mWqLS>

Cite this article as: Gut-Gobert C, Cavallès A, Dixmier A, *et al.* Women and COPD: do we need more evidence? *Eur Respir Rev* 2019; 28: 180055 [<https://doi.org/10.1183/16000617.0055-2018>].

ABSTRACT The increasingly female face of chronic obstructive pulmonary disease (COPD) prevalence among women has equalled that of men since 2008, due in part to increased tobacco use among women worldwide and exposure to biomass fuels. This finding is supported by a number of characteristics. There is evidence of susceptibility to smoking and other airborne contaminants, along with epidemiological and phenotypic manifestations. COPD has thus become the leading cause of death in women in the USA. The clinical presentation is characterised by increasingly pronounced dyspnoea with a marked tendency towards anxiety and depression, undernutrition, nonsmall cell lung cancer (especially adenocarcinoma) and osteoporosis. Quality of life is also more significantly impacted. The theories advanced to explain these differences involve the role played by oestrogens, impaired gas exchange in the lungs and smoking habits. While these differences require appropriate therapeutic responses (smoking cessation, pulmonary rehabilitation, long-term oxygen therapy), barriers to the treatment of women with COPD include greater under-diagnosis than in men, fewer spirometry tests and medical consultations. Faced with this serious public health problem, we need to update and adapt our knowledge to the epidemiological changes.

Provenance: Publication of this peer-reviewed article was sponsored by Novartis, France (article sponsor, *European Respiratory Review* issue 151).

Received: June 11 2018 | Accepted after revision: Sept 21 2018

Copyright ©ERS 2019. ERR articles are open access and distributed under the terms of the Creative Commons Attribution Non-Commercial Licence 4.0.

Introduction

Chronic obstructive pulmonary disease (COPD) is no longer a respiratory disease that predominantly affects men, and less still the elderly. It encompasses several phenotypes and a specific female phenotype cannot be ruled out.

There is a significant delay in diagnosis because women do not seek medical attention for dyspnoea but for fatigue, resulting in major psychosocial consequences and impaired quality of life [1]. The experience of this disease is difficult for these women because COPD is an invisible disease that can cause misunderstanding among family and friends.

In the USA, COPD is the leading cause of death among female smokers [2, 3], ahead of lung cancer and cardiovascular diseases. Under-diagnosis in primary care medicine is still common [4] because the medical profession needs to change its perception and teaching of COPD as a male disease, a misperception that has stuck with them since their early years of medical school. COPD has female-specific clinical manifestations and comorbidities.

The Fletcher curve is still used today to plot the rate of forced expiratory volume in 1 s (FEV₁) decline, despite only male patients being used and despite the identified decline in females [5, 6].

The rise of smoking among women

Catherine de Medici was the first woman in French history to use tobacco, yet consumption by women would long remain marginal. A whiff of scandal surrounded female smokers such as Georges Sand, who would visit literary salons with a pipe or cigar. Not until the 20th century did cigarettes become widespread among women. Female smoking trends rose in tandem with the changing status of women.

Smoking trends among women seem to follow the same pattern as the trend in male smoking after a lag of nearly a century [7]; smoking first started among the upper classes and then spread to working-class men, who became the primary tobacco users. This trend went hand-in-hand with a sharp increase in the number of smokers. It is now thought that smoking among men has reached its peak and is sociologically in decline. For women, this trend began in the interwar years. By smoking, women signified their membership of a privileged social group and their desire for emancipation. This was the fashion statement of the flappers. Between 1950 and 1970, cigarettes permeated every stratum of society and became commonplace.

The phenomenon was aided by advertising [8]. The tobacco industry sought to portray cigarettes as a means of seduction, first by lauding the slimming effects of tobacco and then by filming glamorous actresses like Marlene Dietrich. Many American actresses signed up to these manipulative practices, and their European counterparts followed suit soon after.

However, an encouraging trend is emerging from the West: in the USA, smoking among women is viewed negatively and its harmful health consequences are recognised. Sociologically, it is in decline. And now, Europe seems to be catching up. We can expect a more rapid rate of decline in smoking prevalence among women than among men.

Confronted by this risk of declining tobacco use in Europe, the tobacco industry is looking elsewhere for market share. It has shifted its interest to developing countries, where it is still possible to win over large numbers of new consumers and where smoking prevention measures are underdeveloped. In these countries, women and young people are prime targets. A report by the World Health Organization (WHO) also points to evidence that tobacco advertising increasingly targets young girls [9]. The messages conveyed by advertising are identical to those which have been tried and tested in Western countries; emphasis is placed on the social acceptability of women who smoke and on smoking as a sign of emancipation. An article by HITCHMAN and FONG [10] shows that there is a close correlation between female and male smoking rates and female empowerment. This correlation is clearly associated with the level of economic development. Will the trend towards greater female empowerment also lead to a tobacco epidemic among women [10]? To highlight the significance of the risk posed by rising tobacco use among women worldwide, the theme for the World No Tobacco Day organised by WHO in 2010 was "Gender and tobacco with an emphasis on marketing to women".

Yet the increase in female smoking is not necessarily inevitable. During the 20th century, the prevalence of female smoking declined in China. The smoking rate among women was 25% at the start of the century and increased further in the 1930s, but then experienced a decline in the second half of the 20th century. This reversal coincided with the emergence of new sociocultural norms which frowned on smoking among women. These new norms were conveyed through movements based on traditional values extolling behaviours that promoted a healthier mind and body [11]. The Global Adult Tobacco Survey (GATS) was conducted by the WHO to assess household tobacco consumption in different countries around the world.

In 2010, it was conducted among Chinese households. It showed that 52.9% of men were reported to smoke while only 2.4% of women smoked [12].

Epidemiology of COPD in women

COPD remains under-diagnosed, especially in women [13]. Two studies carried out in North America and Spain show similar results. Female smokers who visit a physician are more than one-third less likely to be diagnosed with COPD than male smokers. Spirometry testing and referral to a pulmonologist are less common for women [4, 14, 15]. Women reportedly suffer a delay in the diagnosis of COPD, explainable in part by voluntary postponement of access to a consultation or in some patients by the prevalence of symptoms of fatigue or depression that point to a different type of treatment [16]. Allowing for these reservations, the prevalence of COPD is currently estimated, depending on the country, to range from 4.5% to 10.2% [17, 18]. Although a 2006 meta-analysis assessed prevalence at 9.8% among men and 5.6% among women [19], the latest data mitigate this difference. The Burden of Obstructive Lung Disease (BOLD) study, in which prevalence was found to be 11.8% among men and 8.5% among women, reports a higher prevalence at the moderate stages among women in the USA, Austria, Iceland and Australia [20]. In 2008, the WHO estimated that 168 million men and 160 million women were affected worldwide [21]. An analysis of data from 1998 to 2009 shows that the prevalence of COPD in the USA increased in women and decreased in men. The same trends were found in Canada, the Netherlands and Austria [22–24]. In the USA, the number of women diagnosed with emphysema has outnumbered that of men since 2011 [24]. An increase in smoking among women in all countries, including France, has been observed in parallel with the exposure of women to indoor smoke as a result of burning biomass fuels in developing countries [25–28]. Women are more susceptible to tobacco, having a more severe disease despite lower cumulative tobacco consumption, earlier onset of COPD, and faster decline in FEV₁, but conversely obtain greater benefit from cessation of exposure [24, 29]. This susceptibility to airborne contaminants involves not only tobacco. CELLI *et al.* [30] found that 80% of nonsmoker COPD patients are women, probably on account of exposure to biomass fuels.

In Spain, the ECCO study evaluated the comorbidity in COPD patients hospitalised in internal medicine departments. The mean number of comorbidities was 3.7 in men and 1.8 in women ($p < 0.05$). Women had a lower prevalence of ischaemic heart disease and alcoholism, but presented more frequently with heart failure, osteoporosis and diabetes [31]. In a recent Swedish study of oxygen-dependent COPD patients, women had significantly higher hypertension, higher rates of depression and higher rates of osteoporosis. The presence of a comorbidity was an independent predictor of mortality and the effect was similar for both sexes; nevertheless, mortality was lower among women [32, 33]. Several authors have found a higher frequency of depression in women and a greater impact on quality of life [34]. Dyspnoea appears strongly associated with depression in women [34–37]. In France, a working group studying a real-life COPD patient population showed that, for a given age and level of airflow obstruction, women have a higher BODE index (body mass index, airflow obstruction, dyspnoea and exercise capacity) due to a lower body mass index and more pronounced dyspnoea, suggesting a poorer overall prognosis [38]. Considering the pulmonary hypertension phenotype in COPD, it is equally present in both sexes as shown by the ASPIRE register, whereas there is a female predominance in pulmonary hypertension in group one [39].

Between 1990 and 2010, COPD became the third leading cause of death in the USA. The COPD mortality rate declined among men in the USA between 1999 (57.0 per 100 000) and 2006 (46.4 per 100 000), whereas there was no significant change in mortality rates for women (35.3 per 100 000 in 1999 and 34.2 per 100 000 in 2006). In 2000, the number of women dying from COPD surpassed the number of men [40]. A recent assessment of mortality rates due to COPD in the cohorts from the National Health and Nutrition Examination Survey (NHANES) I and NHANES III in the USA showed a smaller decrease in the mortality rate among women with moderate or severe COPD than among men (3.0% *versus* 17.8%) [41]. Thus, the survival advantage of women with COPD has been diminishing [3].

Is there a specific COPD-related phenotype in women?

Women with COPD appear to have a different clinico-radiological phenotype from that of men. In smoking- or FEV₁-matched men and women, women present with more symptoms such as dyspnoea [42, 43] or cough [14]. Even if they produce less sputum than men [14, 44], they are more likely to have a chronic bronchitic phenotype. This seems to confirm the histopathological analyses [42]; although women have less severe COPD than men, they have thicker small airway walls (<2 mm in tissue removed during lung volume reduction surgery). Conversely, a computed tomography analysis showed thicker bronchial walls in men [45]. The significance of emphysema in women is also under discussion [19]. In the National Emphysema Treatment Trial (NETT) study of patients with severe COPD before lung volume reduction surgery [42] or in a lung cancer screening study [46], smoking-matched women had less emphysema than men. Women presenting with the same airflow obstruction and lower smoking rates are usually younger

and have more frequent exacerbations [47]. This susceptibility to tobacco smoke in women favours a faster rate of FEV₁ decline [48–51].

Several cohort studies of patients in the Genetic Epidemiology of COPD (COPD Gene) study support the existence of a sex-related genetic component in COPD onset, including a higher risk of early-onset severe COPD in female smokers [52, 53]. The X chromosome may be involved since COPD in a mother increases the risk of COPD onset in a daughter who smokes [52]. Anatomical factors and the concept of dysanapsis may explain why women are at increased risk of airway diseases [54]. During adolescence, female airway growth is small relative to lung growth to men, whose bronchopulmonary growth is more homogeneous, hence the impaction of inhaled substances on a smaller surface [55]. There is greater particle deposition in healthy women than in men [56] especially in the proximal airways [57]. Another hypothesis is that local inflammatory response in the airways is greater in female than in male smokers. Indeed, respiratory bronchiolitis may occur early in young female smokers [58]. Narrowing of the small airways in women with emphysema is associated with greater bronchial wall thickening [42]. There are also significant differential expressions of certain proteins in alveolar macrophages from male and female smokers with COPD [59]. In women with COPD, impairment in lysosomal function would result in macroautophagy inhibition in alveolar macrophages, thereby contributing to airway inflammation. Leptin, a pro-inflammatory mediator present in the airways of female COPD patients, stimulates the production of certain cytokines; its elevated serum level is thought to be associated with an increased prevalence of COPD in women but not in men [60]. In patients matched for airway obstruction, serum levels of interleukin-16 and vascular endothelial growth factor were significantly higher in female COPD patients than in male COPD patients [61]. Bronchial hyperresponsiveness (BHR) is more common in women than in men, and excessive smoking (>20 g·day⁻¹) is associated with a higher risk of significant BHR only in women [62]. BHR is a risk factor for accelerating lung function decline in smokers of both sexes but the rate of FEV₁ decline is more rapid in women [63]. The manner in which smoke is inhaled can change the level of BHR; women tend to inhale more deeply, thus exacerbating BHR [64]. Hormonal factors play a complex role at bronchopulmonary level. The role of oestrogens is well established at several levels: they are involved in maintaining cell structures and in lung elastic recoil that keeps airways open [65]. Accordingly, the level of airway obstruction in post-menopausal women not receiving hormone replacement therapy is greater than in those receiving hormone replacement therapy [66]. The potential anti-oestrogenic effect of cigarette smoke is known [67], and may contribute to impaired lung function in female smokers. After smoke inhalation, chemicals are metabolised in phase I and II, largely mediated by cytochrome P450 (CYP) enzyme inducers. Oestradiol upregulates CYP enzymes of phase I without altering phase II. It increases oxidative stress in the airways making female lungs more susceptible to oxidant damage in response to cigarette smoke [68]. Finally, oestradiol induces MUC5B mucin gene expression in airway epithelial cells in healthy subjects, hypersecretion of the protein being common in chronic respiratory disease [69]. Testosterone aggravates emphysema of the male lung [68]. It increases metalloprotease and elastase effects and alveolar destruction by stimulating neutrophils response.

Nonsmoking-related COPD

COPD among nonsmokers is an increasing area of study and accounts for 25–45% of patients with COPD [70]. The prevalence of nonsmoking-related COPD is higher among women than among men [24, 71–79]. In the international BOLD study, among 4291 nonsmokers aged >40 years, 6.6% had COPD stage I and 5.6% had COPD stage II or higher [80]. Among smokers, the prevalence of COPD was higher in men (17.1% *versus* 13.2%, *p*=0.001) while among nonsmokers, the age distribution was similar in both sexes (5.2% *versus* 6.2%, *p*=0.142) [80].

Exposure to biomass smoke is thought to be the main risk factor for nonsmoking-related COPD worldwide and occurs mainly in women because they cook in poorly ventilated homes in developing countries [24, 70]. A Chinese study of a large group of nonsmokers found COPD prevalence to be 4% in women and 5.1% in men [81]. The women were slightly younger than the men (51.1 years *versus* 54.4 years) and 92% of them were responsible for cooking compared with 29–52% of the men. Cooking and the use of coal or wood for heating were associated with a higher prevalence of airway obstruction. The risk factors for COPD were living in a rural area, older age, lower level of education, lower income, lower body mass index and a doctor-diagnosed history of tuberculosis.

Psychological and social impact of COPD: are men and women equal?

Several quality of life questionnaires are available for the study of COPD patients: Short-form (SF) 36, the Medical Research Council Questionnaire and the St George's Respiratory Questionnaire (SGRQ), the last of which is most commonly used. Several studies based on these tools report greater quality of life deterioration in women than in men (table 1) [1, 24, 82, 83]. A French multicentre study [1] compared the SGRQ, the hospital anxiety and depression (HAD) scale and a motivation scale on smoking cessation

TABLE 1 Studies concerned with quality of life by sex in chronic obstructive pulmonary disease (COPD)

First author [ref.]	Country	Quality of life (tools used)	COPD stage	Females/males n	Results
DE TORRES [82]	Spain, single-centre	SGRQ	GOLD II, III	73/73	Quality of life more impaired in females Dyspnoea and arterial saturation: only factors associated with the SGRQ score, unlike males for whom exercise capacity, dyspnoea and comorbidities are found
KATSURA [83]	Japan, single-centre	SGRQ-SF-36, Philadelphia Geriatric Center morale scale	GOLD II	39/117	SGRQ: highest score for activity, impact and total score among females compared with males SF-36 highest score on perceived health and health compared with 1 year ago for females/males
KAPTEIN [84]	Netherlands	Telephone or e-mail questionnaire: intimate physical contact scale (IPCS) or Respiratory Experiences with Sexuality Profiles (RESP)	COPD, asthma	55 patients (of whom 10 females and 15 males with COPD)	Reduced sexual activity in males and deterioration in quality of sex life More seldom occurrence in women
RODRÍGUEZ-GONZÁLEZ MORO [85]	Spain	SF-12 (used during consultations with pulmonologists and treating physician)	GOLD II, III	1786/1661	Quality of life more impaired in females, particularly concerning the mental and physical aspects of SF-12
RAHERISON [1]	France, multicentre	SGRQ, HADS, Q-MAT (for active smokers)	GOLD II, III, IV	247/183	Poorer anxiety and quality of life scores in females even though lung function impairment was less severe than in males

SGRQ: St George's Respiratory Questionnaire; SF: short-form; GOLD: Global Initiative for Chronic Obstructive Pulmonary Disease; HADS: hospital anxiety and depression scale.

in men and in women. The frequency of cough, sputum and severity of dyspnoea did not differ between sexes. The anxiety score was higher and quality of life more impaired in women. In a multivariate analysis, chronic sputum was associated with a poorer quality of life. A Swedish study highlighted the impact of cough and urinary incontinence on quality of life for women and men with COPD [86]. The study placed importance on the reporting of these symptoms during the interview but did not compare the differences between men and women. Female COPD patients with a heavy smoking history seem to experience a poorer quality of life than men with the same smoking intensity [86]. Sex is overlooked in the quality of life scores. A small cohort study looked at a population of patients with COPD and asthma [84]. Patients were asked to answer two questionnaires by phone or e-mail: the Intimate Physical Contact Scale and the Respiratory Experiences with Sexuality Profiles. The female COPD patients described a poorer “quality of sex life” than asthmatic women. However, the scores appeared better than for the men with COPD, particularly in terms of activity and satisfaction. BECKLAKE and KAUFMANN [87] speculated that men attach more importance to athleticism and place less emphasis on dyspnoea as a major symptom. This societal ideal would explain some differences in terms of quality of life and symptom reporting [24]. Conversely, women are less likely than men to report sputum as a symptom. Loss of muscle mass is greater in patients with moderate-to-severe COPD [88] and muscle mass depletion impacts quality of life. This concept is under discussion and some authors suggest a link between muscle mass loss, decreased carbon monoxide diffusion capacity and static hyperinflation [89]. This would tie in with the more common emphysematous phenotype in women. Different methods are used to assess anxiety and depression in COPD patients. In terms of questionnaires, HAD is the preferred method [1, 34, 35, 90–92]; the structured interview approach or medical diagnoses are less common [90, 93].

The incidence of depression is higher in COPD patients than in the general population. Women are significantly more likely to have anxiety and/or depression than men in the COPD population (table 2) with manifestations that can be severe (phobia, panic attacks and generalised anxiety) [90]. The prevalence of depression is significantly higher among women (OR=1.76) [94]. Thus, VANFLETEREN *et al.* [91] have identified a “psychological” phenotype with a female majority of 55%.

The consequences of anxiety and depression are numerous: smoking cessation is more difficult [95]; dyspnoea is increased [94]; quality of life, as measured by the SGRQ or the SF36, is impaired [1, 34, 90, 99, 96]; and sleep disturbances are more frequent [97–99]. The rate of rejection or abandonment of pulmonary rehabilitation is increased [100, 101] and the commitment to self-management of the disease and implementation of action plans to manage exacerbations is lacking [92]. The increased use of care services is linked to an increased risk of exacerbations [35, 102, 103] and increased risk of hospital readmission at 1 year [35]. For MCGARVEY *et al.* [93], female sex and depression are among the risk factors of frequent exacerbations (>2 the previous year). For FAN *et al.* [96], 3-year mortality is significantly increased among patients with depressive symptoms. Anxiety is one of 12 comorbid conditions contributing to excess mortality among the patients concerned, with a hazard ratio of 13.76 [104].

Lung cancer and COPD in women

Nonsmall cell lung cancer (NSCLC) incidence and mortality have increased in recent years in women [105]. The incidence of NSCLC increased 4–5 times faster than the general population in patients with COPD over the period 1991–2001 in the UK, regardless of sex [106]. COPD itself is an independent risk factor for NSCLC above and beyond its association with smoking [107]. The impact of COPD on mortality is negligible in some studies [108, 109], whereas according to the study by KIRI *et al.* [106], COPD increases 3-year mortality in patients with NSCLC regardless of patient age or sex, with higher mortality rates above all in patients aged >65 years (RR=2.05 against 1.3 in patients aged <65 years). Few studies focus on women with COPD and presenting with NSCLC. In the study by IZQUIERDO *et al.* [109], 47.7% of patients with stages IIIB and IV NSCLC also had COPD, while in the study by LOGANATHAN *et al.* [105], 72.8% of men with NSCLC and 52.5% of women with NSCLC also had COPD, regardless of the TNM (classification of malignant tumours) stage of the disease. In a study of 562 North American women aged

TABLE 2 Prevalence of anxiety and depression in patients with chronic obstructive pulmonary disease (COPD) according to sex

First author [ref.]	Population	Patients n	COPD	Females %	Anxiety	Depression	Anxiety and depression
DI MARCO [34]	Italian	202	Moderate to severe	23.3	38.3/25.2	38.3/12.9	25.5/6.5
LAURIN [90]	Canadian	116	Moderate to severe	53	56/35	24/12	Not provided
RAHERISON [1]	French	430	Moderate to severe	57.4	29.4/16.8	19/4.6	Not provided

Data are presented as females/males %, unless otherwise stated.

18–74 years who were diagnosed with NSCLC, an increased risk of developing NSCLC before 55 years of age was seen in women with COPD (OR=1.67) compared to women without COPD. This effect was not present in the subgroup of African-American women. Emphysema was associated with an increased risk of NSCLC in women with COPD (OR=3.21). A history of chronic bronchitis also increased the risk of NSCLC (OR=1.7) [110]. The median time between COPD diagnosis and NSCLC diagnosis was 13 years regardless of age at diagnosis of NSCLC, but the average age of onset of COPD was 35 years for patients diagnosed with NSCLC before 55 years of age, compared to 51 years if the diagnosis of NSCLC was made after 55 years of age. The authors suggest a possible genetic susceptibility to early onset of lung disease and then NSCLC [110] in female smokers. In support of these findings, several pathophysiological hypotheses exist in the literature. DNA repair capacity is thought to be 10–15% lower in women than in men [111]. Women are said to have higher levels of DNA adducts and more DNA damage [111]. Increased metabolic bioactivation relating to tobacco use (hydrocarbons) was also a proposed hypothesis [111]. Women are thought to have increased expression of CYP (cytochrome P450 gene) enzymes in the distal airways that would be upregulated by oestrogen receptor- α [111]. Finally, oestrogens might influence NSCLC onset regardless of sex [112, 113] by oestrogen receptor- α and - β expression in lung adenocarcinoma cells [112].

Therapeutic care

Analysis of the effects of smoking cessation according to sex has been the subject of several studies involving very large samples. The “Lung Health Study” [49], which set out to assess lung function parameters in patients with mild-to-moderate COPD, found that FEV₁ gain was greater in women than in men at 1 year (3.7% versus 1.6%; $p < 0.001$) if smoking cessation was successful, and at 5 years, although the difference between the two sexes was less marked. Furthermore, the relationship between FEV₁ slope of decline and bronchial hyperreactivity was more significantly correlated in women than in men ($p < 0.001$). These findings confirmed those obtained in two previous Danish [114] and Canadian [50] cohorts, which showed that the greater the women’s past smoking habit, the greater the beneficial effects of smoking cessation [114]. Moreover, the female population appears to be more sensitive to the harmful effects of smoking [115]. However, it would appear that women find it harder than men to stop smoking and are more likely to relapse because of less clinical improvement after smoking cessation and potential weight gain [29, 49]. These factors justify the use of highly tailored cessation strategies designed specifically for women. A background of anxiety and depression complicates smoking cessation. Studies that have investigated the effect of pharmacological smoking cessation aids including bupropion and varenicline seem to find them equally effective in men and women [29, 116], although the efficacy of nicotine transdermal patches is more pronounced in the male population [117]. Women have stronger immune responses than men after a flu vaccination. They have more frequent adverse reactions than men, but the vaccination is more effective [118]. Given the small number of studies devoted to the influence of sex on the medical treatment of COPD, the published results might appear conflicting. Thus, an old study [119] on a small population suggested that, irrespective of age, the inhalation technique proved less effective in women than in men, with 4% of the female population satisfied with its use compared with 43% of males. This finding was not confirmed in a more recent study, which however did not control the extent to which patients were educated on the use of inhalers [120]. Similarly, the findings diverge in terms of compliance; women with mild-to-moderate COPD were reportedly more treatment compliant, whereas compliance was reported to be more satisfactory in men with severe COPD [121]. Few studies focus on the efficacy and tolerability of these treatments by sex. The study by VESTBO *et al.* [122] on salmeterol/fluticasone coadministration found no difference in therapeutic response between men and women in terms of FEV₁, frequency of exacerbations and quality of life. Another study of tiotropium showed identical effects on lung function, respiratory symptoms and quality of life in the female and male population [123]. A similar finding could be extrapolated from the results published in the TORCH study, in which treatment response was unaffected by sex, although the study was not designed to answer this question [124]. The lower proportion of women in the majority of studies makes it difficult to evaluate the pharmacology and effects of the different therapeutic classes used in COPD. We have little data on the differences in sex-based physiology and respiratory anatomy to make a potential effect on dose delivery and on the efficacy of inhaled therapies. However, the likelihood of a deterioration in clinical respiratory status (symptoms and exacerbations) on discontinuation of inhaled corticosteroids seems greater in female than male patients with COPD [125]. Finally, special attention should be paid to inhaled corticosteroid use in women given the the increased risk of osteoporosis, justifying supplementation with with vitamin D, calcium and bisphosphonates.

Osteoporosis is characterised by compromised bone strength and microarchitectural deterioration of bone tissue, leading to bone fragility and increased fracture risk. Its prevalence in COPD ranges from 24% to 69% [126]. The risk factors for COPD, smoking and menopause, are compounded in women with COPD. Oral corticosteroids are a recognised risk factor at a cumulative prednisone equivalent dose >7.5 mg daily for more than three consecutive months. The effects of long-term use of high-dose inhaled corticosteroids

TABLE 3 Osteoporosis risk factors

Major factors	Minor factors
Systemic corticosteroids for >3 months-year ⁻¹	Body mass index <21
Prior vertebral compression fracture	Active smoker
	Chronic alcoholism
	Aged >65 years
	Hip fracture
	Rib fracture
	Menopause
	Sedentariness
	FEV ₁ <50% of theoretical value

FEV₁: forced expiratory volume in 1 s.

have been reported previously [127, 128]. In post-menopausal women, the risk factors identified in addition to fracture risk were: body mass index <19, onset of menopause before the age of 40 years, and history of femoral neck fracture in a first-degree relative [129]. Pulmonary rehabilitation, which includes smoking cessation and exercise training, plays a key role in the treatment and prevention of osteoporosis; its beneficial effects on bone mineral density, strength muscle, balance and fall risk have been proven [107]. Calcium and vitamin D supplementation are effective in reducing fracture risk with a vitamin D dose-dependent effect only when combined with calcium [130]. A daily dose of 800 IU of vitamin D and 1 g of calcium is recommended with a T-score more than -1 with three minor risk criteria or one major risk criterion (table 3) [131, 132]. The value of antiresorptive therapy with bisphosphonates has been demonstrated in post-menopausal women in the steroid-induced forms of osteoporosis and in obstructive respiratory disease [133, 134]. Teriparatide appears beneficial in the treatment of glucocorticoid-induced osteoporosis but its value in COPD patients has yet to be established [133]. There are as yet no specific recommendations for COPD patients, let alone women with COPD [129, 135–137].

The majority of studies investigating the effects of oxygen therapy specifically for the treatment of COPD in men and women seem to have found a benefit in women. The study by MIYAMOTO *et al.* [138] in 1995 found that women on long-term oxygen therapy experienced significantly better survival, with a persistent difference after controlling for age, blood gas analysis and lung function. These results were confirmed by a Swedish cohort study of 5689 patients on oxygen therapy and followed up for >10 years, with a relative risk of excess mortality among men of 1.21 [139]. In other words, first year survival was 77% in women and 69% in men. More recent still, another cohort study in a larger population, whose mortality was assessed according to their comorbid conditions, confirmed the mortality reduction among women [32]. This latest study made it possible to establish that, although the significance of the comorbidities found in these patients is an independent predictor of mortality, female mortality was lower, even after adjusting for these comorbidities.

The explanation for this difference is not fully clear given that there was no difference in FEV₁ between the two sexes at the start of oxygen therapy. Treatment initiation does not appear earlier in women, especially since their arterial oxygen tension was lower than that of men when therapy was indicated. It is difficult to ascertain whether compliance with oxygen therapy differed given that it was rarely controlled. It is possible that the smoking cessation usually achieved through the introduction of oxygen therapy has been more beneficial for women, as described earlier in this chapter. Differences in environmental and occupational exposure, together with phenotypic differences accounting for the greater frequency of emphysema in men, could be implicated.

The improvement in exercise capacity and quality of life after pulmonary rehabilitation appears identical in men and women [140]. These programmes appear to be beneficial for women as they take into account many factors, including the role played by therapeutic education and its additional positive impact on anxiety and depression. However, FOY *et al.* [141] found that, despite a similar improvement in both sexes in terms of symptoms and quality of life after 3 months of exercise training, these benefits persisted only in men at the 18-month assessment. These results should be interpreted with caution in the absence of data on the exercises performed to maintain the gains achieved. Moreover, the expected benefits of pulmonary rehabilitation might differ between the sexes.

Conclusion

COPD in women is a major public health issue. It is biologically plausible that women are more vulnerable to nicotine addiction or environmental factors than men. More evidence is required, particularly to elucidate the role played by hormones.

In developing countries, COPD affects women differently to men, not only because of nicotine addiction, but also because of environmental exposure to biomass fuels.

The risk of death associated with active smoking among women increases with the number of cigarettes smoked and the age at which they become active smokers.

The perception of COPD has changed. It increasingly affects women, who differ in terms of both clinical presentation of symptoms and exacerbations and radiological presentation. Primary care professionals and pulmonologists must rise to the challenge of improving treatment through early diagnosis and shared knowledge of the specific features of COPD in women in primary care medicine.

To conclude, women remain under-represented in controlled clinical trials and few studies have investigated therapeutic response to COPD treatment according to sex.

Acknowledgements: We were assisted in the preparation of this article by Matrix Consultants (Paris, France).

Conflict of interest: C. Gut-Gobert reports personal fees from Novartis, Chiesi and GSK, outside the submitted work. A. Cavailles has nothing to disclose. A. Dixmier reports non-financial support from Novartis, outside the submitted work. S. Guillot has nothing to disclose. S. Jouneau reports grants and personal fees (fees, funding or reimbursement for national and international conferences, boards, expert or opinion groups and research projects) from Actelion, AIRB, AstraZeneca, BMS, Boehringer, Chiesi, Gilead, GSK, LVL, Mundipharma, Novartis, Pfizer, Roche and Savara-Serendex, outside the submitted work. C. Leroyer has nothing to disclose. S. Marchand-Adam reports personal fees from Novartis, Roche and Boehringer, outside the submitted work. D. Marquette has nothing to disclose. J-C. Meurice reports grants, personal fees and nonfinancial support from ResMed, grants from Philips, personal fees from Fisher Paykel, and personal fees and nonfinancial support from Novartis and Orkyn, outside the submitted work. N. Desvigne reports other fees from Novartis, outside the submitted work. H. Morel has nothing to disclose. C. Person-Tacnet has nothing to disclose. C. Raheison reports personal fees from Novartis, AstraZeneca, Chiesi, GSK and Boehringer Ingelheim, outside the submitted work.

Support statement: The study was supported by Novartis Pharma AG (Paris, France).

References

- 1 Raheison C, Tillie-Leblond I, Prudhomme A, *et al.* Clinical characteristics and quality of life in women with COPD: an observational study. *BMC Womens Health* 2014; 14: 31.
- 2 Pirie K, Peto R, Beral V. The first generation in which many women began smoking – authors' reply. *Lancet* 2013; 381: 1455–1456.
- 3 Thun MJ, Carter BD, Feskanich D, *et al.* 50-year trends in smoking-related mortality in the United States. *N Engl J Med* 2013; 368: 351–364.
- 4 Chapman KR, Tashkin DP, Pye DJ. Gender bias in the diagnosis of COPD. *Chest* 2001; 119: 1691–1695.
- 5 Fletcher C, Peto R. The natural history of chronic airflow obstruction. *Br Med J* 1977; 1: 1645–1648.
- 6 Kohansal R, Martinez-Cambor P, Agustí A, *et al.* The natural history of chronic airflow obstruction revisited: an analysis of the Framingham offspring cohort. *Am J Respir Crit Care Med* 2009; 180: 3–10.
- 7 Lopez AD, Collishaw NE, Piha T. A descriptive model of the cigarette epidemic in developed countries. *Tob Control* 1994; 3: 242–247.
- 8 Anderson SJ, Glantz SA, Ling PM. Emotions for sale: cigarette advertising and women's psychosocial needs. *Tob Control* 2005; 14: 127–135.
- 9 WHO. Les femmes et la santé: la réalité d'aujourd'hui, le programme de demain. Geneva, World Health Organization, 2009.
- 10 Hitchman SC, Fong GT. Gender empowerment and female-to-male smoking prevalence ratios. *Bull World Health Organ* 2011; 89: 195–202.
- 11 Hermalin AI, Lowry D. The age prevalence of smoking among Chinese women: a case of arrested diffusion? Population studies center research report 10-718. Michigan, University of Michigan Institute for Social Research, 2010.
- 12 WHO. Global Adult Tobacco Survey (GATS), Fact Sheet China: 2010. Geneva, World Health Organization, 2010.
- 13 Ancochea J, Miravittles M, García-Río F, *et al.* Underdiagnosis of chronic obstructive pulmonary disease in women: quantification of the problem, determinants and proposed actions. *Arch Bronconeumol* 2013; 49: 223–229.
- 14 Watson L, Vestbo J, Postma DS, *et al.* Gender differences in the management and experience of chronic obstructive pulmonary disease. *Respir Med* 2004; 98: 1207–1213.
- 15 Miravittles M, de la Roza C, Naberan K, *et al.* [Attitudes toward the diagnosis of chronic obstructive pulmonary disease in primary care]. *Arch Bronconeumol* 2006; 42: 3–8.
- 16 Martinez CH, Raparla S, Plauschinat CA, *et al.* Gender differences in symptoms and care delivery for chronic obstructive pulmonary disease. *J Womens Health (Larchmt)* 2012; 21: 1267–1274.
- 17 Tilert T, Dillon C, Paulose-Ram R, *et al.* Estimating the U.S. prevalence of chronic obstructive pulmonary disease using pre- and post-bronchodilator spirometry: the National Health and Nutrition Examination Survey (NHANES) 2007–2010. *Respir Res* 2013; 14: 103.
- 18 Soriano JB, Ancochea J, Miravittles M, *et al.* Recent trends in COPD prevalence in Spain: a repeated cross-sectional survey 1997–2007. *Eur Respir J* 2010; 36: 758–765.
- 19 Halbert RJ, Natoli JL, Gano A, *et al.* Global burden of COPD: systematic review and meta-analysis. *Eur Respir J* 2006; 28: 523–532.
- 20 Buist AS, McBurnie MA, Vollmer WM, *et al.* International variation in the prevalence of COPD (the BOLD Study): a population-based prevalence study. *Lancet* 2007; 370: 741–750.

- 21 Vos T, Flaxman AD, Naghavi M, *et al.* Years lived with disability (YLDs) for 1160 sequelae of 289 diseases and injuries 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2012; 380: 2163–2196.
- 22 Gershon AS, Wang C, Wilton AS, *et al.* Trends in chronic obstructive pulmonary disease prevalence, incidence, and mortality in Ontario, Canada, 1996 to 2007: a population-based study. *Arch Intern Med* 2010; 170: 560–565.
- 23 Bischoff EW, Schermer TR, Bor H, *et al.* Trends in COPD prevalence and exacerbation rates in Dutch primary care. *Br J Gen Pract* 2009; 59: 927–933.
- 24 Aryal S, Diaz-Guzman E, Mannino DM. Influence of sex on chronic obstructive pulmonary disease risk and treatment outcomes. *Int J Chron Obstruct Pulmon Dis* 2014; 9: 1145–1154.
- 25 Raheison C, Girodet PO. Epidemiology of COPD. *Eur Respir Rev* 2009; 18: 213–221.
- 26 Raheison C, Biron E, Nocent-Ejnaini C, *et al.* [Are there specific characteristics of COPD in women?]. *Rev Mal Respir* 2010; 27: 611–624.
- 27 Liu S, Zhou Y, Wang X, *et al.* Biomass fuels are the probable risk factor for chronic obstructive pulmonary disease in rural South China. *Thorax* 2007; 62: 889–897.
- 28 Spilka S, Le Nézet O, Ngantcha M, *et al.* Les drogues à 17 ans: analyses de l'enquête ESCAPAD 2014. www.ofdt.fr/publications/collections/periodiques/lettre-tendances/les-drogues-17-ans-analyse-de-lenquete-escapad-2014-tendances-100-mai-2015/ Date last updated: May, 2015.
- 29 Aryal S, Diaz-Guzman E, Mannino DM. COPD and gender differences: an update. *Transl Res* 2013; 162: 208–218.
- 30 Celli BR, Halbert RJ, Nordyke RJ, *et al.* Airway obstruction in never smokers: results from the Third National Health and Nutrition Examination Survey. *Am J Med* 2005; 118: 1364–1372.
- 31 Almagro P, López García F, Cabrera F, *et al.* Comorbidity and gender-related differences in patients hospitalized for COPD. The ECCO study. *Respir Med* 2010; 104: 253–259.
- 32 Ekstrom MP, Jørgensen C, Strom KE. Comorbidity and sex-related differences in mortality in oxygen-dependent chronic obstructive pulmonary disease. *PLoS One* 2012; 7: e35806.
- 33 Agusti A, Calverley PM, Celli B, *et al.* Characterisation of COPD heterogeneity in the ECLIPSE cohort. *Respir Res* 2010; 11: 122.
- 34 Di Marco F, Verga M, Reggente M, *et al.* Anxiety and depression in COPD patients: the roles of gender and disease severity. *Respir Med* 2006; 100: 1767–1774.
- 35 Gudmundsson G, Gislason T, Janson C, *et al.* Depression, anxiety and health status after hospitalisation for COPD: a multicentre study in the Nordic countries. *Respir Med* 2006; 100: 87–93.
- 36 Gudmundsson G, Gislason T, Janson C, *et al.* Risk factors for rehospitalisation in COPD: role of health status, anxiety and depression. *Eur Respir J* 2005; 26: 414–419.
- 37 Fuhrman C, Roche N, Vergnenegre A, *et al.* [Chronic bronchitis: prevalence and quality of life. Analysis of data from the French Health Interview Survey 2002-2003]. *Rev Mal Respir* 2009; 26: 759–768.
- 38 Roche N, Deslée G, Caillaud D, *et al.* Impact of gender on COPD expression in a real-life cohort. *Respir Res* 2014; 15: 20.
- 39 Hurdman J, Condliffe R, Elliot CA, *et al.* Pulmonary hypertension in COPD: results from the ASPIRE registry. *Eur Respir J* 2013; 41: 1292–1301.
- 40 Camp PG, Goring SM. Gender and the diagnosis, management, and surveillance of chronic obstructive pulmonary disease. *Proc Am Thorac Soc* 2007; 4: 686–691.
- 41 Ford ES, Mannino DM, Zhao G, *et al.* Changes in mortality among US adults with COPD in two national cohorts recruited from 1971–1975 and 1988–1994. *Chest* 2012; 141: 101–110.
- 42 Martinez FJ, Curtis JL, Sciruba F, *et al.* Sex differences in severe pulmonary emphysema. *Am J Respir Crit Care Med* 2007; 176: 243–252.
- 43 de Torres JP, Casanova C, Hernández C, *et al.* Gender and COPD in patients attending a pulmonary clinic. *Chest* 2005; 128: 2012–2016.
- 44 Cydulka RK, Rowe BH, Clark S, *et al.* Gender differences in emergency department patients with chronic obstructive pulmonary disease exacerbation. *Acad Emerg Med* 2005; 12: 1173–1179.
- 45 Grydeland TB, Dirksen A, Coxson HO, *et al.* Quantitative computed tomography: emphysema and airway wall thickness by sex, age and smoking. *Eur Respir J* 2009; 34: 858–865.
- 46 Sverzellati N, Calabrò E, Randi G, *et al.* Sex differences in emphysema phenotype in smokers without airflow obstruction. *Eur Respir J* 2009; 33: 1320–1328.
- 47 de Torres JP, Casanova C, Montejo de Garcini A, *et al.* Gender and respiratory factors associated with dyspnea in chronic obstructive pulmonary disease. *Respir Res* 2007; 8: 18.
- 48 Naberan K, Azpeitia A, Cantoni J, *et al.* Impairment of quality of life in women with chronic obstructive pulmonary disease. *Respir Med* 2012; 106: 367–373.
- 49 Connett JE, Murray RP, Buist AS, *et al.* Changes in smoking status affect women more than men: result of the Lung Health Study. *Am J Epidemiol* 2003; 157: 973–979.
- 50 Xu X, Li B, Wang L. Gender difference in smoking effects on adult pulmonary function. *Eur Respir J* 1994; 7: 477–483.
- 51 Chen Y, Horne SL, Dosman JA. Increased susceptibility to lung dysfunction in female smokers. *Am Rev Respir Dis* 1991; 143: 1224–1230.
- 52 Foreman MG, Zhang L, Murphy J, *et al.* Early-onset chronic obstructive pulmonary disease is associated with female sex, maternal factors, and African American race in the COPDGen Study. *Am J Respir Crit Care Med* 2011; 184: 414–420.
- 53 Silverman EK, Weiss ST, Drazen JM, *et al.* Gender-related differences in severe, early-onset chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2000; 162: 2152–2158.
- 54 Sheel AW, Guenette JA, Yuan R, *et al.* Evidence for dysanapsis using computed tomographic imaging of the airways in older ex-smokers. *J Appl Physiol* 2009; 107: 1622–1628.
- 55 Merkus PJ, Borsboom GJ, Van Pelt W, *et al.* Growth of airways and air spaces in teenagers is related to sex but not to symptoms. *J Appl Physiol* 1993; 75: 2045–2053.
- 56 Kim CS, Hu SC. Regional deposition of inhaled particles in human lungs: comparison between men and women. *J Appl Physiol* 1998; 84: 1834–1844.
- 57 Bennett WD, Zeman KL, Kim C. Variability of fine particle deposition in healthy adults: effect of age and gender. *Am J Respir Crit Care Med* 1996; 153: 1641–1647.

- 58 Sayiner A, Hague C, Ajlan A, *et al.* Bronchiolitis in young female smokers. *Respir Med* 2013; 107: 732–738.
- 59 Kohler M, Sandberg A, Kjellqvist S, *et al.* Gender differences in the bronchoalveolar lavage cell proteome of patients with chronic obstructive pulmonary disease. *J Allergy Clin Immunol* 2013; 131: 743–751.
- 60 Ali Assad N, Sood A. Leptin, adiponectin and pulmonary diseases. *Biochimie* 2012; 94: 2180–2189.
- 61 de Torres JP, Casanova C, Pinto-Plata V, *et al.* Gender differences in plasma biomarker levels in a cohort of COPD patients: a pilot study. *PLoS One* 2011; 6: e16021.
- 62 Leynaert B, Bousquet J, Henry C, *et al.* Is bronchial hyperresponsiveness more frequent in women than in men? A population-based study. *Am J Respir Crit Care Med* 1997; 156: 1413–1420.
- 63 Tashkin DP, Altose MD, Connett JE, *et al.* Methacholine reactivity predicts changes in lung function over time in smokers with early chronic obstructive pulmonary disease. The Lung Health Study Research Group. *Am J Respir Crit Care Med* 1996; 153: 1802–1811.
- 64 Taylor DR, Reid WD, Paré PD, *et al.* Cigarette smoke inhalation patterns and bronchial reactivity. *Thorax* 1998; 43: 65–70.
- 65 Massaro D, Massaro GD. Toward therapeutic pulmonary alveolar regeneration in humans. *Proc Am Thorac Soc* 2006; 3: 709–712.
- 66 Carlson CL, Cushman M, Enright PL, *et al.* Hormone replacement therapy is associated with higher FEV1 in elderly women. *Am J Respir Crit Care Med* 2001; 163: 423–428.
- 67 Key TJ, Pike MC, Brown JB, *et al.* Cigarette smoking and urinary oestrogen excretion in premenopausal and post-menopausal women. *Br J Cancer* 1996; 74: 1313–1316.
- 68 Tam A, Morrish D, Wadsworth S, *et al.* The role of female hormones on lung function in chronic lung diseases. *BMC Womens Health* 2011; 3: 11–24.
- 69 Choi HJ, Chung YS, Kim HJ, *et al.* Signal pathway of 17beta-estradiol-induced MUC5B expression in human airway epithelial cells. *Am J Respir Cell Mol Biol* 2009; 40: 168–178.
- 70 Salvi SS, Barnes PJ. Chronic obstructive pulmonary disease in non-smokers. *Lancet* 2009; 374: 733–743.
- 71 Hardie JA, Vollmer WM, Buist AS, *et al.* Respiratory symptoms and obstructive pulmonary disease in a population aged over 70 years. *Respir Med* 2005; 99: 186–195.
- 72 Lamprecht B, Schirnhofner L, Kaiser B, *et al.* Non-reversible airway obstruction in never smokers: results from the Austrian BOLD study. *Respir Med* 2008; 102: 1833–1838.
- 73 Lindström M, Kotaniemi J, Jönsson E, *et al.* Smoking, respiratory symptoms, and diseases: a comparative study between northern Sweden and northern Finland: report from the FinEsS study. *Chest* 2001; 119: 852–861.
- 74 Viegi G, Pedreschi M, Pistelli F, *et al.* Prevalence of airways obstruction in a general population: European Respiratory Society vs American Thoracic Society definition. *Chest* 2000; 117: 339S–345S.
- 75 Zhou Y, Wang C, Yao W, *et al.* COPD in Chinese nonsmokers. *Eur Respir J* 2009; 33: 509–518.
- 76 Ehrlich RI, White N, Norman R, *et al.* Predictors of chronic bronchitis in South African adults. *Int J Tuberc Lung Dis* 2004; 8: 369–376.
- 77 Kim DS, Kim YS, Jung KS, *et al.* Prevalence of chronic obstructive pulmonary disease in Korea: a population-based spirometry survey. *Am J Respir Crit Care Med* 2005; 172: 842–847.
- 78 Cerveri I, Accordini S, Verlato G, *et al.* Variations in the prevalence across countries of chronic bronchitis and smoking habits in young adults. *Eur Respir J* 2001; 18: 85–92.
- 79 von Hertzen L, Reunanen A, Impivaara O, *et al.* Airway obstruction in relation to symptoms in chronic respiratory disease – a nationally representative population study. *Respir Med* 2000; 94: 356–363.
- 80 Lamprecht B, McBurnie MA, Vollmer WM, *et al.* COPD in never smokers: results from the population-based burden of obstructive lung disease study. *Chest* 2011; 139: 752–763.
- 81 Smith M, Li L, Augustyn M, *et al.* Prevalence and correlates of airflow obstruction in ~317,000 never-smokers in China. *Eur Respir J* 2014; 44: 66–77.
- 82 de Torres JP, Casanova C, Hernández C, *et al.* Gender associated differences in determinants of quality of life in patients with COPD: a case series study. *Health Qual Life Outcomes* 2006; 4: 72.
- 83 Katsura H, Yamada K, Wakabayashi R, *et al.* Gender-associated differences in dyspnoea and health-related quality of life in patients with chronic obstructive pulmonary disease. *Respirology* 2007; 12: 427–432.
- 84 Kaptein AA, Scharloo M, Fischer MJ, *et al.* 50 years of psychological research on patients with COPD – road to ruin or highway to heaven? *Respir Med* 2009; 103: 3–11.
- 85 Rodríguez-González Moro JM, Izquierdo JL, Antón E, *et al.* Health-related quality of life in outpatient women with COPD in daily practice: the MUVICE Spanish study. *Respir Med* 2009; 103: 1303–1312.
- 86 Hrisanfow E, Häggglund D. Impact of cough and urinary incontinence on quality of life in women and men with chronic obstructive pulmonary disease. *J Clin Nurs* 2013; 22: 97–105.
- 87 Becklake MR, Kauffmann F. Gender differences in airway behaviour over the human life span. *Thorax* 1999; 54: 1119–1138.
- 88 Vermeeren MA, Creutzberg EC, Schols AM, *et al.* Prevalence of nutritional depletion in a large out-patient population of patients with COPD. *Respir Med* 2006; 100: 1349–1355.
- 89 Verhage TL, Heijdra Y, Molema J, *et al.* Associations of muscle depletion with health status. Another gender difference in COPD? *Clin Nutr* 2011; 30: 332–338.
- 90 Laurin C, Lavoie KL, Bacon SL, *et al.* Sex differences in the prevalence of psychiatric disorders and psychological distress in patients with COPD. *Chest* 2007; 132: 148–155.
- 91 Vanfleteren LE, Spruit MA, Groenen M, *et al.* Clusters of comorbidities based on validated objective measurements and systemic inflammation in patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2013; 187: 728–735.
- 92 Bos-Touwen I, Schuurmans M, Monninkhof EM, *et al.* Patient and disease characteristics associated with activation for self-management in patients with diabetes, chronic obstructive pulmonary disease, chronic heart failure and chronic renal disease: a cross-sectional survey study. *PLoS One* 2015; 10: e0126400.
- 93 McGarvey L, Lee AJ, Roberts J, *et al.* Characterisation of the frequent exacerbator phenotype in COPD patients in a large UK primary care population. *Respir Med* 2015; 109: 228–237.
- 94 Hanania NA, Müllerova H, Locantore NW, *et al.* Evaluation of COPD Longitudinally to Identify Predictive Surrogate Endpoints (ECLIPSE) study investigators. Determinants of depression in the ECLIPSE chronic obstructive pulmonary disease cohort. *Am J Respir Crit Care Med* 2011; 183: 604–611.

- 95 Jiménez-Ruiz CA, Andreas S, Lewis KE, *et al.* Statement on smoking cessation in COPD and other pulmonary diseases and in smokers with comorbidities who find it difficult to quit. *Eur Respir J* 2015; 46: 61–79.
- 96 Fan VS, Ramsey SD, Giardino ND, *et al.* Sex, depression, and risk of hospitalization and mortality in chronic obstructive pulmonary disease. *Arch Intern Med* 2007; 167: 2345–2353.
- 97 Schneider C, Jick SS, Bothner U, *et al.* COPD and the risk of depression. *Chest* 2010; 137: 341–347.
- 98 Scharf SM, Maimon N, Simon-Tuval T, *et al.* Sleep quality predicts quality of life in chronic obstructive pulmonary disease. *Int J Chron Obstruct Pulmon Dis* 2010; 6: 1–12.
- 99 Budhiraja R, Parthasarthy S, Habib MP, *et al.* Insomnia in patients with COPD. *Sleep* 2012; 35: 369–375.
- 100 Garrod R, Marshall J, Barley E, *et al.* Predictors of success and failure in pulmonary rehabilitation. *Eur Respir J* 2006; 27: 788–794.
- 101 Busch AM, Scott-Sheldon LAJ, Pierce J, *et al.* Depressed mood predicts pulmonary rehabilitation completion among women, but not men. *Respir Med* 2014; 108: 1007–1013.
- 102 Atlantis E, Fahey P, Cochrane B, *et al.* Bidirectional associations between clinically relevant depression or anxiety and COPD: a systematic review and meta-analysis. *Chest* 2013; 144: 766–777.
- 103 Laurin C, Moullec G, Bacon SL, *et al.* Impact of anxiety and depression on chronic obstructive pulmonary disease exacerbation risk. *Am J Respir Crit Care Med* 2012; 185: 918–923.
- 104 Divo M, Cote C, de Torres JP, *et al.* Comorbidities and risk of mortality in patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2012; 186: 155–161.
- 105 Loganathan RS, Stover DE, Shi W, *et al.* Prevalence of COPD in women compared to men around the time of diagnosis of primary lung cancer. *Chest* 2006; 129: 1305–1312.
- 106 Kiri VA, Soriano J, Visick G, *et al.* Recent trends in lung cancer and its association with COPD: an analysis using the UK GP Research Database. *Prim Care Respir J* 2010; 19: 57–61.
- 107 Cavallès A, Brinchault-Rabin G, Dixmier A, *et al.* Comorbidities of COPD. *Eur Respir Rev* 2013; 22: 454–475.
- 108 Mina N, Soubani AO, Cote ML, *et al.* The relationship between chronic obstructive pulmonary disease and lung cancer in African American patients. *Clin Lung Cancer* 2012; 13: 149–156.
- 109 Izquierdo JL, Resano P, El Hachem A, *et al.* Impact of COPD in patients with lung cancer and advanced disease treated with chemotherapy and/or tyrosine kinase inhibitors. *Int J Chron Obstruct Pulmon Dis* 2014; 9: 1053–1058.
- 110 Schwartz AG, Cote ML, Wenzlaff AS, *et al.* Chronic obstructive lung diseases and risk of non-small cell lung cancer in women. *J Thorac Oncol* 2009; 4: 291–299.
- 111 Ben-Zaken Cohen S, Paré PD, Man SF, *et al.* The growing burden of chronic obstructive pulmonary disease and lung cancer in women: examining sex differences in cigarette smoke metabolism. *Am J Respir Crit Care Med* 2007; 176: 113–120.
- 112 Marquez-Garban DC, Mah V, Alavi M, *et al.* Progesterone and estrogen receptor expression and activity in human non-small cell lung cancer. *Steroids* 2011; 76: 910–920.
- 113 Quoix E, Menecier B. [Lung cancer in women]. *Rev Mal Respir* 2005; 22: 55–62.
- 114 Lange P, Groth S, Nyboe GJ, *et al.* Effects of smoking and changes in smoking habits on the decline of FEV₁. *Eur Respir J* 1989; 2: 811–816.
- 115 Langhammer A, Johnsen R, Holmen J, *et al.* Cigarette smoking gives more respiratory symptoms among women than among men. The Nord-Trøndelag Health Study (HUNT). *J Epidemiol Community Health* 2000; 54: 917–922.
- 116 Tashkin DP, Rennard S, Hays JT, *et al.* Effects of varenicline on smoking cessation in patients with mild to moderate COPD: a randomized controlled trial. *Chest* 2011; 139: 591–599.
- 117 Han MK, Postma D, Mannino DM, *et al.* Gender and chronic obstructive pulmonary disease: why it matters. *Am J Respir Crit Care Med* 2007; 176: 1179–1184.
- 118 Klein SL, Pekosz A. Sex-based biology and the rational design of influenza vaccination strategies. *J Infect Dis* 2014; 209: Suppl 3, S114–S119.
- 119 Goodman DE, Israel E, Rosenberg M, *et al.* The influence of age, diagnosis, and gender on proper use of metered-dose inhalers. *Am J Respir Crit Care Med* 1994; 150: 1256–1261.
- 120 Hashmi A, Soomro JA, Menon A, *et al.* Incorrect inhaler technique compromising quality of life of asthmatic patients. *J Medicine* 2012; 13: 16–21.
- 121 Dales RE, Mehdizadeh A, Aaron SD, *et al.* Sex differences in the clinical presentation and management of airflow obstruction. *Eur Respir J* 2006; 28: 319–322.
- 122 Vestbo J, Soriano JB, Anderson JA, *et al.* Gender does not influence the response of the combination of salmeterol and fluticasone propionate in COPD. *Respir Med* 2004; 98: 1045–1050.
- 123 O'Donnell DE, Flüge T, Gerken F, *et al.* Effects of tiotropium on lung hyperinflation, dyspnoea and exercise tolerance in COPD. *Eur Respir J* 2004; 23: 832–840.
- 124 Calverley PM, Anderson JA, Celli B, *et al.* Salmeterol and fluticasone propionate and survival in chronic obstructive pulmonary disease. *N Engl J Med* 2007; 356: 775–789.
- 125 Schermer TR, Hendriks AJ, Chavannes NH, *et al.* Probability and determinants of relapse after discontinuation of inhaled corticosteroids in patients with COPD treated in general practice. *Prim Care Respir J* 2004; 13: 48–55.
- 126 Graat-Verboom L, Spruit MA, van den Borne BE, *et al.* Correlates of osteoporosis in chronic obstructive pulmonary disease: an underestimated systemic component. *Respir Med* 2009; 103: 1143–1151.
- 127 Drummond MB, Dasenbrook EC, Pitz MW, *et al.* Inhaled corticosteroids in patients with stable chronic obstructive pulmonary disease: a systematic review and meta-analysis. *JAMA* 2008; 300: 2407–2416.
- 128 Richey F, Bousquet J, Ehrlich GE, *et al.* Inhaled corticosteroids effects on bone in asthmatic and COPD patients: a quantitative systematic review. *Osteoporos Int* 2003; 14: 179–190.
- 129 Haute Autorité de Santé. Les médicaments de l'ostéoporose. www.has-sante.fr/portail/jcms/c_1751307/fr/les-medicaments-de-l-osteoporose Date last updated: June, 2014.
- 130 Avenell A, Gillespie WJ, Gillespie LD, *et al.* Vitamin D and vitamin D analogues for preventing fractures associated with involutional and post-menopausal osteoporosis. *Cochrane Database Syst Rev* 2009; 2: CD000227.
- 131 Langhammer A, Forsmo S, Syversen U. Long-term therapy in COPD: any evidence of adverse effect on bone? *Int J Chron Obstruct Pulmon Dis* 2009; 4: 365–380.
- 132 Qaseem A, Snow V, Shekelle P, *et al.* Screening for osteoporosis in men: a clinical practice guideline from the American College of Physicians. *Ann Intern Med* 2008; 148: 680–684.

- 133 Grossman JM, Gordon R, Ranganath VK, *et al.* American College of Rheumatology 2010 recommendations for the prevention and treatment of glucocorticoid-induced osteoporosis. *Arthritis Care Res (Hoboken)* 2010; 62: 1515–1526.
- 134 Smith BJ, Laslett LL, Pile KD, *et al.* Randomized controlled trial of alendronate in airways disease and low bone mineral density. *Chron Respir Dis* 2004; 1: 131–137.
- 135 Briot K, Cortet B, Thomas T, *et al.* 2012 update of French guidelines for the pharmacological treatment of postmenopausal osteoporosis. *Joint Bone Spine* 2012; 79: 304–313.
- 136 Romme EA, Geusens P, Lems WF, *et al.* Fracture prevention in COPD patients; a clinical 5-step approach. *Respir Res* 2015; 16: 32.
- 137 Compston J, Bowring C, Cooper A, *et al.* Diagnosis and management of osteoporosis in postmenopausal women and older men in the UK: National Osteoporosis Guideline Group (NOGG) update 2013. *Maturitas* 2013; 75: 392–396.
- 138 Miyamoto K, Aida A, Nishimura M, *et al.* Gender effect on prognosis of patients receiving long-term home oxygen therapy. The Respiratory Failure Research Group in Japan. *Am J Respir Crit Care Med* 1995; 152: 972–976.
- 139 Franklin KA, Gustafson T, Ranstam J, *et al.* Survival and future need of long-term oxygen therapy for chronic obstructive pulmonary disease – gender differences. *Respir Med* 2007; 101: 1506–1511.
- 140 de Torres JP, Casanova C, Cote CG, *et al.* Women with chronic obstructive pulmonary disease: an emerging phenotype of the disease. *Therapy* 2009; 6: 821–830.
- 141 Foy CG, Rejeski WJ, Berry MJ, *et al.* Gender moderates the effects of exercise therapy on health-related quality of life among COPD patients. *Chest* 2001; 119: 70–76.