



RATIONALITY OF THE LOCUS OF SINGULARITIES OF THE GENERAL GOUGH-STEWART PLATFORM

Michel Coste, Seydou Moussa

► **To cite this version:**

Michel Coste, Seydou Moussa. RATIONALITY OF THE LOCUS OF SINGULARITIES OF THE GENERAL GOUGH-STEWART PLATFORM. 2019. hal-02083002

HAL Id: hal-02083002

<https://hal-univ-rennes1.archives-ouvertes.fr/hal-02083002>

Submitted on 2 Apr 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Rationality of the locus of singularities of the general Gough-Stewart platform

Michel Coste*and Seydou Moussa†

April 2, 2019

Abstract

We prove that the set of singular configurations of a general Gough Stewart platform has a rational parametrization. We introduce a reciprocal twist mapping which, for a general orientation of the platform, realizes the cubic surface of singularities as the blowing up of a quadric surface in five points.


Keywords: parallel robot, singularities, cubic surface

AMS classification: 70B15, 14E08, 14J26

Introduction

Gough-Stewart platform is the most well-known parallel robot with six degrees of freedom. It consists in a platform linked to a fixed base via six limbs whose lengths are controlled by actuated prismatic joints ; each limb is attached to the base by a universal joint and to the platform by a spherical joint (6-UPS in the standard notation [8]).

Figure 1: A Gough-Stewart platform



The problem of avoidance of singular configurations (in which one loses control on at least one degree of freedom) is crucial for parallel

*Univ Rennes, CNRS, IRMAR - UMR 6625, F-35000 Rennes, France

†Univ. Dan Dicko Dankoulodo, Dép. Mathématiques, Maradi, Niger

robots, and many papers have been devoted to the study of the set Sing of singular configurations of the Gough-Stewart platform. A basic fact is the characterization of singular configurations as those for which the systems of Plücker coordinates of the six limbs of the platform are linearly dependant (see [8]). This leads to a description of Sing by a polynomial equation in terms of position variables and orientation variables. This equation is of degree 3 in the position variables for a general architecture (this degree may drop for special architectures, see [7, 10]). Hence, for a generic (in the sense of algebraic geometry) orientation, the singular configurations form a cubic surface Σ in 3-dimensional space.

A way to describe the set of singular configurations of a robot is to give a rational parametrization of this set, when this is possible; this has been done for 3-RPR planar robots in [6], which stresses the advantage of rationality. In the case of the Gough-Stewart platform, a rational parametrization has been found for special architecture in [1], and the rationality of Sing has been proved for the general case of planar base and platform base in [3]. The rationality in this case follows from the existence of a singular point at infinity of the cubic surface Σ . This argument cannot be used for the general Gough-Stewart platform : the projective closure Σ^h of Σ is in general a nonsingular cubic surface. It is known (see [11]) that the rationality of a nonsingular cubic projective surface is related to special features of the set of the 27 lines on this surface. Experiments using exact computations over rational numbers show that the polynomial of degree 27 whose roots corresponds to the lines splits in factors of degrees 2, 5, 10, 10, where the lines corresponding to the factor of degree 2 are skew. This gives an evidence for rationality, which has to be confirmed by a proof. The key to the proof is given by a consideration from kinematics: to each singular configuration of the platform one can associate a line of reciprocal twists, which expresses the infinitesimal rigid motion which can no longer be controlled in the singular configuration. We prove that, for a general architecture, this reciprocal twist mapping is a regular mapping from the projective cubic surface Σ^h to a quadric surface Q (in the 5-dimensional projective space of twists). We prove moreover that the reciprocal twist mapping is the blowing-up of the quadric surface Q in five points, and the five exceptional divisor in the cubic surface Σ^h are the lines corresponding to the factor of degree 5 mentioned above. The singularities which form these five exceptional divisors are those for which all six limbs of the platform have a common secant line, and the reciprocal twist is a twist of rotation about this secant line.

We have proved in this way the rationality of the locus of singularities of the general Gough-Stewart platform. But, more than that, the information obtained about the reciprocal twist mapping may be of interest for the kinematics of this robot.

The paper is organized as follows. In Section 1 we describe the singularity locus and explain how the problem of its rational parametrization reduces to the problem of the rationality of the cubic surface Σ defined over the field of functions of the rotation group $\text{SO}(3)$; we recall the relation between the rationality of this surface and the structure of the set of lines on this surface, and give the example of a computation about these lines which shows an evidence for rationality. In section 2 we introduce the reciprocal twist mapping and show that it gives a birational equivalence between Σ and a quadric surface Q , which proves the rationality of Σ . In section 3 we extend the reciprocal twist mapping to the projective closure Σ^h of Σ and show that the extended mapping is the blowing-up of

Q in five points; we make precise the kinematic significance of the exceptional divisors of this blowing-up. We conclude by collecting some open questions raised by our approach.

1 The singularity locus of a Gough-Stewart platform, and the question of its rational parametrization

We consider a Gough-Stewart platform with an arbitrary architecture. For $i = 1, \dots, 6$, we denote by A_i the vector of coordinates of the center of the universal joint of the i -th limb on the base, in the fixed orthonormal frame, and by b_i the vector of coordinates of the center of the spherical joint on the mobile platform in the orthonormal frame attached to this platform. We may assume $A_1 = b_1 = (0, 0, 0)^T$ and use it whenever it is convenient.

1.1 Equation of the singularity locus in $SE(3)$

We recall the derivation of the equation of the singularity hypersurface, mainly in order to fix notation. The computation is very similar to the one in [4].

The group $SE(3)$ of rigid motions in 3-space acts on the mobile platform by the transformation $X \mapsto RX + P$ where R is the rotation matrix and $P = (x, y, z)^T$ is the translation vector. So the coordinates of the joints on the mobile platform are, in the fixed frame, $Rb_i + P$ for $i = 1 \dots 6$; we set $B_i = Rb_i$ and $C_i = B_i - A_i$.

The Plücker coordinates of the limbs w.r.t. the fixed frame are 6-dimensional vectors whose first three coordinates are $C_i + P$ and last three coordinates (the moment w.r.t. the origin) $A_i \times (C_i + P)$. It is well known and explained in [8] that the Gough-Stewart platform is in a singular configuration if and only if the Plücker coordinates of the six limbs are linearly dependant. This is expressed by the vanishing of the determinant of the 6×6 matrix whose rows are the Plücker coordinates of the limbs:

$$\text{Jac} = \begin{pmatrix} C_1 + P \\ \vdots \\ C_6 + P \\ A_1 \times (C_1 + P) \\ \vdots \\ A_6 \times (C_6 + P) \end{pmatrix}_{i=1, \dots, 6}^T. \quad (1)$$

We denote by $\text{Sing} \subset SE(3)$ the hypersurface of singular configurations, whose equation is $\det(\text{Jac}) = 0$. The following result is well known.

Theorem 1 *For a generic Gough-Stewart platform, the singularity locus Sing in $SE(3)$ has an equation which is of degree 3 with respect to P .*

Proof: We compute $\det(\text{Jac})$ using the generalized Laplace expansion w.r.t. the first three rows of Jac . We use the notation $[U, V, W]$ to denote the mixed product of the three 3-dimensional vectors U, V, W (i.e. the determinant whose columns are U, V, W).

$$\det(\text{Jac}) = \sum_{1 \leq i_1 < i_2 < i_3 \leq 6} (-1)^{i_1+i_2+i_3} [C_{i_1} + P, C_{i_2} + P, C_{i_3} + P] [A_{j_1} \times (C_{j_1} + P), A_{j_2} \times (C_{j_2} + P), A_{j_3} \times (C_{j_3} + P)], \quad (2)$$

where $j_1 < j_2 < j_3$ are the integers between 1 and 6 different from i_1, i_2, i_3 . The mixed products $[C_{i_1} + P, C_{i_2} + P, C_{i_3} + P]$ have degree at most 1 w.r.t.

P , while the mixed products $[A_{j_1} \times (C_{j_1} + P), A_{j_2} \times (C_{j_2} + P), A_{j_3} \times (C_{j_3} + P)]$ have degree at most 2, because the vectors $A_{j_1} \times P, A_{j_2} \times P, A_{j_3} \times P$ are linearly dependant. So the degree of $\det(\text{Jac})$ w.r.t. P is at most 3. The computation in a example (see later) shows that it is actually 3 for a generic Gough-Stewart platform. \square

1.2 Cayley parametrization for rotation matrices

Let

$$U = \begin{pmatrix} 0 & -r & q \\ r & 0 & -p \\ -q & p & 0 \end{pmatrix}$$

be a skew-symmetric matrix. Since 1 is not an eigenvalue of U , $I - U$ is invertible. The matrix $R = (I + U)(I - U)^{-1}$ is a rotation matrix

$$R(p, q, r) = \frac{1}{\Delta} \begin{pmatrix} 1 + p^2 - q^2 - r^2 & 2(pq - r) & 2(pr + q) \\ 2(pq + r) & 1 - p^2 + q^2 - r^2 & 2(qr - p) \\ 2(pr - q) & 2(qr + p) & 1 - p^2 - q^2 + r^2 \end{pmatrix}, \quad (3)$$

where $\Delta = 1 + p^2 + q^2 + r^2$. If the vector $(p, q, r)^\top$ is not the zero vector, it spans the axis of the rotation R . The tangent of the half-angle of the rotation is $\sqrt{\Delta - 1}$. The Cayley parametrization is a rational parametrization of all rotation matrices, except the half-turns. These half-turns are obtained as limits as $p^2 + q^2 + r^2$ tends to infinity; alternatively, one can use the homogeneous Euler-Rodrigues parametrization (that is, parametrization with quaternions) with one more variable.

Note that we can recover rationally p, q, r from the rotation matrix $R(p, q, r)$. Indeed

$$p = \frac{R_{3,2} - R_{2,3}}{1 + \text{tr}(R)}, \quad q = \frac{R_{1,3} - R_{3,1}}{1 + \text{tr}(R)}, \quad r = \frac{R_{2,1} - R_{1,2}}{1 + \text{tr}(R)}, \quad (4)$$

where $\text{tr}(R) = R_{1,1} + R_{2,2} + R_{3,3}$. This shows:

Proposition 2 *The Cayley parametrization induces an isomorphism between the field $\mathbb{R}(\text{SO}(3))$ of rational functions on $\text{SO}(3)$ and the field of rational functions in three independent variables $\mathbb{R}(p, q, r)$.*

The preceding result says, in terms of algebraic geometry, that the variety $\text{SO}(3)$ is a rational variety over \mathbb{R} .

1.3 The cubic surface Σ

The equation $\det(\text{Jac}) = 0$ defines the algebraic variety of singular configurations $\text{Sing} \subset \text{SE}(3)$. For a fixed $R \in \text{SO}(3)$, the equation $\det(\text{Jac}) = 0$ in the variables x, y, z defines a cubic surface $\text{Sing}_R \subset \mathbb{R}^3$, and Sing may be viewed as the family of these cubic surfaces parametrized by $\text{SO}(3)$. We can also view the equation $\det(\text{Jac}) = 0$ as an equation in the three variables x, y, z with coefficients in the field $\mathbb{R}(\text{SO}(3))$ (or the field $\mathbb{R}(p, q, r)$, according to Proposition 2). As such, this is the equation of a cubic surface Σ defined over $\mathbb{R}(\text{SO}(3))$. This surface Σ will be the main object of study in the rest of this article. In the language of algebraic geometry, Σ is the generic fibre of the family of surfaces Sing_R parametrized by $\text{SO}(3)$.

In order to prove that the algebraic variety Sing is rational over \mathbb{R} , it suffices to prove that Σ is rational over $\mathbb{R}(\text{SO}(3))$, since $\text{SO}(3)$ itself is rational over \mathbb{R} , as shown in Proposition 2. This will be the aim of the

next section. So we shall work over the field $\mathbb{R}(\text{SO}(3))$. We shall feel free to use the classical formulas of vector algebra relating dot product, cross product, mixed product over $\mathbb{R}(\text{SO}(3))$ as if we were in \mathbb{R}^3 .

A general criterion for rationality of a cubic surface over a field k is due to Swinnerton-Dyer. It is related to the geometry of lines on the cubic surface. A very classical result of algebraic geometry says that a smooth projective cubic surface over a field k has 27 lines, which are defined over the algebraic closure of k (see for instance [5], Theorem 4.9 p.402).

Theorem 3 ([11]) *Let Γ be a smooth projective cubic surface over a field k (of characteristic 0). Call S_n a set of n of the 27 lines on Γ , mutually skew, which is stable under conjugation over k ; i.e., the union of these lines is defined over k . Then Γ is k -rational iff it has a k -rational point and a S_2 , or a S_3 , or a S_6 .*

We shall first investigate experimentally whether the projective closure Σ^h of the cubic surface Σ may satisfy this criterion. The existence of a $\mathbb{R}(\text{SO}(3))$ -rational point on Σ is certain, since $P = (0, 0, 0)^\top$ is clearly always on Σ (when we assume $A_1 = b_1 = (0, 0, 0)^\top$). So the problem is only the splitting of the set of 27 lines over $\mathbb{R}(\text{SO}(3))$.

In principle, we could do the computation leaving the geometry of the Gough-Stewart platform free and the orientation R free. This would be a too heavy computation. Instead, we fix the geometry of the platform by choosing rational coordinates and pick a rotation matrix R also with rational coefficients, so that all the computations will be exact computations over \mathbb{Q} , concerning the surface Sing_R defined over \mathbb{Q} . If the set of lines on Σ splits over $\mathbb{R}(\text{SO}(3))$ then the set of lines on Sing_R will split accordingly over \mathbb{Q} . Experiments show the presence of a S_2 defined over \mathbb{Q} on Sing_R , with the same pattern of splitting for different choices of R and different geometries of the platform. Note that for a general cubic surface defined over \mathbb{Q} , the set of 27 lines does not split over \mathbb{Q} .

1.4 A case study

We fix the geometry of the Gough-Stewart platform as follows:

$$A_1 = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} A_2 = \begin{pmatrix} 2 \\ 0 \\ 0 \end{pmatrix} A_3 = \begin{pmatrix} 0 \\ 2 \\ -1 \end{pmatrix} A_4 = \begin{pmatrix} 0 \\ 1 \\ 2 \end{pmatrix} A_5 = \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} A_6 = \begin{pmatrix} 6 \\ 3 \\ 0 \end{pmatrix}$$

$$b_1 = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} b_2 = \begin{pmatrix} 2 \\ 3 \\ -1 \end{pmatrix} b_3 = \begin{pmatrix} 0 \\ 1 \\ 4 \end{pmatrix} b_4 = \begin{pmatrix} 1 \\ 3 \\ 1 \end{pmatrix} b_5 = \begin{pmatrix} 1 \\ 3 \\ -1 \end{pmatrix} b_6 = \begin{pmatrix} 2 \\ 4 \\ -3 \end{pmatrix}$$

We also choose Cayley parameters $p = 0$, $q = 0$, $r = 0$ for the rotation matrix $R(p, q, r)$, so that R is the identity matrix and $B_i = b_i$ for $i = 1, \dots, 6$

The set of singular configurations for the chosen rotation R is the cubic surface Sing_R with equation

$$\begin{aligned} &80x^3 - 107yx^2 - 47zx^2 - 376x^2 - 9y^2x - 301yx + 95zxy - 1392x \\ &- 96z^2x + 643zx - 68y^3 + 98zy^2 - 78y^2 + 426y + 708zy \\ &+ 78z^2y + 234z^2 - 24z^3 + 1410z \end{aligned} \quad (5)$$

In order to check that the projective closure $(\text{Sing}_R)^h$ of this cubic surface is nonsingular, we homogenize the cubic equation with homogeneization

variable w and verify that the ideal generated by the partial derivatives of the homogenized equation w.r.t. w, x, y, z contains the fifth power of the ideal generated by w, x, y, z .

There are 27 lines on this cubic surface; it happens that they are all real for the present example, but this is not the case for other choices. These lines can be computed in the following way: we write a parametrization of a line with parameter t and indeterminate coefficients a, b, c, d

$$x = t, \quad y = a + bt, \quad z = c + dt.$$

Substituting this parametrization in the equation of the cubic surface yields a degree 3 polynomial in t whose coefficients depend on a, b, c, d . The ideal generated by these four coefficients is the ideal of lines on the surface (assuming that no line is contained in the plane at infinity nor parallel to the (y, z) -plane). We check that the quotient of $\mathbb{Q}[a, b, c, d]$ by this ideal is indeed a finite extension of degree 27. The Groebner basis of the ideal with respect to the lexicographic order on a, b, c, d contains a polynomial of degree 27 in d , and a, b, c are rational polynomials in d in the quotient. This polynomial is the product of four irreducible factors over \mathbb{Q} , of degrees respectively 2, 5, 10 and 10. The factors of degree 2 and 5 are

$$\begin{aligned} F_2 &= 2796d^2 + 4137d - 56 \\ F_5 &= 14853594d^5 + 160133255d^4 - 6870509d^3 \\ &\quad - 1145865348d^2 + 1491086416d - 515006656 \end{aligned} \tag{6}$$

We explore the coplanarity relations between the lines corresponding to the different factors. In order to do this, we make computations over the quadratic extension of \mathbb{Q} given by F_2 and we use the fact that the lines with coefficients a, b, c, d and a', b', c', d' are coplanar if and only if

$$(a - a')(d - d') - (b - b')(c - c') = 0.$$


It can be checked that

- the two lines corresponding to the factor F_2 are skew,
- the five lines corresponding to the factor F_5 are those among the 27 which meet both lines of F_2 ,
- the ten lines corresponding to one of the factor of degree 10 are those which meet exactly one of the lines of F_2 (five each),
- the ten lines corresponding to the remaining factor are those which meet none of the lines of F_2 .

The same factorization pattern appears for generic choices of architecture and orientation (with rational parameters). The two lines corresponding to factor F_2 form a S_2 over \mathbb{Q} . This is a strong indication that Σ itself has a S_2 over $\mathbb{R}(\text{SO}(3))$, and hence is rational over this field. We shall prove this fact in the following section.

It is well-known how to obtain a rational parametrization from a S_2 . Pick a point on the cubic surface; there is a unique line through this point meeting the two lines of the S_2 , intersection of the planes containing the point and a line of the S_2 . The set of lines meeting the two lines of the S_2 can be seen as a quadric surface, and we obtain in this way a regular mapping from the cubic surface to a quadric surface; this regular mapping is a birational isomorphism, actually the blowing-up of the quadric surface

Figure 2: The cubic surface Sing_R , the two lines corresponding to F_2 in black, the five lines corresponding to F_5 in red



in the five points corresponding to the five lines contained in the cubic surface and meeting the two lines of the S_2 . For more detail on this, see for instance [9]. We shall in the next section show how kinematics provides a regular birational morphism from Σ to a quadric surface, and we shall prove in Section 2 That the projective closure Σ^h is indeed the blowing up of this quadric surface in five points.

2 The reciprocal twist at a singular configuration

2.1 Solutions of homogeneous systems of corank 1

Recall that a twist $\begin{pmatrix} \Omega \\ V \end{pmatrix}$ is said to be reciprocal to a screw $\begin{pmatrix} F \\ M \end{pmatrix}$ when their reciprocal product $F \cdot V + \Omega \cdot M$ is zero. If a Gough Stewart platform is in a singular configuration such that the rank of the family of systems of Plücker coordinates $\begin{pmatrix} F_i(P) \\ M_i(P) \end{pmatrix}$ of the six limbs is equal to five, then there is a line of twists reciprocal to all $\begin{pmatrix} F_i(P) \\ M_i(P) \end{pmatrix}$.

Let $A = (a_{i,j})_{i,j=1,\dots,n}$ be a square matrix of size n whose entries are polynomials in $X = (x_1, \dots, x_d)$ with coefficients in the field k .

Lemma 4 *If $\det(A) = 0$ is a nonsingular hypersurface in d -dimensional affine space k^d , then the matrix A has rank $n - 1$ at every point of this hypersurface. Moreover, if $\text{sol}(A)$ denotes the line of solutions S of $AS = 0$ for every A such that $\det(A) = 0$, then $A \mapsto \text{sol}(A)$ is a regular mapping from the hypersurface $\det(A) = 0$ to the projective space $\mathbb{P}^{n-1}(k)$.*

Proof. The derivative of $\det(A)$ with respect to x_ℓ can be expressed as

$$\frac{\partial \det(A)}{\partial x_\ell} = \sum_{j=1}^n \sum_{i=1}^n \text{cof}(A)_{i,j} \frac{\partial a_{i,j}}{\partial x_\ell}, \quad (7)$$

where $\text{cof}(A)_{i,j}$ is the cofactor of $a_{i,j}$ in matrix A . Assume that there is a point in the affine space of dimension d where the rank of A is $< n - 1$. Then all minors of dimension $n - 1$ of the matrix A vanish at this point and the formula above shows that all partial derivatives of $\det(A)$ also vanish at this point. This contradicts the assumption that $\det(A) = 0$ is a nonsingular hypersurface.

Let U_i be the Zariski open subset of the nonsingular hypersurface $\det(A) = 0$ where all rows of A but the i -th are linearly independent. By the first part of the lemma, $(U_i)_{i=1,\dots,n}$ is a Zariski open cover of $\det(A) = 0$. For any A in U_i , the line of solutions of $AS = 0$ is spanned by the vector with coordinates $\text{cof}(A)_{i,j}$ for $j = 1, \dots, n$; these cofactors are polynomial in the coefficients of A , hence also polynomials in X . In the end, we obtain a regular mapping from the hypersurface $\det(A) = 0$ in k^d to $\mathbb{P}^{n-1}(k)$. \square

The aim of this paper is to prove the rationality of Sing for a general Gough-Stewart platform, without assuming planarity of the base or of the platform.

Proposition 5 *Assume that Σ is a nonsingular hypersurface of $\mathbb{R}(\text{SO}(3))^3$. Then, for every singular configuration in Σ , the system of Plücker coordinates of the six limbs is of rank 5 in the space of screws. The mapping which associates, to each singular configuration, the line of twists reciprocal to the Plücker coordinates of the six limbs is a regular mapping from Σ to $\mathbb{P}^5(\mathbb{R}(\text{SO}(3)))$, which we denote by Rec.*

Proof. Lemma 4 shows that the rank of the matrix Jac of the Plücker coordinates of the six limbs is indeed 5 for each singular configuration P . The matrix Jac is the matrix of the system expressing the reciprocity of the twist $\begin{pmatrix} \Omega \\ V \end{pmatrix}$ to the systems of Plücker coordinates $\begin{pmatrix} F_i(P) \\ M_i(P) \end{pmatrix}$ of the six limbs. The second part of Lemma 4 shows that the mapping Rec associating to each singular configuration P the line of screws reciprocal to the Plücker coordinates of the six limbs is indeed a regular mapping from Σ to $\mathbb{P}^5\mathbb{R}(\text{SO}(3))$. \square

We are going to show in the following that the image of Rec is contained in a quadric surface in a 3-dimensional subspace of $\mathbb{P}^5(\mathbb{R}(\text{SO}(3)))$ and that Rec is a birational equivalence with this quadric surface.

2.2 The image of Rec is contained in a quadric surface

The equations expressing that the twist $\begin{pmatrix} \Omega \\ V \end{pmatrix}$ is reciprocal to the systems of Plücker coordinates of the limbs are

$$(C_i + P) \cdot V + (A_i \times (C_i + P)) \cdot \Omega = 0, \quad i = 1, \dots, 6. \quad (8)$$

This is equivalent to:

$$V \cdot C_i + [\Omega, A_i, C_i] + (\Omega \times A_i + V) \cdot P = 0, \quad i = 1, \dots, 6, \quad (9)$$

where $[\Omega, A_i, C_i]$ is the mixed product of vectors Ω, A_i, C_i .

We are going to eliminate P from this system of equations. The first step is to obtain two linear equations in V, Ω where P does not appear. In equation $(9)_1$, $A_1 = C_1 = 0$ so $(9)_1$ is $V \cdot P = 0$. Subtracting this first equation from the other ones we get:

$$V \cdot C_i + [\Omega, A_i, C_i] + (\Omega \times A_i) \cdot P = 0, \quad i = 2, \dots, 6. \quad (10)$$

The family of vectors $A_i, i = 2, \dots, 6$ has rank at most 3. Up to reordering the indices, we may assume that A_5 and A_6 are linear combinations of A_2, A_3 and A_4 . Set $A_5 = \alpha_2 A_2 + \alpha_3 A_3 + \alpha_4 A_4$ and $A_6 = \beta_2 A_2 + \beta_3 A_3 + \beta_4 A_4$. Computing $(10)_5 - \sum_{i=2}^4 \alpha_i (10)_i$ and $(10)_6 - \sum_{i=2}^4 \beta_i (10)_i$ we obtain respectively

$$V \cdot (C_5 - \sum_{i=2}^4 \alpha_i C_i) + [\Omega, A_5, C_5] - \sum_{i=2}^4 \alpha_i [\Omega, A_i, C_i] = 0 \quad (11)$$

and

$$V \cdot (C_6 - \sum_{i=2}^4 \beta_i C_i) + [\Omega, A_6, C_6] - \sum_{i=2}^4 \beta_i [\Omega, A_i, C_i] = 0 \quad (12)$$

which are linear homogeneous equations in Ω, V .

The second step in the elimination of P is to obtain a quadratic equation in V, Ω without P . Set $\ell_2 = [\Omega, A_3, A_4]$, $\ell_3 = [\Omega, A_4, A_2]$, $\ell_4 = [\Omega, A_2, A_3]$.

Lemma 6 $\ell_2(\Omega \times A_2) + \ell_3(\Omega \times A_3) + \ell_4(\Omega \times A_4) = 0$.

Proof. It suffices to check this formula of vector algebra assuming A_2, A_3, A_4 linearly independent. The dot product of the left handside with any of A_2, A_3, A_4 is easily seen to be zero, hence, it is the zero vector. \square

The linear combination of $(10)_2, (10)_3, (10)_4$ with coefficients ℓ_2, ℓ_3, ℓ_4 yields the following homogeneous quadratic equation in Ω, V :

$$\begin{aligned} & [\Omega, A_3, A_4](V \cdot C_2 + [\Omega, A_2, C_2]) + [\Omega, A_4, A_2](V \cdot C_3 + [\Omega, A_3, C_3]) \\ & + [\Omega, A_2, A_3](V \cdot C_4 + [\Omega, A_4, C_4]) = 0. \end{aligned} \quad (13)$$

Equations (11), (12) and (13) are the homogeneous equations of a quadric surface Q in $\mathbb{P}^5(\mathbb{R}(\text{SO}(3)))$, the projectivisation of the space of twists. We shall check in the case study that this projective quadric surface is nonsingular for a generic Gough-Stewart platform. We have proved

Proposition 7 *The image of the regular mapping Rec is contained in the quadric surface Q given by equations (11), (12) and (13).*

2.3 The birational equivalence

We obtained the system of equations (11), (12) and (13) from the system of equations in P, V and Ω expressing that the screw $\begin{pmatrix} \Omega \\ V \end{pmatrix}$ is reciprocal to the Plücker coordinates of the six limbs for the singular configuration P . Let Ω, V be the homogeneous coordinates of a point on the quadric surface Q such that $[V, \Omega \times A_2, \Omega \times A_3] \neq 0$. Generically, the quadric Q is nonsingular and the set of Ω, V such that $[V, \Omega \times A_2, \Omega \times A_3] \neq 0$ is a dense Zariski open subset of Q . The condition $[V, \Omega \times A_2, \Omega \times A_3] \neq 0$ implies that $\ell_4 = [\Omega, A_2, A_3] \neq 0$; indeed $[\Omega, A_2, A_3] = 0$ implies that $\Omega \times A_2$ and $\Omega \times A_3$ are colinear. Since $\ell_4 \neq 0$, the system of equations (9_{*i*}) for $i = 1, \dots, 6$ is equivalent to the system of equations (9₁, 10₂, 10₃, 11, 12, 13). The variables P appear only in the first three equations of this system and these equation can be rewritten as

$$\begin{aligned} V \cdot P &= 0 \\ (\Omega \times A_2) \cdot P &= -V \cdot C_2 - [\Omega, A_2, C_2] \\ (\Omega \times A_3) \cdot P &= -V \cdot C_3 - [\Omega, A_3, C_3]. \end{aligned} \quad (14)$$

This is a linear system in P which is a Cramer system since we have taken (Ω, V) in the quadric surface Q such that its determinant $[V, \Omega \times A_2, \Omega \times A_3]$ does not vanish. Hence there is a unique P satisfying the system (14), and this P is given by the Cramer formulas, i.e. as a rational function (of degree 3) in Ω, V . This P is on the surface Σ , since P, V, Ω satisfy the reciprocity equations, which implies that the rank of the system of Plücker coordinates of the limbs for P is < 6 . So we get a rational mapping, which we denote by Pos , from Q to Σ . The uniqueness of the solution for the Cramer system (14) implies that the composition $\text{Pos} \circ \text{Rec}$ is the identity of Σ . The fact that there is only a line of screws reciprocal to the Plücker coordinates of the limbs for a configuration P in Σ shows that the composition $\text{Rec} \circ \text{Pos}$ is the identity on Q .

In conclusion, we have proved the following theorem.

Theorem 8 *The regular mapping $\text{Rec} : \Sigma \rightarrow Q$ is a birational equivalence to the quadric surface $Q \in \mathbb{P}^5(\mathbb{R}(\text{SO}(3)))$, defined over $\mathbb{R}(\text{SO}(3))$.*

Corollary 9 *The cubic surface Σ is rational over $\mathbb{R}(\text{SO}(3))$.*

Proof. The cubic surface Σ has the $\mathbb{R}(\text{SO}(3))$ -rational point $P = 0$. Hence, the quadric surface Q has the $\mathbb{R}(\text{SO}(3))$ -rational point $\text{Rec}(0)$. It follows that Q is rational over $\mathbb{R}(\text{SO}(3))$ and since Σ is $\mathbb{R}(\text{SO}(3))$ -birational to Q , it is also rational over $\mathbb{R}(\text{SO}(3))$. \square

Theorem 10 *For a generic Gough-Stewart platform, the hypersurface $\text{Sing} \subset \text{SE}(3)$ of singular configurations is rational over \mathbb{R} .*

Proof. As already mentioned, this is a consequence of Corollary 9 and of the fact that $\text{SO}(3)$ is rational over \mathbb{R} . \square

2.4 Case study continued

We continue here the computations for the cubic surface Sing_R where the geometry of the Gough-Stewart platform and the orientation are those of section 1.4. Let us denote by $(\omega_1, \omega_2, \omega_3, v_1, v_2, v_3)$ the coordinates of a twist. We compute the variety of reciprocal twists as follows: we form the ideal \mathfrak{Rec} generated by the reciprocal products of the twist with the Plücker coordinates of the limbs and the Jacobian determinant, then we eliminate the variables x, y, z . The ideal obtained is the homogeneous ideal generated by

$$\begin{aligned} &72 \omega_1 - 42 \omega_2 + 16 v_1 + 21 v_3 + 35 v_2, \\ &18 \omega_3 + 30 \omega_2 - 26 v_1 - 15 v_3 + 5 v_2, \\ &168 \omega_2^2 + 80 \omega_2 v_1 + 522 \omega_2 v_3 - 296 v_1 v_3 - 159 v_3^2 \\ &\quad - 518 \omega_2 v_2 + 264 v_1 v_2 + 212 v_3 v_2 - 165 v_2^2 \end{aligned} \tag{15}$$

There are two linear equations and one quadratic: this is the ideal of a quadric surface in $\mathbb{P}^5(\mathbb{R})$.

We compute formulas for the reciprocal twist (only defined up to a scalar factor, as a point in $\mathbb{P}^5(\mathbb{R})$) in the following way: eliminating variables $\omega_1, \omega_2, \omega_3, v_3$ from the ideal \mathfrak{Rec} , we obtain an ideal which contains a polynomial which is linear homogeneous in v_1, v_2 and quadratic in x, y, z . This gives v_2/v_1 as a rational function of degree 2 in x, y, z . Proceeding in the same way to compute $v_3/v_1, \omega_1/v_1, \omega_2/v_1, \omega_3/v_1$ and chasing

denominator (the same for all expressions), we arrive at

$$\begin{aligned}
v_1 &= -80x^2 + 55xy - 109xz + 140y^2 - 14yz - 84z^2 \\
&\quad + 376x + 774y + 20z + 1392 \\
v_2 &= 52x^2 - 131xy + 164xz + 68y^2 - 62yz + 24z^2 \\
&\quad - 473x + 78y - 130z - 426 \\
v_3 &= 156x^2 - 245xy + 180xz - 36y^2 - 102yz + 24z^2 \\
&\quad - 663x - 578y - 234z - 1410 \\
\omega_1 &= -53x^2 + 92xy - 108xz + 338x + 248y - 48z + 736 \\
\omega_2 &= -53xy + 92y^2 - 108yz - 3x + 496y - 300z + 732 \\
\omega_3 &= -53xz + 92yz - 108z^2 + 127x - 212y + 370z - 266
\end{aligned} \tag{16}$$

These formulas are degree 2 polynomials in x, y, z . The formulas define a point in $\mathbb{P}^5(\mathbb{R})$ except when they all vanish. We compute the ideal generated by these quadratic polynomials in x, y, z and the cubic equation of the surface of singularities. We find that this ideal is precisely the ideal of the two lines corresponding to factor F_2 .

Nevertheless, the regular mapping which associates to a point (x, y, z) of the cubic singularity surface, a point of $\mathbb{P}^5(\mathbb{R})$ representing the line of reciprocal twists is also well defined on these two lines. Indeed, at every point in the surface of singularities there is a nonzero cofactor in the Jacobian matrix, and the cofactors on the same row as this nonzero cofactor are the coordinates of a nonzero reciprocal twist. These cofactors are polynomials of degree 3 in x, y, z . We shall explain in section 3.2 why we obtain degree 2 in the formulas above instead of degree 3.


We next compute the image of the five lines corresponding to factor F_5 by the reciprocal twist mapping. This is again the computation of an elimination ideal, and the computed ideal is zero-dimensional, of degree 5. It is actually the ideal of five distinct points of the quadric, all real in the case under consideration. The reciprocal twists corresponding to these five points are self-reciprocal. The figure 2.4 is in the 3-space containing the quadric, in its affine chart given by $\omega_2 = 1$, using coordinates v_1, v_2, v_3 . The quadric image of the reciprocal twist mapping is in blue, the five points in red, and the intersection of the quadric of self-reciprocal twists with the 3-space is in green.

It is also possible to compute formulas for the inverse rational mapping from the quadric surface in $\mathbb{P}^5(\mathbb{R})$ to Sing_R : for instance, eliminating the variables y, z in the ideal $\mathfrak{R}\mathfrak{e}\mathfrak{c}$, we obtain x as a rational function of degree 2 in the V, Ω . Actually we compute formulas for the rational mapping to the projective closure $(\text{Sing}_R)^h$ in $\mathbb{P}^3(\mathbb{R})$. We use homogeneous coordinates $(w : x : y : z)$ for $\mathbb{P}^3(\mathbb{R})$, with $w = 0$ as plane at infinity.

$$\begin{aligned}
w &= 60v_3^2 + 83v_1v_3 + 42\omega_2v_1 - 20v_3v_2 - 35v_1v_2 + 72\omega_2v_2 - 120\omega_2v_3 - 16v_1^2 \\
x &= 162\omega_2v_3 + 54\omega_2v_2 - 48v_1v_3 - 48v_1v_2 - 213v_2^2 + 63v_3^2 - 42v_3v_2 \\
y &= -54\omega_2v_1 + 48v_1^2 + 288\omega_2v_3 - 357v_1v_3 - 144v_3^2 + 213v_1v_2 - 48v_3v_2 \\
z &= -162\omega_2v_1 - 288\omega_2v_2 - 63v_1v_3 + 399v_1v_2 + 48v_2^2 + 48v_1^2 + 144v_3v_2
\end{aligned} \tag{17}$$

Note that these formulas are of degree 2, whereas the Cramer formulas invoked in the argument for birationality would give degree 3. It is possible to check that formulas (16) and (17) give actually a birational isomorphism between $(\text{Sing}_R)^h$ and the quadric surface in $\mathbb{P}^5(\mathbb{R})$. One can also

Figure 3: The quadric surface of reciprocal twists in blue



compute the indetermination points of formulas (17) on the quadric surface: one finds five points, which are precisely the image of the five lines corresponding to F_5 . This indicates that the “reciprocal twist” regular mapping from Σ to Q may be extended to a regular mapping from the projective closure Σ^h to Q , which is the blowing-up of Q in five points. We shall prove this in the following section after investigating the points at infinity of the singularity locus, i.e. $\Sigma^h \setminus \Sigma$.

3 The projective closure of the singularity surface

We already know that the projective closure Σ^h of Σ in $\mathbb{P}^3(\mathbb{R}(\text{SO}(3)))$ is a nonsingular projective cubic surface for a generic Gough-Stewart platform. Indeed, we checked in section 1.4 that $(\text{Sing}_R)^h$ is nonsingular.

3.1 Singularity condition at infinity

The formula (2) can be rewritten as a sum indexed by the symmetric group \mathfrak{S}_6 , with $\epsilon(\sigma)$ denoting the signature of a permutation σ :

$$\det(\text{Jac}) = \frac{1}{36} \sum_{\sigma \in \mathfrak{S}_6} \epsilon(\sigma) [C_{\sigma(1)} + P, C_{\sigma(2)} + P, C_{\sigma(3)} + P] \\ [A_{\sigma(4)} \times (C_{\sigma(4)} + P), A_{\sigma(5)} \times (C_{\sigma(5)} + P), A_{\sigma(6)} \times (C_{\sigma(6)} + P)]. \quad (18)$$

The equation of the projective closure Σ^h is the homogenization of $\det(\text{Jac})$, say with homogenization variable w . The equation of its intersection with the plane at infinity $w = 0$ is the homogeneous part of degree 3 of $\det(\text{Jac})$. Using multilinearity in (18), we find that this homogeneous part is

$$H_3 = \frac{1}{4} \sum_{\sigma \in \mathfrak{S}_6} \epsilon(\sigma) [C_{\sigma(1)}, C_{\sigma(2)}, P] [A_{\sigma(4)} \times C_{\sigma(4)}, A_{\sigma(5)} \times P, A_{\sigma(6)} \times P]. \quad (19)$$

Since

$$[A_{\sigma(4)} \times C_{\sigma(4)}, A_{\sigma(5)} \times P, A_{\sigma(6)} \times P] = [A_{\sigma(4)}, C_{\sigma(4)}, P] [A_{\sigma(5)}, A_{\sigma(6)}, P], \quad (20)$$

we arrive to

$$\begin{aligned} H_3 &= \frac{1}{4} \sum_{\sigma \in \mathfrak{S}_6} \epsilon(\sigma) [C_{\sigma(1)}, C_{\sigma(2)}, P] [A_{\sigma(4)}, C_{\sigma(4)}, P] [A_{\sigma(5)}, A_{\sigma(6)}, P] \\ &= \frac{-1}{4} \sum_{\sigma \in \mathfrak{S}_6} \epsilon(\sigma) [A_{\sigma(1)}, A_{\sigma(2)}, P] [A_{\sigma(3)}, C_{\sigma(3)}, P] [C_{\sigma(4)}, C_{\sigma(5)}, P] \end{aligned} \quad (21)$$

In this formula we can replace the C with the B :

$$H_3 = \frac{-1}{4} \sum_{\sigma \in \mathfrak{S}_6} \epsilon(\sigma) [A_{\sigma(1)}, A_{\sigma(2)}, P] [A_{\sigma(3)}, B_{\sigma(3)}, P] [B_{\sigma(4)}, B_{\sigma(5)}, P] \quad (22)$$

The equality is proved using $C_i = B_i - A_i$ and Grassmann-Plücker relations such as:

$$\begin{aligned} [A_{\sigma(1)}, A_{\sigma(2)}, P] [A_{\sigma(4)}, A_{\sigma(5)}, P] - [A_{\sigma(1)}, A_{\sigma(4)}, P] [A_{\sigma(2)}, A_{\sigma(5)}, P] \\ + [A_{\sigma(1)}, A_{\sigma(5)}, P] [A_{\sigma(2)}, A_{\sigma(4)}, P] = 0 \end{aligned} \quad (23)$$

We summarize what we have seen in the following

Proposition 11 *The singularities at infinity of the Gough-Stewart platform in the orientation R are the directions of $P = (x \ y \ z)^T$ satisfying*

$$\sum_{\sigma \in \mathfrak{S}_6} \epsilon(\sigma) [A_{\sigma(1)}, A_{\sigma(2)}, P] [A_{\sigma(3)}, B_{\sigma(3)}, P] [B_{\sigma(4)}, B_{\sigma(5)}, P] = 0,$$

where $B_i = Rb_i$.

We can reformulate this condition as a condition concerning two planar hexagons. Let π_P the orthogonal projection in the direction of P . Then for $i = 1, \dots, 6$, we denote $\pi_P(A_i) = \alpha_i$ the projection of the vertices of the base and $\pi_P(B_i) = \beta_i$ the projection of the vertices of the platform after rotation R . So we have two hexagons in the plane orthogonal to P . The condition of singularity at infinity is given by:

$$\sum_{\sigma \in \mathfrak{S}_6} \epsilon(\sigma) [\alpha_{\sigma(1)}, \alpha_{\sigma(2)}] [\beta_{\sigma(3)}, \beta_{\sigma(4)}] [\alpha_{\sigma(5)}, \beta_{\sigma(5)}] = 0, \quad (24)$$

where the bracket denotes here 2×2 determinants. This condition is invariant under the following transformations:

1. The same affine transformation applied to all α_i 's and β_i 's. The determinants are multiplied by the determinant of linear part of the transformation.
2. A homothety (of ratio $k \neq 0$) on one of the hexagons. The product of determinants are multiplied by k^3 .
3. A translation (of vector u) on one of the hexagons. The invariance follows from Plücker relations.

4. Replace the β'_i s by any linear combination of β_i and α_i .

In addition, when the hexagons have three distinct vertices in common ($\alpha_1 = \beta_1, \alpha_2 = \beta_2, \alpha_3 = \beta_3$), the condition is satisfied if and only if the three lines (α_i, β_i) for $i = 4, 5, 6$ are concurrent or parallel.

3.2 Reciprocal twist at infinity

Recall that the twist $\begin{pmatrix} \Omega \\ V \end{pmatrix}$ is reciprocal to the system of Plücker coordinates of the six limbs if and only if

$$(C_i + P) \cdot V + (A_i \times (C_i + P)) \cdot \Omega = 0, \quad i = 1, \dots, 6. \quad (25)$$

The matrix of this system is Jac. In a singular configuration, the rank of this system is generically 5. In this case, if the rows of Jac with index $\neq i$ are linearly independent, the line of solutions of the system of equations is spanned by the vector whose coordinates are $\text{cof}(\text{Jac})_{i,j}$ for $j = 1, \dots, 6$. This gives formulas of degree 3 in P . The line of solutions is also spanned by the vector whose coordinates are $\sum_{i=1}^6 \text{cof}(\text{Jac})_{i,j}$ for $j = 1, \dots, 6$, if this vector is nonzero. It turns out that these sums of cofactors are polynomials of degree 2 in P ; this explains what we have seen in the case study in formulas (16).

The sum by columns of the cofactors in the first three columns is the vector

$$T_1 = \sum_{1 \leq i_1 < i_2 < i_3 \leq 6} (-1)^{i_1+i_2+i_3} \left(\sum_{\text{cyc}} (C_{i_2} + P) \times (C_{i_3} + P) \right) \\ [(A_{j_1} \times (C_{j_1} + P), A_{j_2} \times (C_{j_2} + P), A_{j_3} \times (C_{j_3} + P))] \quad (26)$$

where $j_1 < j_2 < j_3$ are the integers between 1 and 6 different from i_1, i_2, i_3 and the cyclic sum is taken over the powers of the cycle (i_1, i_2, i_3) . It will be more convenient to write the sum T_1 as a sum over all permutations in \mathfrak{S}_6 .

$$T_1 = \frac{1}{12} \sum_{\sigma \in \mathfrak{S}_6} \left(\varepsilon(\sigma) (C_{\sigma(2)} + P) \times (C_{\sigma(3)} + P) \right. \\ \left. [A_{\sigma(4)} \times (C_{\sigma(4)} + P), A_{\sigma(5)} \times (C_{\sigma(5)} + P), A_{\sigma(6)} \times (C_{\sigma(6)} + P)] \right) \quad (27)$$

We develop this expression using multilinearity in the cross-product and in the mixed product. We remark that all terms containing only four among the six indices $\sigma(i)$ disappear in the sum over all permutations (due to the transposition on the missing indices). We remark also that the mixed product

$$[A_{\sigma(4)} \times P, A_{\sigma(5)} \times P, A_{\sigma(6)} \times P]$$

is zero because the three vectors are linearly dependant. This explains why all terms of degree 3 disappear in the sum and why the only remaining

terms of degree 2 are

$$\begin{aligned}
V_\infty &= \frac{1}{4} \sum_{\sigma \in \mathfrak{S}_6} \varepsilon(\sigma) [A_{\sigma(4)} \times C_{\sigma(4)}, A_{\sigma(5)} \times P, A_{\sigma(6)} \times P] C_{\sigma(2)} \times C_{\sigma(3)} \\
&= \frac{1}{4} \sum_{\sigma \in \mathfrak{S}_6} \varepsilon(\sigma) [A_{\sigma(4)}, C_{\sigma(4)}, P] [A_{\sigma(5)}, A_{\sigma(6)}, P] C_{\sigma(2)} \times C_{\sigma(3)} \\
&= \frac{-1}{4} \sum_{\sigma \in \mathfrak{S}_6} \varepsilon(\sigma) [A_{\sigma(1)}, A_{\sigma(2)}, P] [A_{\sigma(3)}, C_{\sigma(3)}, P] C_{\sigma(4)} \times C_{\sigma(5)}
\end{aligned} \tag{28}$$

The sum by columns of the cofactors in the last three columns is the vector

$$\begin{aligned}
T_2 &= \sum_{1 \leq i_1 < i_2 < i_3 \leq 6} (-1)^{i_1+i_2+i_3} \left([C_{i_1} + P, C_{i_2} + P, C_{i_3} + P] \right. \\
&\quad \left. \sum_{\text{cyc}} (A_{j_1} \times (C_{j_1} + P)) \times (A_{j_2} \times (C_{j_2} + P)) \right)
\end{aligned} \tag{29}$$

where $j_1 < j_2 < j_3$ are the integers between 1 and 6 different from i_1, i_2, i_3 and the cyclic sum is taken over the powers of the cycle (j_1, j_2, j_3) . Here also it will be more convenient to write the sum T_2 as a sum over all permutations in \mathfrak{S}_6 .

$$\begin{aligned}
T_2 &= \frac{1}{12} \sum_{\sigma \in \mathfrak{S}_6} \varepsilon(\sigma) [C_{\sigma(1)} + P, C_{\sigma(2)} + P, C_{\sigma(3)} + P] \\
&\quad (A_{\sigma(4)} \times (C_{\sigma(4)} + P)) \times (A_{\sigma(5)} \times (C_{\sigma(5)} + P))
\end{aligned} \tag{30}$$

We develop this expression using multilinearity in the mixed product and in the cross-product. Here also, terms where only four indices among the $\sigma(i)$ are present disappear in the sum over all permutations. This explains why all terms of degree 3 in P disappear, and why the terms of degree 2 in P which remain are only:

$$\begin{aligned}
\Omega_\infty &= \frac{1}{12} \left(\sum_{\sigma \in \mathfrak{S}_6} \varepsilon(\sigma) [C_{\sigma(1)}, C_{\sigma(2)}, C_{\sigma(3)}] (A_{\sigma(4)} \times P) \times (A_{\sigma(5)} \times P) \right) \\
&= \frac{1}{12} \left(\sum_{\sigma \in \mathfrak{S}_6} \varepsilon(\sigma) [C_{\sigma(1)}, C_{\sigma(2)}, C_{\sigma(3)}] [A_{\sigma(4)}, A_{\sigma(5)}, P] \right) P
\end{aligned} \tag{31}$$

Note that Ω_∞ is the product of the vector P by a linear form in P ; this confirms the result of the computation in case study 2.4.

The notations V_∞ and Ω_∞ are explained by the fact that $\begin{pmatrix} \Omega_\infty \\ V_\infty \end{pmatrix}$ is actually the reciprocal twist at infinity, as explained in the following

Proposition 12 *For a generic Gough-Stewart platform, the reciprocal twist mapping $\text{Rec} : \Sigma \rightarrow Q$ can be extended to a regular mapping $\text{Rec}^h : \Sigma^h \rightarrow Q$. The image by Rec^h of a point at infinity in the direction P in Σ^h is given by the homogeneous second degree formulas (28) for V_∞ and (31) for Ω_∞ , provided that V_∞ and Ω_∞ do not both vanish at P .*

Proof. We know that $\begin{pmatrix} T_2 \\ T_1 \end{pmatrix}$ give degree 2 formulas for Rec at points of Σ where T_1 and T_2 do not both vanish. Hence, the homogeneous quadratic part $\begin{pmatrix} \Omega_\infty \\ V_\infty \end{pmatrix}$ give a regular extension of Rec at points at infinity of Σ^h in the direction of P , when Ω_∞ and V_∞ do not both vanish. We note that, in the plane at infinity, Ω_∞ vanish on a line (whose equation is the linear form $L(P)$ such that $\Omega_\infty = L(P)P$), while each component of V_∞ vanishes on a conic. We check in the case study 3.4 that these curves have two points in common which are on the cubic curve of the points at infinity of Σ^h ; these two points are, of course, the points at infinity of the two lines on Sing_R^h corresponding to F_2 . We check also that the formulas for the reciprocal twist at infinity given by the homogeneous part of degree 3 of the cofactors of Jac on the first line do not vanish at these two points. This shows that for a generic Gough-Stewart platform, the regular extension Rec^h is well defined at every point at infinity of Σ^h : when both Ω_∞ and V_∞ vanish, it is given by the homogeneous part of degree 3 of the cofactors of Jac on the first line. \square

Proposition 13 *For a generic Gough-Stewart platform, the reciprocal twists at points at infinity of Σ^h in the direction P are all self-reciprocal. They are the twists of a rotation with axis parallel to P , or of a translation orthogonal to P .*

Proof. From formulas (28) for V_∞ and Proposition (11), we obtain $V_\infty \cdot P = 0$ for all point at infinity of Σ^h in direction P . Since $\Omega_\infty = L(P)P$, the statements of the Proposition follow. \square

3.3 Σ^h as the blowing-up in five points of a quadric surface

Theorem 14 *For a generic Gough-Stewart platform, the reciprocal twist regular mapping $\text{Rec}^h : \Sigma^h \rightarrow Q$ is the blowing-up of the projective quadric surface Q in five points. The five exceptional divisors are five lines on Σ^h , forming a S_5 intersecting both lines of a S_2 (using notations of Theorem 3), both S_5 and S_2 being defined over $\mathbb{R}(\text{SO}(3))$. A point of Σ belongs to one of the lines of the S_5 if and only if its reciprocal twist is the twist of a pure rotation; the axis of this rotation is then a line intersecting the six limbs of the platform.*

Proof. We know from Proposition 12 that $\text{Rec}^h : \Sigma^h \rightarrow Q$ is a regular on the whole of Σ^h , and we know from Theorem 8 that it is a birational equivalence. From this we deduce that Rec^h is the composition of a sequence of blowing-ups at points (see for instance [5], corollary 5.4 p.411). Since the divisor class group of a nonsingular projective cubic surface is \mathbb{Z}^7 ([5] Proposition 4.8 p.401) and the divisor class group of a nonsingular projective quadric surface is \mathbb{Z}^2 , Rec^h is the blowing-up of Q in five points. These five points are distinct points of Q , as checked in the case study, so the five exceptional divisors have self-intersection equal to -1 and are mutually skew lines on Σ^h . Since Rec^h is defined over $\mathbb{R}(\text{SO}(3))$, the five exceptional divisors form a S_5 over $\mathbb{R}(\text{SO}(3))$. There are two possible types of S_5 , one of them characterized as the five lines intersecting both lines of an S_2 (see [11]). Actually, the five exceptional divisors obtained by blowing up five points in a quadric form a S_5 of this type; we defer the proof of this fact to Lemma 15 below.


Since the reciprocal twist is constant along each line of the S_5 , it is equal to the reciprocal twist of the point at infinity of this line. Hence, by Proposition 13, it is the twist of a rotation or of a translation. If the reciprocal twist at $P \in \Sigma$ were the twist $\begin{pmatrix} 0 \\ V \end{pmatrix}$ of a translation in the direction of vector V then, for every vector U orthogonal to V , the twist $\begin{pmatrix} 0 \\ V \end{pmatrix}$ would be reciprocal to the systems of Plücker coordinates of all limbs in position $P + U$ and hence $P + U$ would be in Σ . This cannot be since Σ contains no plane. This shows that the reciprocal twist at each point of Σ belonging to a line of the S_5 is the twist of a pure rotation; the reciprocity conditions means that every limb of the platform intersects (or is parallel to) the axis of this rotation.

Reciprocally, suppose that the reciprocal twist at a point $P \in \Sigma$ is the twist $\begin{pmatrix} \Omega \\ V \end{pmatrix}$ of a pure rotation, i.e. $\Omega \neq 0$ and $\Omega \cdot V = 0$. Then, for all scalars λ , $P + \lambda\Omega$ is still in Σ , with reciprocal twist $\begin{pmatrix} \Omega \\ V \end{pmatrix}$. Hence the line of $P + \lambda\Omega$ is one of the exceptional divisors in the S_5 . \square

Lemma 15 *The five exceptional divisors obtained by blowing up a quadric surface in five distinct points form an S_5 with two transversals forming an S_2 .*

Proof. Let p_1, \dots, p_4, q be the five distinct points on the quadric surface. Let d_5, d_6 be the two lines on the quadric through q . Blow up q ; the strict transforms of d_5 and d_6 (which we still denote by d_5 and d_6) have self-intersection -1 and one can blow them down to obtain a projective plane with six distinct points $p_1, \dots, p_4, d_5, d_6$ (abusing notation, we denote with the same letter points and the exceptional divisors above them). The five exceptional divisors of the blowing up of the quadric are, in the blowing up of the plane at the six points $p_1, p_2, p_3, p_4, d_5, d_6$, the exceptional divisors above p_1, p_2, p_3, p_4 and the strict transform of the line $(d_5 d_6)$. The S_5 so obtained has two transversals, which are the strict transforms of two conics in the plane through p_1, p_2, p_3, p_4, d_5 and p_1, p_2, p_3, p_4, d_6 , respectively. \square

Figure 4: Blowing up the quadric vs blowing up the plane



3.4 Case study, the end

The part at infinity of the cubic surface, which is $\text{Sing}_R^\infty = \text{Sing}_R^h \setminus \text{Sing}_R$, is the nonsingular cubic curve in the projective plane with homogeneous coordinates x, y, z given by the equation

$$80x^3 - 107yx^2 - 47zx^2 - 9y^2x + 95zxy - 96z^2x - 68y^3 + 98zy^2 + 78z^2y - 24z^3 \quad (32)$$


The formulas (16) for the reciprocal twist extend to the following homogeneous quadratic formulas:

$$\begin{aligned} \Omega_\infty &= (-53x + 92y - 108z) \begin{pmatrix} x \\ y \\ z \end{pmatrix} \\ V_\infty &= \begin{pmatrix} -80x^2 + 55xy - 109xz + 140y^2 - 14yz - 84z^2 \\ 52x^2 - 131xy + 164xz + 68y^2 - 62yz + 24z^2 \\ 156x^2 - 245xy + 180xz - 36y^2 - 102yz + 24z^2 \end{pmatrix} \end{aligned} \quad (33)$$

These formulas have two points of indetermination on Sing_R^∞ which are, of course, the points at infinity of the two lines of indetermination for the formulas (16) on Sing_R corresponding to F_2 .

The figure 3.4 shows the two points of indetermination in the affine part of the plane at infinity given by $x = 1$. The solid black curve is the cubic Sing_R^∞ , the dashed brown line is the line whose equation is the linear form factor of Ω_∞ , and the dotted conics in red, green and blue are those whose equations are the components of V_∞ . All these curves concur to the indetermination points marked with diamonds.

Figure 5: Points of indetermination at infinity of the quadratic formulas for the reciprocal twist



We now check that the reciprocal twist mapping is also defined at the indetermination points by other formulas: precisely, we use the degree 3 homogeneous part of the cofactors of the coefficients in the last line of the Jacobian matrix Jac:

$$\begin{aligned} \Omega_\infty &= (-x^2 + 7yx - 12zx - 6z^2 + 7y^2 - 7zy) \begin{pmatrix} x \\ y \\ z \end{pmatrix} \\ V_\infty &= \begin{pmatrix} -4yx^2 - 12zx^2 + 17y^2x + zxy - 18z^2x + 4y^3 + 7zy^2 - 5z^2y - 6z^3 \\ 4x^3 - 17yx^2 + 14zx^2 - 4y^2x + 11zxy + 6z^2x + 2zy^2 + 4z^2y \\ 12x^3 - 15yx^2 + 18zx^2 - 18y^2x - zxy + 6z^2x - 2y^3 - 4zy^2 \end{pmatrix} \end{aligned} \quad (34)$$

The homogeneous ideal generated by the ideal of indetermination of the quadratic formulas (33) and the quadratic form $-x^2 + 7yx - 12zx - 6z^2 + 7y^2 - 7zy$ factor of Ω_∞ in (34) contains the third power of the maximal ideal generated by x, y, z . This shows that the reciprocal twist mapping is well defined as a regular map on the whole of Sing_R^h .

Conclusion

We have proved the rationality of the locus of singular configurations of the general Gough-Stewart platform. We have moreover related the rationality with the reciprocal twist mapping, which has a kinematic relevance; the lines of reciprocal twists form a quadric surface, and this fact may be useful for further studies on the singularities of a Gough-Stewart platform. We have also related a group of five lines on the cubic surface of singularities with special singular configurations for which the reciprocal twist is the twist of a pure rotation. This interplay between the classical algebraic geometry of cubic surfaces and kinematic properties of a parallel robot is rather fascinating.

We conclude with two questions encountered in this paper for which we have no satisfactory answer.

The first question concerns the characterization of singularities at infinity in section 3.1. We have seen that it can be expressed in terms of a relation between two planar hexagons (Equation 24). We have not been able to uncover the geometric significance of this relation. In the case of the planar 3-RPR, the singularities at infinity may be characterized by the fact that two triples of aligned points can be transformed one into another by an affine mapping [2].

The second question is related to the fact that the reciprocal twist mapping extends to a regular mapping defined on the whole projective closure of the surface of singularities. We have followed a rather cumbersome way, with the help of a computation in a specific example, to show that this is indeed generically the case. It would be much nicer if the following assertion were true.

Let $A(t)X = 0$ be a homogeneous linear system of n equations in n unknowns with coefficients $a_{i,j}(t)$ polynomials of degree 1 in parameters $t = (t_1, \dots, t_p)$. Assume that the projective closure S^h (in the p -dimensional projective space) of the set S of parameters t such that $\det(A(t)) = 0$ is a smooth projective hypersurface. Then the mapping which associates to $t \in S$ the line of solutions of $A(t)X = 0$ extends to a regular mapping from S^h to the $n - 1$ -dimensional projective space.

This assertion is not obvious when the degree of $\det(A(t))$ is strictly smaller than n , which happens in our case since the degree of the equation

of the surface of singularities w.r.t. the position variables is 3 instead of 6. We have no idea whether this assertion holds true.

References

- [1] Sandipan Bandyopadhyay and Ashitava Ghosal. Geometric characterization and parametric representation of the singularity manifold of a 6-6 Stewart platform manipulator. *Mechanism and Machine Theory*, 41(11):1377 – 1400, 2006.
- [2] Michel Coste. Asymptotic singularities of planar parallel 3-RPR manipulators. In Jadran Lenarcic and Manfred Husty, editors, *Latest Advances in Robot Kinematics*, pages 35–42, Dordrecht, 2012. Springer Netherlands.
- [3] Michel Coste and Seydou Moussa. On the rationality of the singularity locus of a Gough-Stewart platform - biplanar case. *Mechanism and Machine Theory*, 87:82 – 92, 2015.
- [4] Karine Doyon, Clment Gosselin, and Philippe Cardou. A vector expression of the constant-orientation singularity locus of the Gough-Stewart platform. *Journal of Mechanisms and Robotics*, 5:034502–4, 2013/06/24.
- [5] R. Hartshorne. *Algebraic Geometry*. Graduate Texts in Mathematics. Springer New York, 2013.
- [6] Manfred Husty and Clment Gosselin. On the singularity surface of planar 3-RPR parallel mechanisms. *Mechanics Based Design of Structures and Machines*, 36(4):411–425, 2008.
- [7] A. Karger. Stewart-Gough platforms with simple singularity surface. In Jadran Lennarčič and B. Roth, editors, *Advances in Robot Kinematics*, pages 247–254, Dordrecht, 2006. Springer Netherlands.
- [8] J.P. Merlet. *Parallel Robots*. Solid Mechanics and Its Applications. Springer Netherlands, 2006.
- [9] D. Mumford. *Algebraic Geometry I: Complex Projective Varieties*. Classics in Mathematics. Springer Berlin Heidelberg, 1995.
- [10] Georg Nawratil. Stewart Gough platforms with non-cubic singularity surface. *Mechanism and Machine Theory*, 45(12):1851 – 1863, 2010.
- [11] H. P. F. Swinnerton-Dyer. The birationality of cubic surfaces over a given field. *Michigan Math. J.*, 17(4):289–295, 12 1970.