



**HAL**  
open science

# Nontechnical Skills in Neurosurgery A Systematic Review of the Literature

Pierre-Louis Henaux, Pierre Jannin, Laurent Riffaud

► **To cite this version:**

Pierre-Louis Henaux, Pierre Jannin, Laurent Riffaud. Nontechnical Skills in Neurosurgery A Systematic Review of the Literature. *World Neurosurgery*, 2019, 130, pp.e726-e736. 10.1016/j.wneu.2019.06.204 . hal-02280254

**HAL Id: hal-02280254**

**<https://univ-rennes.hal.science/hal-02280254>**

Submitted on 16 Dec 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Non-technical skills in Neurosurgery : a systematic review of the literature**

Pierre-Louis Hénaux<sup>1,2</sup>, MD, PhD; Pierre Jannin<sup>2</sup>, PhD ; Laurent Riffaud<sup>1,2</sup>, MD, PhD

<sup>1</sup>Department of Neurosurgery, Rennes University Hospital, 2, rue Henri Le Guilloux 35033 Rennes, Cedex 9, France

<sup>2</sup>INSERM MediCIS, Unit U1099 LTSI, Rennes 1 University, Faculty of Medicine, 2 Avenue du Professeur Léon Bernard, 35043 Rennes Cedex, France

Authors' e-mail adress : [pierre-louis.henaux@etudiant.univ-rennes1.fr](mailto:pierre-louis.henaux@etudiant.univ-rennes1.fr); [pierre.jannin@univ-rennes1.fr](mailto:pierre.jannin@univ-rennes1.fr); [laurent.riffaud@chu-rennes.fr](mailto:laurent.riffaud@chu-rennes.fr)

Corresponding author : Dr. Pierre-Louis Henaux

Department of Neurosurgery, Pontchaillou University Hospital,

2, rue Henri Le Guilloux 35033 Rennes Cedex 9, France

Phone Number: + 33 2 99 28 42 77

Fax: + 33 2 99 28 41 80

Email: [pierre-louis.henaux@etudiant.univ-rennes1.fr](mailto:pierre-louis.henaux@etudiant.univ-rennes1.fr)

**Key-Words**

Non-Technical Skills; Neurosurgery; Interpersonal Skills; Cognitive Skills; Patient Safety

**Abbreviations list**

ACGME: Accreditation Council for Graduate Medical Education

CPD: continuing professional development

NOTECHS: NonTECHnical Skills

NOTSS: Non Technical Skills for Surgeons

NTS: non-technical skills

OTAS: Observational Teamwork Assessment for Surgery

TS: technical skills

UE: undesirable events

ACCEPTED MANUSCRIPT

**ABSTRACT****Background**

Undesirable events in the neurosurgery operating theatre are surprisingly less often the result of a technical error than of a dysfunction linked to non-technical skills (NTS). The essential aim of our study was a systematic literature study published on NTS in neurosurgery. The secondary objective was identification of a list of NTS more specific to neurosurgery in order to define on that basis the training needs of neurosurgery trainees.

**Methods**

MEDLINE and The Cochrane Database of Systematic Reviews were searched according to the PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analysis) statement. Research initially identified 2132 articles. 21 were eligible for systematic study. Data were extracted from the articles regarding study design, sample size, NTS assessed, assessment tools and the key results were collected.

**Results**

Interpersonal skills (communication, teamwork), cognitive skills (decision making, situation awareness) and personal resource factors (coping with stress or fatigue) were specifically identified. No article used assessment tools such as NOTECHS, NOTSS or OTAS. They were carried out in a real environment in 11 cases, in a simulated environment in 9 cases and during theoretical teaching in 1 case.

**Conclusions**

Very few studies have been carried out concerning neurosurgical NTS, despite increasing numbers of articles over the last few years on NTS in other domains of surgery. Society today is concentrating more and more on the quality and safety of medical care. The development and

application of NTS assessment tools is therefore essential in order to provide assistance in the training of future neurosurgeons.

ACCEPTED MANUSCRIPT

**INTRODUCTION**

Neurosurgery is a discipline requiring knowledge of numerous hyperspecialist domains such as vascular, tumoral, malformation, degenerative and spinal pathologies. As well as this declarative knowledge of neurosurgical pathologies and their associated indications for operation, the surgeon has to acquire technical skills both in accordance with good practice and under ongoing development, for example microsurgery, computer-assisted surgery<sup>1-7</sup> and robotics<sup>8</sup>. Even with perfect mastery of this knowledge and of the technical skills (TS) involved, a certain number of undesirable events (UE) are to be observed in the neurosurgery operating theatre<sup>9-16</sup>. These events may have irreversible consequences involving serious functional deficits or even life-threatening situations<sup>17,18</sup>. Surprisingly, they are less often the result of a technical error than of a dysfunction linked to non-technical skills (NTS)<sup>19</sup>. This relationship between NTS and UE has led the surgical community to explore in greater depth safety and quality issues with increasing numbers of publications on NTS over the last few years. NTS can be divided into two groups according to the taxonomy developed by Yule et al.<sup>20</sup>. The first includes interpersonal skills: communication, leadership, teamwork, briefing, planning and preparation, resource management, requests for advice and feedback, attitudes to pressure, stress and fatigue. The second group covers cognitive skills including awareness of the situation, mental disposition, risk assessment, anticipation of problems, decision-making, adaptation strategies, flexibility or workload. In real conditions, interpersonal skills and cognitive skills may be involved simultaneously. For example, poor communication between the surgeon and the anesthesiologist during an unexpected haemorrhage in the operating theatre (NTS involved: communication and situational awareness) may have life-threatening consequences. Publications on this subject still rely for the moment on non-standardized methods. NTS assessment is rendered problematical by a methodology difficult to put into practice (necessitating robust psychometric testing) involving multiple variables which are themselves not easily controlled<sup>21-30</sup>. However, it is certain that NTS exercise a positive or negative influence on TS depending on whether or not they are applied<sup>19</sup>. The safety and quality of care is a universal and

increasing preoccupation among the population<sup>9,17,31-34</sup>: surgeons have taken this into account and, as in other sensitive domains such as the army<sup>35</sup> or navy, they are beginning to develop “firewall” tools such as checklists<sup>9,36-40</sup>. These NTS, still ill-defined, are only just beginning to emerge in surgeon training objectives and are even less present in the confidential world of neurosurgery<sup>25,41-54</sup>.

The essential aim of our study was a systematic study of the literature published on NTS in the neurosurgical domain. The secondary objective was identification, from the existing literature, of a list of NTS more specific to neurosurgery in order to define on that basis the training needs of neurosurgery trainees.

**MATERIAL AND METHODS**

We conducted a systematic review following the PRISMA statement<sup>55</sup>. Ethics committee approval was not required for our research protocol. We did not register our review protocol in the PROSPERO database.

Using the databases PubMed-MEDLINE and The Cochrane Database of Systematic Reviews, we combined two A and B categories using the Boolean operator “AND”, Category A was the following : Adaptation, Psychologic OR Adjustment, Psychological OR Behavior, Adaptive OR Coping Behavior OR Coping Skills OR Psychological Adaptation OR Psychological Adjustment OR skills, Category B was: neurosurgery OR neurosurgical procedures. We defined the following limits: publication date between January 1980 and December 2018 ; articles concerning studies on humans, excluding those on animals ; publications in English and French. The final research was carried out on 2<sup>nd</sup> January 2019. The lists of article references included were also researched to find other references such as grey literature (unpublished studies with limited distribution, for example conference abstracts) and published studies not identified using our initial search strategy. This led us to add 2 articles for which we made a complete assessment of the text. We assessed the collected quotations, sifting through the titles and abstracts to identify the pertinent articles on the basis of the following predefined inclusion criteria :

1) Data concerning NTS OR interpersonal behaviour OR leadership OR teamwork OR planning and preparation OR feedback OR stress and pressure OR fatigue OR cognitive skills OR anticipation of problems OR decision-making OR flexibility OR adaptation strategies

AND

2) Data concerning the neurosurgical domain

AND

3) Data on training needs OR teaching OR simulation systems including robotics. These criteria were chosen to identify the articles specifically examining NTS in neurosurgery, the resulting training needs and the integration of these skills to simulation systems.

The data was extracted from the articles concerned using a structured data résumé (Figure 1), to ensure coherent assessment of each article. Using this method, details of the study design, the size of the sample, the NTS assessed, the assessment tools and the key results were collected.

ACCEPTED MANUSCRIPT

**RESULTS****Selection of articles**

The diagram summarizing the research strategy is shown in Figure 2.

Research initially identified 2132 articles. In all, 21 articles were eligible for systematic study.

**Populations studied**

12 articles concerned analysis of neurosurgery residents<sup>43,44,54,56-64</sup>. 12 others studied senior neurosurgeons<sup>33,35,58,62,64-71</sup> and 3 others directors of training (also senior neurosurgeons)<sup>44,58,61</sup>. Six others dealt with the whole team of a neurosurgery operating theatre<sup>35,65,66,68-70</sup> and one article studied the whole team of a neurosurgery department<sup>33</sup>.

**Study designs**

7 were experimental studies<sup>56,57,59,60,63,64,71</sup>, 7 were surveys<sup>43,44,54,58,61,62,66</sup>, 5 were interventional studies<sup>33,35,65,67,69</sup>, and 2 were observational studies<sup>68,70</sup>.

**Environment**

These studies were carried out :

- in a real environment in 11 cases : 10 in the operating theatre<sup>33,35,44,58,61,65,66,68-70</sup>, 1 in a neurosurgery department<sup>33</sup>, and 4 in consultation<sup>44,58,61,67</sup>,
- in a simulated environment in 9 cases<sup>54,56,57,59,60,62-64,71</sup>,
- during theoretical teaching in 1 case<sup>43</sup>.

**Non-technical surgical skills assessed**

Interpersonal NTS were specifically assessed in 9 studies<sup>33,43,44,59,60,63,67,68,71</sup>.

Cognitive NTS were specifically evaluated in 9 studies<sup>44,56,57,62,64–66,69,70</sup>. Finally, neurosurgery NTS were assessed overall in 6 studies without differentiation between the two types<sup>35,54,57,58,61,69</sup>.

**Assessment tools and key results of each study**

None of the articles selected used assessment tools such as NOTECHS, NOTSS or OTAS, all of which have all been established as valid in the literature concerning assessment of NTS in surgery.

*Studies in a real environment*

Neurosurgery interns complained of lack of teaching on behaviour and interpersonal relations in a questionnaire designed by Cusimano *et al.*<sup>44</sup> given to a population of 30 interns. Open questions with content analysis and double-scale quantification were carried out both among neurosurgery interns and among heads of teaching programmes. These revealed a positive correlation between excessive workload and voluntary withdrawal, and between dismissal and interpersonal skills during the neurosurgery internship. According to Pathak *et al.*<sup>67</sup> Senior neurosurgeons demonstrated good productivity and flexibility, but appeared to have difficulty in remaining objective and making themselves understood in consultation. This study was carried out during neurosurgery consultations, and the sample included 12 senior neurosurgeons with at least 10 years' experience. The surgeon was assessed by an outside observer using an assessment scale including : productivity, flexibility, objectivity and comprehensibility. Fargen *et al.*<sup>69</sup> studied the effect of the introduction of a checklist for neurointerventional procedures in neurovascular pathology. 71 procedures were carried out 4 weeks before the introduction of the checklist and 60 others 4 weeks after it. The checklist brought about a significant improvement in communication within the team and a significant reduction in the number of undesirable events. Ferroli *et al.*<sup>35</sup> assessed a notification system inspired by aviation and applied to the neurosurgical domain. The entire team of a

neurosurgery department participated in the study and reported incidents occurring in the service.

14 incidents were reported and analyzed. 9 were due to human factors and 3 to questions of organization. Lau *et al.*<sup>33</sup> assessed and installed a safety education video in a neurosurgery department. It had been made over several months by a multidisciplinary team including neurosurgeons, anaesthetists, nurses and professionals from the quality and video domains. This team ensured that all the safety parameters involved in a neurosurgical procedure were collected and formalized in video format. McLaughlin *et al.*<sup>66</sup> examined the interest of establishing a checklist before the beginning of an operation. 98,9% of those questioned judged this to be in the interest of the patient's safety, and 97,8% thought it increased team spirit. All members of the team judged that the surgeon should be present at the time of the checklist and according 76,3% thought that the surgeon should carry it out in person. Osvald *et al.*<sup>65</sup> studied the impact of the introduction of a checklist in a neurosurgery operating theatre. 8795 surgical procedures were carried out without checklist between January 2007 and December 2010, with two error reports : one procedure on the wrong side of the skull and one at the wrong level on the lumbar vertebrae. Between January 2011 and June 2012 no errors of side or level were reported. Couat *et al.*<sup>68</sup> through an observational study of the operating theatre using video analysis and ethographical methods, demonstrated that 33% of errors committed in the operating theatre were at least partly caused by poor communication or organisation management. Michinov *et al.*<sup>70</sup> in an operating theatre observational study of deep brain stimulation showed that explicit coordination, awareness of the situation and leadership are the NTS most involved in deep brain stimulation by the neurosurgeon. What is more, these aspects of behaviour differ according to the time of operation and the other members of the theatre team with whom the surgeon is communicating. Brandman *et al.*<sup>61</sup> also used a survey which underlined the fact that heads of programmes considered NTS more important than did the interns, with an emphasis on the disclosure of bad news. Interns focussed more on obtaining the patients' consent. Both heads of programmes and interns considered the communication of red flags as critical. The survey carried out by Khan *et al.*<sup>58</sup> made it clear that according to interns the current environment

does not leave enough place for the teaching of NTS. For them NTS appears more important than it does for their tutors, both quantitatively and qualitatively. Teachers judge that too much teaching time is spent on NTS, and students think the contrary.

#### *Studies in a simulated environment*

Sakamoto *et al.*<sup>71</sup> carried out an experimental study on 5th-year medical students which showed that expert feedback improved technical performance in surgical anastomosis, without lengthening operating time. Trainees with feedback achieved higher scores than neurosurgeons. Tanweer *et al.*<sup>62</sup> demonstrated through a survey that neurosurgical decision-making is essentially similar from both the patient's and the neurosurgeon's point of view. Morineau *et al.*<sup>64</sup> showed a significant association between expert decision-making and conflict reduction and management of conflict control based on knowledge essentially contributed by experts. Harnof *et al.*<sup>63</sup> in an experimental study of neurosurgical interns in simulated interviews with patients and their families showed the realistic qualities of the actors and their efficiency in NTS communication training. Haji *et al.*<sup>59</sup> showed the importance of simulation in the teaching of teamwork, communication, dealing with pressure, stress and fatigue to first-year neurosurgery interns. Hunt *et al.*<sup>60</sup>, also using simulation, demonstrated that leadership and teamwork are essential for a successful career in neurosurgery. Training camps improved the trainee's awareness of his own behaviour and motivation. Three-quarters of participants understood how to apply this in daily life. Ganju *et al.* specifically studied the effects of fatigue on neurosurgery interns<sup>56</sup>. They subjected 7 neurosurgery interns to 4 sessions of 6 pre- and post-duty surgical exercises in a simulated environment. Fatigue appeared to reduce performance to a certain extent, particularly in "day after duty" interns. Memory and attention were the functions most impaired. Fluidity of gesture and operation duration were also affected by fatigue. Selden *et al.*<sup>54</sup> carried out an opinion poll on 186 neurosurgery interns following practical and theoretical teaching of multiple NTS, in a simulated environment. The teaching of NTS received 100% approval from trainee neurosurgeons, particularly the sections on leadership. Finally, Ciporen *et al.*<sup>57</sup> in an experimental study on interns in neurosurgery and anaesthetics showed no

significant difference in NTS between the two groups. Understanding information and dealing with pressure were important elements. Leadership was more obvious among the neurosurgical interns, but not significantly so. The anaesthetist interns were better at collecting information and considering different options.

*Study during theoretical teaching*

Pettit *et al.*<sup>43</sup> used an opinion poll on theoretical teaching of one hour per month given to 11 neurosurgery interns. Pre-tests and post-tests revealed both quantitative and qualitative improvements in knowledge concerning leadership.

**DISCUSSION**

This systematic review of the literature on NTS is to our knowledge the first which specifically concerns neurosurgery. Analysis of these skills is scattered through the literature, and they are frequently assessed individually without use of an up-to-date unified domain taxonomy. Articles reporting on the assessment of one or several NTS appear very limited in number (21). Specific training needs are difficult to define for the moment since these NTS have not yet been properly formalized. However, the much more abundant literature on the subject in other surgical disciplines offers the possibility of interpolation with the neurosurgical domain.

**Methodological limits of existing studies in neurosurgery**

The methodology of the studies selected after our research did not offer a high standard of proof. Few of them used specific statistical tests. We found no articles using specific surgery NTS assessment methods whose robustness and validity had been demonstrated in the literature. The 3 principal scales currently used for objective assessment of NTS are the OTAS (Observational Teamwork Assessment for Surgery)<sup>72</sup>, the NOTECHS (NonTECHnical Skills)<sup>21</sup> and the NOTSS (Non Technical Skills for Surgeons)<sup>73</sup>. The OTAS<sup>72</sup> essentially assesses the work of the team on the basis of 5 criteria : communication, leadership, cooperation, coordination and surveillance. The NOTECHS<sup>21</sup> classifies NTS in 4 categories : leadership and management, teamwork and cooperation, problem-solving, and decision-making and awareness of the situation. A score of 5 points is given for each category with a scale from 1 (very low) to 5 (very high). The NOTSS corresponds to interviews with consultant surgeons who show five main categories of NTS (awareness of the situation, decision-making, management tasks, leadership, communication and team spirit). Within each category there are 2 or 3 NTS elements depending on the surgical speciality, leading to a score covering 14 separate elements of skill. In some studies all the NTS are assessed together and in others only one or a few of the skills are examined. These factors make it difficult to interpret the results of all these studies. The ideal solution would be an overarching assessment scale with proportionally divided sub-sections to allow all the parameters of the NTS to

be taken into account. Ten of these studies were carried out in a simulated environment or during teaching. The 11 others were done in a simulated environment, in the operating theatre or in consultation. There exist no studies comparing neurosurgery NTS in a simulated environment with a real environment such as the operating theatre. The scope of the evaluation of these NTS proposed here focused on the healthcare activity in real or simulated environment but does not include all the NTS required in this profession.

The methodological heterogeneity of the studies included in our work made it impossible to assess risk of bias in individual studies, summary measures, synthesis of results, risk of bias across studies and additional analysis as suggested in the PRISMA statement.

### **Literature on surgical NTS**

Although there are numerous synonyms in the literature describing surgical NTS, the 4 essential domains most often described are communication, decision-making, leadership and teamwork<sup>19-21,73-80</sup>. There also exists a subdivision of surgical NTS into two major groups : cognitive skills and interpersonal skills<sup>20</sup>.

Several specific skills among the interpersonal NTS have been studied separately, in particular stress<sup>18</sup>. Stress can apparently have both positive and negative effects. A moderate level of stress<sup>24</sup> results in improved performance through increase in concentration. It can however endanger levels of performance when it becomes overwhelming. Pressure gives similar results to stress in a majority of cases. Surprisingly, fatigue (through sleep loss particularly post-duty)<sup>56</sup> can bring about contradictory results. Some studies clearly show a slowdown in task performance, clumsiness and even numbers of supplementary errors, and difficulty in concentrating. Other studies highlight

contrary conclusions, with more rapid procedure time, better gesture ergonomics and fewer errors. An element of disinhibition is probably the reason for this, with fewer conflicts in a fatigue situation, and more pragmatic attitudes. Therefore, assessment in stress or fatigue conditions is very important in order to have objective data on this topic. There is a major issue in the context of the respect of the weekly working hours imposed by the European directives, while at the same time neurosurgery is impaired by an insufficient surgical demography in France.

Interpersonal communication<sup>69</sup> and communication with the patient are among the least studied areas in the literature. Most articles highlight the beneficial effects of feedback from a more experienced surgeon in terms of quality, gesture ergonomics, operation time and numbers of errors. In the rest of the literature NTS are analysed as a whole<sup>17,19-24,29,72,73,75-77,79,81-83</sup>, using scales to examine the different skills. Overall, a close link is shown to exist between teamwork and the number of undesirable events occurring in the operating theatre. These articles underline the necessity for maximum communication among the whole team before, during and after a surgical procedure in order to reduce the risk of errors<sup>17,28,32-35,69,78</sup>.

### **Repercussions and prospects**

The utility of a good knowledge of NTS and of their assessment was demonstrated in the work of Hull *et al* in 2012<sup>19</sup>. NTS appear to have a positive influence on TS : they apparently improve the quality of care and patient outcome by improving safety and efficiency in the operating theatre. This study was carried out on a panel of several different types of specialist surgery.

Good mastery of NTS in neurosurgery has obvious pedagogical implications. A better understanding and definition of these skills in each specialist surgical domain, particularly in neurosurgery, would enable those responsible for teaching to identify more clearly the training

needs for surgeons in that domain. The next step would be inclusion of NTS as specific subjects in educational programmes<sup>25,41,50-52,54,64</sup>. This is beginning to be done in the United States by the Accreditation Council for Graduate Medical Education (ACGME)<sup>53</sup>. The ACGME indicated for each of the specialties the "milestones" to be achieved by residents in training in each of the six "core competencies" required in the training course. Those dedicated to neurosurgery are particularly detailed in the March 2018 version of these recommendations<sup>84</sup>. Including NTS learning in neurosurgery can be envisaged at two points in the training period : during internship (initial training) and/or as ongoing education in the context of continuing professional development (CPD). Confining learning of these skills exclusively to one or another particular stage of the learning process seems unduly simplistic. Placing the acquisition of these skills at the beginning of internship training would be more valuable, since these are general skills which can be applied from the outset of daily surgical activity. They do not necessarily require previous specific declarative knowledge such as complex anatomical elements, which could represent a technical obstacle for a young resident operating, for example, on a brain tumour in a difficult location. It also appears more logical for students to acquire these skills as early as possible to optimize patient safety, for the same reason as they learn basic technical skills (for example, how to hold an instrument properly). It is also essential to include NTS in CPD, since acquisition of these skills is not yet systematic in early training. Surgery, and neurosurgery in particular, is often taught by mentoring, the main teaching tool being the role model. In spite of many hours spent in the operating theatre or with patients, a neurosurgery intern may at times find it difficult to understand the difficulties facing the surgeon and the cognitive conflicts he experiences<sup>64</sup>. On the other hand, the surgeon may have difficulty in transmitting the skills he has learnt by experience, which seem natural to him since he cannot formalize them and make them explicit. Understanding NTS more clearly, formalizing them and rendering them explicit, will greatly improve the training of young neurosurgeons by helping them to identify problems at an early stage, so as to optimize patient safety<sup>85</sup> and accelerate their learning curve.

NTS are also an important subject in the research domain. Initially, applying recognised measuring scales (NOTECHS, NOTSS, OTAS) to neurosurgery could render it possible to pursue the study of their feasibility and assess their reliability and variability in a specific context. Neurosurgery is a complex and specific surgical discipline which nonetheless shares similar difficulties and NTS with other surgical specialties, while retaining its own particular aspects. Studying NTS in neurosurgery opens the possibility of enriching the taxonomy in this domain. The concepts of morbidity and mortality within neurosurgical procedures are of extreme importance. Many operations may carry a risk of life-threatening consequences or of life-changing functional handicaps. They involve cognitive conflicts which are difficult for the surgeon to manage, the more so as the procedure often takes place in the context of an emergency. One example is a patient's recovery from anaesthesia after an operation for cranio-cerebral pathology. Several different scenarios are possible : 1) A patient without preoperative neurological signs may on recovery present new disabilities which may or may not disappear (induced handicap) ; 2) A patient with no preoperative vigilance disorders may not recover consciousness after anaesthesia and risks falling into a chronic coma state ; 3) Fortunately in the majority of cases the patient recovers consciousness without new disabilities, or with an improvement compared to preoperative neurological signs. It can easily be understood that in these different situations stress management, risk assessment and pressure are not the same. Stress management continues until the patient's recovery from anaesthesia ; it does not end when the surgical procedure is finished.

Different methods of intervention to improve safety in operating theatres have been appearing over the last few years, particularly since 2009 with the Surgical Safety Checklist<sup>36</sup>. Other specific checklists have been developed in neurosurgical specialist areas such as vascular neurosurgery or deep brain stimulation<sup>40</sup>. In terms of public health, surgical errors are responsible for a significant increase in morbidity, a deterioration in the quality of life and increased expenditure<sup>10,12-17,28,33,34,36,38,86</sup>. These checklists have been developed<sup>36-40</sup> in order to reduce the number of avoidable errors, with the constant purpose of improving the quality of care and the safety of

patients.

ACCEPTED MANUSCRIPT

**CONCLUSIONS**

After this systematic review of the literature, it appears that very few studies have been carried out concerning neurosurgical NTS, in spite of increasing numbers of articles over the last few years on NTS in other domains of surgery. Moreover, this literature is very diverse in comparison with studies on TS. The complexity of the methodology required to ensure robust assessment tools is probably the reason for this disparity. Society today is concentrating more and more on the quality and safety of medical care, particularly in the domain of neurosurgery. The development and application of these assessment tools is therefore essential. They will also provide important and still underestimated assistance in the training of future neurosurgeons, thus ensuring that all the skills necessary to the profession are effectively transmitted.

**Disclosure**

The authors report no conflict of interest concerning the materials or methods used in this study or the findings specified in this paper.

**Funding sources**

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

**Figure Legends**

**Figure 1** Résumé of structured data for article analysis

**Figure 2** Diagram of analysis of articles

## REFERENCES

1. Alaraj A, Charbel FT, Birk D, Tobin M, Tobin M, Luciano C, et al. Role of cranial and spinal virtual and augmented reality simulation using immersive touch modules in neurosurgical training. *Neurosurgery*. 2013;72 Suppl 1:115–23.
2. Chan S, Conti F, Salisbury K, Blevins NH. Virtual reality simulation in neurosurgery: technologies and evolution. *Neurosurgery*. 2013;72 Suppl 1:154–64.
3. de Ribaupierre S, Wilson TD. Construction of a 3-D anatomical model for teaching temporal lobectomy. *Comput Biol Med*. 2012;42:692–6.
4. Fargen KM, Siddiqui AH, Veznedaroglu E, Turner RD, Ringer AJ, Mocco J. Simulator based angiography education in neurosurgery: results of a pilot educational program. *J Neurointerventional Surg*. 2012;4:438–41.
5. Filho FVG, Coelho G, Cavalheiro S, Lyra M, Zymberg ST. Quality assessment of a new surgical simulator for neuroendoscopic training. *Neurosurg Focus*. 2011;30:E17.
6. Haji FA, Dubrowski A, Drake J, de Ribaupierre S. Needs assessment for simulation training in neuroendoscopy: a Canadian national survey. *J Neurosurg*. 2013;118:250–7.
7. Malone HR, Syed ON, Downes MS, D’Ambrosio AL, Quest DO, Kaiser MG. Simulation in neurosurgery: a review of computer-based simulation environments and their surgical applications. *Neurosurgery*. 2010;67:1105–16.
8. Nathoo N, Pesek T, Barnett GH. Robotics and neurosurgery. *Surg Clin North Am*. 2003;83:1339–50.
9. Ladak A, Spinner RJ. Redefining “wrong site surgery” and refining the surgical pause and checklist: taking surgical safety to another level. *World Neurosurg*. 2014;81:e33-35.
10. Mitchell P, Nicholson CL, Jenkins A. Side errors in neurosurgery. *Acta Neurochir (Wien)*. 2006;148(12):1289–92; discussion 1292.
11. Walcott BP, Redjal N, Coumans J-VCE. Infection following operations on the central nervous system: deconstructing the myth of the sterile field. *Neurosurg Focus*. 2012;33:E8.

12. Wong JM, Ziewacz JE, Panchmatia JR, Bader AM, Pandey AS, Thompson BG, et al. Patterns in neurosurgical adverse events: endovascular neurosurgery. *Neurosurg Focus*. 2012;33:E14.
13. Wong JM, Ziewacz JE, Ho AL, Panchmatia JR, Kim AH, Bader AM, et al. Patterns in neurosurgical adverse events: open cerebrovascular neurosurgery. *Neurosurg Focus*. 2012;33:E15.
14. Wong JM, Ziewacz JE, Ho AL, Panchmatia JR, Bader AM, Garton HJ, et al. Patterns in neurosurgical adverse events: cerebrospinal fluid shunt surgery. *Neurosurg Focus*. 2012;33:E13.
15. Wong JM, Panchmatia JR, Ziewacz JE, Bader AM, Dunn IF, Laws ER, et al. Patterns in neurosurgical adverse events: intracranial neoplasm surgery. *Neurosurg Focus*. 2012;33:E16.
16. Wong JM, Bader AM, Laws ER, Popp AJ, Gawande AA. Patterns in neurosurgical adverse events and proposed strategies for reduction. *Neurosurg Focus*. 2012;33:E1.
17. Ajlan AM, Harsh GR. The human factor and safety attitudes in neurosurgical operating rooms. *World Neurosurg*. 2015;83:46–8.
18. Arora S, Hull L, Sevdalis N, Tierney T, Nestel D, Woloshynowych M, et al. Factors compromising safety in surgery: stressful events in the operating room. *Am J Surg*. 2010;199:60–5.
19. Hull L, Arora S, Aggarwal R, Darzi A, Vincent C, Sevdalis N. The Impact of Nontechnical Skills on Technical Performance in Surgery: A Systematic Review. *J Am Coll Surg*. 2012;214:214–30.
20. Yule S, Flin R, Paterson-Brown S, Maran N. Non-technical skills for surgeons in the operating room: a review of the literature. *Surgery*. 2006;139:140–9.
21. Sharma B, Mishra A, Aggarwal R, Grantcharov TP. Non-technical skills assessment in surgery. *Surg Oncol*. 2011;20:169–77.
22. Arora S, Ahmed M, Paige J, Nestel D, Runnacles J, Hull L, et al. Objective structured assessment of debriefing: bringing science to the art of debriefing in surgery. *Ann Surg*. 2012;256:982–8.
23. Arora S, Miskovic D, Hull L, Moorthy K, Aggarwal R, Johannsson H, et al. Self vs expert

assessment of technical and non-technical skills in high fidelity simulation. *Am J Surg.*

2011;202:500–6.

24. Hull L, Arora S, Kassab E, Kneebone R, Sevdalis N. Assessment of stress and teamwork in the operating room: an exploratory study. *Am J Surg.* 2011;201:24–30.
25. Hull L, Arora S, Symons NRA, Jalil R, Darzi A, Vincent C, et al. Training faculty in nontechnical skill assessment: national guidelines on program requirements. *Ann Surg.* 2013;258:370–5.
26. Mitchell L, Flin R, Yule S, Mitchell J, Coutts K, Youngson G. Evaluation of the Scrub Practitioners' List of Intraoperative Non-Technical Skills system. *Int J Nurs Stud.* 2012;49:201–11.
27. Mitchell L, Flin R, Yule S, Mitchell J, Coutts K, Youngson G. Development of a behavioural marker system for scrub practitioners' non-technical skills (SPLINTS system). *J Eval Clin Pract.* 2013;19:317–23.
28. Russ S, Arora S, Wharton R, Wheelock A, Hull L, Sharma E, et al. Measuring safety and efficiency in the operating room: development and validation of a metric for evaluating task execution in the operating room. *J Am Coll Surg.* 2013;216:472–81.
29. Russ S, Hull L, Rout S, Vincent C, Darzi A, Sevdalis N. Observational teamwork assessment for surgery: feasibility of clinical and nonclinical assessor calibration with short-term training. *Ann Surg.* 2012;255:804–9.
30. Yule S, Flin R, Paterson-Brown S, Maran N, Rowley D. Development of a rating system for surgeons' non-technical skills. *Med Educ.* 2006;40:1098–104.
31. Fallah A, Ibrahim GM, Bernstein M. The “SAAFE” neurosurgical sign-out. *World Neurosurg.* 2014;81:e21-23.
32. Hull L, Arora S, Amaya AC, Wheelock A, Gaitán-Duarte H, Vincent C, et al. Building global capacity for patient safety: a training program for surgical safety research in developing and transitional countries. *Int J Surg Lond Engl.* 2012;10:493–9.
33. Lau CY, Greysen SR, Mistry RI, Han SJ, Mummaneni PV, Berger MS. Creating a culture of

safety within operative neurosurgery: the design and implementation of a perioperative safety

video. *Neurosurg Focus*. 2012;33:E3.

34. Muzumdar D. Safety in the operating room: neurosurgical perspective. *Int J Surg Lond Engl*. 2007;5:286–8.

35. Ferroli P, Caldiroli D, Acerbi F, Scholtze M, Piro A, Schiariti M, et al. Application of an aviation model of incident reporting and investigation to the neurosurgical scenario: method and preliminary data. *Neurosurg Focus*. 2012;33:E7.

36. Haynes AB, Weiser TG, Berry WR, Lipsitz SR, Breizat A-HS, Dellinger EP, et al. A Surgical Safety Checklist to Reduce Morbidity and Mortality in a Global Population. *N Engl J Med*. 2009;360:491–9.

37. Da Silva-Freitas R, Martín-Laez R, Madrazo-Leal CB, Villena-Martin M, Valduvico-Juaristi I, Martínez-Agüeros JÁ, et al. Establishment of a modified surgical safety checklist for the neurosurgical patient: Initial experience in 400 cases. *Neurocir Astur Spain*. 2012;23:60–9.

38. McConnell DJ, Fargen KM, Mocco J. Surgical checklists: A detailed review of their emergence, development, and relevance to neurosurgical practice. *Surg Neurol Int*. 2012;3:2.

39. Rahmathulla G, Recinos PF, Traul DE, Avitsian R, Yunak M, Harper NT, et al. Surgical briefings, checklists, and the creation of an environment of safety in the neurosurgical intraoperative magnetic resonance imaging suite. *Neurosurg Focus*. 2012;33:E12.

40. Zuckerman SL, Green CS, Carr KR, Dewan MC, Morone PJ, Mocco J. Neurosurgical checklists: a review. *Neurosurg Focus*. 2012;33:E2.

41. Long DM. Competency-based residency training: the next advance in graduate medical education. *Acad Med J Assoc Am Med Coll*. 2000;75:1178–83.

42. Flin R, Yule S, Paterson-Brown S, Maran N, Rowley D, Youngson G. Teaching surgeons about non-technical skills. *The Surgeon*. 2007;5:86–9.

43. Pettit JE, Dahdaleh NS, Albert GW, Greenlee JD. Neurosurgery resident leadership development: an innovative approach. *Neurosurgery*. 2011;68:546–50; discussion 550.

44. Cusimano MD, Yonke AM, Tucker WS. An analysis of attrition from Canadian neurosurgery residency programs. *Acad Med J Assoc Am Med Coll.* 1999;74:925–31.
45. Dedy NJ, Bonrath EM, Zevin B, Grantcharov TP. Teaching nontechnical skills in surgical residency: a systematic review of current approaches and outcomes. *Surgery.* 2013;154:1000–8.
46. Dumont TM, Horgan MA. The surgical skills laboratory residency interview: an enjoyable alternative. *J Surg Educ.* 2012;69:407–10.
47. Fallah A, Ebrahim S, Haji F, Gillis C, Girgis F, Howe K, et al. Surgical activity of first-year canadian neurosurgical residents. *Can J Neurol Sci.* 2010;37:855–60.
48. Fernando N, McAdam T, Cleland J, Yule S, McKenzie H, Youngson G. How can we prepare medical students for theatre-based learning? *Med Educ.* 2007;41:968–74.
49. Haase J, Boisen E. Neurosurgical training: more hours needed or a new learning culture? *Surg Neurol.* 2009;72(1):89–95.
50. Heros RC. Neurosurgical education: the “other” competencies. The 2003 presidential address. *J Neurosurg.* 2003;99:623–9.
51. Kabwama S. Opportunities to improve undergraduate neurosurgery career interest, knowledge and skills: lessons from the BAPRAS undergraduate courses. *Br J Neurosurg.* 2011;25:783–4.
52. Kaufman HH, Wiegand RL, Tunick RH. Teaching surgeons to operate--principles of psychomotor skills training. *Acta Neurochir (Wien).* 1987;87:1–7.
53. Mazzola CA, Lobel DA, Krishnamurthy S, Bloomgarden GM, Benzil DL. Efficacy of neurosurgery resident education in the new millennium: the 2008 Council of State Neurosurgical Societies post-residency survey results. *Neurosurgery.* 2010;67:225–32; discussion 232-233.
54. Selden NR, Origitano TC, Burchiel KJ, Getch CC, Anderson VC, McCartney S, et al. A National Fundamentals Curriculum for Neurosurgery PGY1 Residents: The 2010 Society of Neurological Surgeons Boot Camp Courses. *Neurosurgery.* 2012;70:971–81.
55. Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gøtzsche PC, Ioannidis JPA, et al. The

PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate

healthcare interventions: explanation and elaboration. *BMJ*. 2009;339:b2700.

56. Ganju A, Kahol K, Lee P, Simonian N, Quinn SJ, Ferrara JJ, et al. The effect of call on neurosurgery residents' skills: implications for policy regarding resident call periods. *J Neurosurg*. 2011;116:478–82.
57. Ciporen J, Gillham H, Noles M, Dillman D, Baskerville M, Haley C, et al. Crisis Management Simulation: Establishing a Dual Neurosurgery and Anesthesia Training Experience. *J Neurosurg Anesthesiol*. 2018;30:65–70.
58. Khan A, Doke T, Boeris D. Nurturing the Next Generation of Neurosurgeons: How Important Are Nontechnical Skills? *World Neurosurg*. 2018;120:e227–33.
59. Haji FA, Clarke DB, Matte MC, Brandman DM, Brien S, Ribaupierre S de, et al. Teaching for the Transition: the Canadian PGY-1 Neurosurgery 'Rookie Camp.' *Can J Neurol Sci*. 2015;42:25–33.
60. Hunt MA, Heilman CB, Shutrán M, Wu JK. Commentary: An Introduction to Leadership Self-Assessment at the Society of Neurological Surgeons Post-Graduate Year 1 Boot Camp: Observations and Commentary. *Neurosurgery*. 2017;80:E201–4.
61. Brandman DM, Haji FA, Matte MC, Clarke DB. Needs Assessment for Incoming PGY-1 Residents in Neurosurgical Residency. *Can J Neurol Sci J Can Sci Neurol*. 2015;42:17–24.
62. Tanweer O, Wilson TA, Kalhorn SP, Golfinos JG, Huang PP, Kondziolka D. Neurosurgical decision making: personal and professional preferences. *J Neurosurg*. 2015;122:678–91.
63. Harnof S, Hadani M, Ziv A, Berkenstadt H. Simulation-based interpersonal communication skills training for neurosurgical residents. *Isr Med Assoc J IMAJ*. 2013;15:489–92.
64. Morineau T, Morandi X, Moëllic NL, Diabira S, Riffaud L, Haegelen C, et al. Decision Making During Preoperative Surgical Planning. *Hum Factors J Hum Factors Ergon Soc*. 2009;51:67–77.
65. Oszvald Á, Vatter H, Byhahn C, Seifert V, Güresir E. "Team time-out" and surgical safety-

experiences in 12,390 neurosurgical patients. *Neurosurg Focus*. 2012;33:E6.

66. McLaughlin N, Winograd D, Chung HR, Van de Wiele B, Martin NA. Impact of the Time-Out Process on Safety Attitude in a Tertiary Neurosurgical Department. *World Neurosurg*. 2014;82:567–74.
67. Pathak A, Pathak N, Kak VK. Ideal ward round making in neurosurgical practice. *Neurol India*. 2000;48:216–22.
68. Couat J-F, Cegarra J, Rodsphon T, Geeraerts T, Lelardeux C, Sol J-C, et al. A prospective video-based observational and analytical approach to evaluate management during brain tumour surgery at a university hospital. *Neurochirurgie*. 2013;59:142–8.
69. Fargen KM, Velat GJ, Lawson MF, Firment CS, Mocco J, Hoh BL. Enhanced staff communication and reduced near-miss errors with a neurointerventional procedural checklist. *J Neurointerventional Surg*. 2013;5:497–500.
70. Michinov E, Jamet E, Dodeler V, Haegelen C, Jannin P. Assessing neurosurgical non-technical skills: an exploratory study of a new behavioural marker system. *J Eval Clin Pract*. 2014;20:582–8.
71. Sakamoto Y, Okamoto S, Shimizu K, Araki Y, Hirakawa A, Wakabayashi T. Comparative Prospective Study of Microvascular Anastomosis Training by Self-Learning or with Expert Instruction. *World Neurosurg*. 2018;118:e818–24.
72. Hull L, Arora S, Kassab E, Kneebone R, Sevdalis N. Observational teamwork assessment for surgery: content validation and tool refinement. *J Am Coll Surg*. 2011;212:234-243.e1-5.
73. Yule S, Flin R, Maran N, Rowley D, Youngson G, Paterson-Brown S. Surgeons' Non-technical Skills in the Operating Room: Reliability Testing of the NOTSS Behavior Rating System. *World J Surg*. 2008;32:548–56.
74. Flin R, Youngson G, Yule S. How do surgeons make intraoperative decisions? *Qual Saf Health Care*. 2007;16:235–9.
75. Henrickson Parker S, Flin R, McKinley A, Yule S. The Surgeons' Leadership Inventory

(SLI): a taxonomy and rating system for surgeons' intraoperative leadership skills. *Am J Surg*.

2013;205:745–51.

76. Neil-Dwyer G, Varma TR, Lang DA. Mechanisms to improve teamworking in neurosurgery. *Acta Neurochir Suppl*. 2001;78:135–8.
77. Pauley K, Flin R, Yule S, Youngson G. Surgeons' intraoperative decision making and risk management. *Am J Surg*. 2011;202:375–81.
78. Sekhar LN, Mantovani A. Teamwork mentality in neurosurgical teams to improve patient safety. *World Neurosurg*. 2015;83:41–3.
79. Yule S, Paterson-Brown S. Surgeons' non-technical skills. *Surg Clin North Am*. 2012;92:37–50.
80. Zuckerman SL, France DJ, Green C, Leming-Lee S, Anders S, Mocco J. Surgical debriefing: a reliable roadmap to completing the patient safety cycle. *Neurosurg Focus*. 2012;33:E4.
81. Yule S, Flin R, Paterson-Brown S, Maran N, Rowley D. Development of a rating system for surgeons' non-technical skills. *Med Educ*. 2006;40:1098–104.
82. Henrickson Parker S, Yule S, Flin R, McKinley A. Towards a model of surgeons' leadership in the operating room. *BMJ Qual Saf*. 2011;20:570–9.
83. Parker SH, Yule S, Flin R, McKinley A. Surgeons' leadership in the operating room: an observational study. *Am J Surg*. 2012;204:347–54.
84. NeurologicalSurgeryMilestones.pdf [Internet]. [cited 2019 Mar 27]. Available from: <https://www.acgme.org/Portals/0/PDFs/Milestones/NeurologicalSurgeryMilestones.pdf?ver=2019-02-28-093643-277>
85. Lane-Fall MB, Davis JJ, Clapp JT, Myers JS, Riesenber LA. What Every Graduating Resident Needs to Know About Quality Improvement and Patient Safety: A Content Analysis of 26 Sets of ACGME Milestones. *Acad Med J Assoc Am Med Coll*. 2018;93:904–10.
86. Marcus H, Vakharia V, Kirkman MA, Murphy M, Nandi D. Practice Makes Perfect? The Role of Simulation-Based Deliberate Practice and Script-Based Mental Rehearsal in the Acquisition

ACCEPTED MANUSCRIPT

**Table 1.** Characteristics of the studies assessing non-technical skills in neurosurgery

First author	Year of publication	Population	Type of study	Environment	Non-technical skills assessed	Tools for assessment	Key results
Cusimano <sup>44</sup>	1999	Teaching hospital neurosurgeons and interns in neurosurgery	Qualitative	Real	Interpersonal skills and workload	Qualitative analysis of questionnaires Double scale content analysis	Positive correlation between excess workload and voluntary withdrawal and between dismissal and interpersonal skills during the neurosurgery internship
Pathak <sup>67</sup>	2000	Professors of neurosurgery	Interventional study	Real environment and consultations	Leadership	External observation scale for patterns of leadership and language	Good productivity and flexibility, moderate objectivity and comprehensibility
Morineau <sup>64</sup>	2009	Neurosurgery interns, registrars and professors	Experimental study	Simulation	Decision-making	Number of conflicts and checks reported by each neurosurgeon Skills /Rule / Knowledge approach	Significant link between the expert undertaking tasks and checking of conflicts, Regulation of conflict control essentially based on experts' knowledge
Pettit <sup>43</sup>	2011	Neurosurgery interns	Survey	Theoretical teaching	Leadership	Opinion poll	Training in the importance of leadership and the means to develop it.
Ganju <sup>56</sup>	2012	Neurosurgery inters	Semi-experimental study	Simulation	Dealing with fatigue, cognitive skills	Fluidity of gestures, time taken and cognitive errors	No significant reduction in skills during simulated surgical procedures after on-call service.
Selden <sup>54</sup>	2012	Professors of neurosurgery and first-year neurosurgery interns	Survey	Simulation	Multiple non-technical skills	Opinion poll	Teaching appreciated by interns for the improvement of their knowledge and their technical and non-technical skills .

Ferrolì <sup>35</sup>	2012	Neurosurgery operating theatre team	Interventional study	Real environment – operating theatre	Multiple non-technical skills	Incidence of undesirable events and analysis of their causes by methods originating in aviation	The majority of incidents were linked to human or organisational factors
Lau <sup>33</sup>	2012	Duty neurosurgery team	Interventional study	Real environment	Communication and teamwork	Analysis of perioperative safety video	The perioperative safety video improves safety in the operating theatre and raises the team's awareness of safety issues
Oszvald <sup>65</sup>	2012	Neurosurgery operating theatre team	Interventional study	Real event – operating theatre	Communication, teamwork preparation, risk assessment, anticipation of problems	Number of undesirable events	The surgical safety checklist improves communication within the team and reduces the number of undesirable events
Harnof <sup>63</sup>	2013	Neurosurgery interns	Experimental study	Simulation – interviews with patients and their families	Communication	Feedback questionnaires (Likert scale)	Realistic actors, efficient training in interpersonal communication skills
Couat <sup>68</sup>	2013	Neurosurgery operating theatre team	Observational study	Real environment - operating theatre	Communication, resource management	Analysis of videos and direct observation using ethnographical methods (notes taken from observation on site)	33% of errors in the operating theatre are at least partly due to poor communication or organisational management
Fargen <sup>69</sup>	2013	Neurosurgery operating theatre team	Interventional study	Real environment - operating theatre	Communication, teamwork preparation, risk assessment, anticipation of problems	Communication and number of undesirable events	The surgical safety checklist improves communication within the team and reduces the number of undesirable events
McLaughlin <sup>66</sup>	2013	Neurosurgery operating theatre team	Survey	Real environment - operating	Communication, teamwork preparation, risk	Opinion poll	The surgical safety checklist improves communication within the team and reduces the number of undesirable events

				theatre	assessment, anticipation of problems		
Michinov <sup>70</sup>	2014	Neurosurgery operating theatre team and patient	Observational study	Real environment - operating theatre	Cooperation and teamwork, awareness of the situation, explicit coordination, decision-making and leadership	Behavioural marker system for assessing neurosurgeons' non-technical skills (BMS - NNTS)	Explicit coordination, awareness of the situation and leadership are the non-technical skills most implicated in deep brain stimulation surgery by the neurosurgeon. Moreover these behaviours vary with the time of the operation and the other member of the surgical team with whom he is communicating.
Tanweer <sup>62</sup>	2015	Neurosurgery interns, registrars and professors	Survey	Simulation	Decision-making	Questionnaire	In neurosurgical decision-making the point of view of the patient and of the neurosurgeon are essentially equivalent.
Brandman <sup>61</sup>	2015		Survey	Real environment - common practice	All non-technical skills	Online questionnaire, Likert scale, open-ended questions	Programme directors placed non-technical communication skills at a higher level than interns, with emphasis on announcing bad news. Interns focussed more on obtaining patients' consent. Both programme directors and interns considered red-flag communication as critically important
Haji <sup>59</sup>	2015	First-year neurosurgery interns	Pre- and post-test model	Simulation	Teamwork, communication, dealing with pressure, stress and fatigue	Multiple-choice questionnaire	Significant differences between pre- and post-test but not on the retention test
Hunt <sup>60</sup>		Neurosurgery interns	Pre- and post-test model	Simulation	Leadership	Questionnaire and personality assessment (Hogan Personality Inventory)	Leadership and teamwork are essential for success in a neurosurgeon's career, This training session improves interns'

							awareness of their own behaviour and motivation. ¾ of trainees understood how to apply this in daily practice.
Khan <sup>58</sup>	2018	Neurosurgery interns and teaching neurosurgeons	Survey	Real environment : teaching/ curriculum	All non-technical skills	Questionnaire with quantitative and qualitative analysis	Interns feel that the current environment leaves little place for the acquisition of non-technical skills. They give greater importance to the learning of non-technical skills than their teachers do, both quantitatively and qualitatively. Teachers feel that too much time is given to non-technical skills, and the trainees believe the contrary.
Sakamoto <sup>71</sup>	2018	5th-year medical students and neurosurgeons	Experimental study	Simulation	Feedback	Comparison of a surgical act of vascular anastomosis with and without feedback	Expert feedback improves the technical performance of anastomosis without prolonging operation time. Trainees with feedback had higher scores than the neurosurgeons.
Ciporen <sup>57</sup>	2018	Neurosurgery and anaesthetic interns	Experimental study	Simulation	Awareness of the situation, decision-making, communication, teamwork, leadership, dealing with pressure	Non-technical skills observation scale	No significant difference in non-technical skills between interns in neurosurgery and anaesthetics. Understanding information and dealing with pressure were important. Leadership was more important for neurosurgery interns, but not significantly. Anaesthetics interns performed better in collecting information and considering different options,

**Name of first author**

**Year of publication**

**Population**

**Study design**

**Environment**

**Non-technical skills assessed**

*Interpersonal skills*

Communication

Leadership

Teamwork

Briefing/Planning/Preparation

Resource management

Seeking advice and feedback

Coping with pressure/stress/fatigue

*Cognitive skills*

Situation awareness

Mental readiness

Risk assessment

Anticipation of problems

Decision-making

Adaptation strategies and flexibility

Workload distribution

**Assessment tools for NTS**

NOTSS|

NOTECHS

OTAS

Others

**Key results**





# PRISMA 2009 Checklist

Section/topic	#	Checklist item	Reported on page #
<b>TITLE</b>			
Title	1	Identify the report as a systematic review, meta-analysis, or both.	1
<b>ABSTRACT</b>			
Structured summary	2	Provide a structured summary including, as applicable: background; objectives; data sources; study eligibility criteria, participants, and interventions; study appraisal and synthesis methods; results; limitations; conclusions and implications of key findings; systematic review registration number.	1
<b>INTRODUCTION</b>			
Rationale	3	Describe the rationale for the review in the context of what is already known.	5
Objectives	4	Provide an explicit statement of questions being addressed with reference to participants, interventions, comparisons, outcomes, and study design (PICOS).	5
<b>METHODS</b>			
Protocol and registration	5	Indicate if a review protocol exists, if and where it can be accessed (e.g., Web address), and, if available, provide registration information including registration number.	7
Eligibility criteria	6	Specify study characteristics (e.g., PICOS, length of follow-up) and report characteristics (e.g., years considered, language, publication status) used as criteria for eligibility, giving rationale.	7
Information sources	7	Describe all information sources (e.g., databases with dates of coverage, contact with study authors to identify additional studies) in the search and date last searched.	7
Search	8	Present full electronic search strategy for at least one database, including any limits used, such that it could be repeated.	7
Study selection	9	State the process for selecting studies (i.e., screening, eligibility, included in systematic review, and, if applicable, included in the meta-analysis).	7
Data collection process	10	Describe method of data extraction from reports (e.g., piloted forms, independently, in duplicate) and any processes for obtaining and confirming data from investigators.	7
Data items	11	List and define all variables for which data were sought (e.g., PICOS, funding sources) and any assumptions and simplifications made.	8
Risk of bias in individual studies	12	Describe methods used for assessing risk of bias of individual studies (including specification of whether this was done at the study or outcome level), and how this information is to be used in any data synthesis.	NC
Summary measures	13	State the principal summary measures (e.g., risk ratio, difference in means).	NC
Synthesis of results	14	Describe the methods of handling data and combining results of studies, if done, including measures of consistency (e.g., $I^2$ ) for each meta-analysis.	NC



## PRISMA 2009 Checklist

Section/topic	#	Checklist item	Reported on page #
Risk of bias across studies	15	Specify any assessment of risk of bias that may affect the cumulative evidence (e.g., publication bias, selective reporting within studies).	NC
Additional analyses	16	Describe methods of additional analyses (e.g., sensitivity or subgroup analyses, meta-regression), if done, indicating which were pre-specified.	NC
<b>RESULTS</b>			
Study selection	17	Give numbers of studies screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally with a flow diagram.	9
Study characteristics	18	For each study, present characteristics for which data were extracted (e.g., study size, PICOS, follow-up period) and provide the citations.	10-13
Risk of bias within studies	19	Present data on risk of bias of each study and, if available, any outcome level assessment (see item 12).	NC
Results of individual studies	20	For all outcomes considered (benefits or harms), present, for each study: (a) simple summary data for each intervention group (b) effect estimates and confidence intervals, ideally with a forest plot.	10-13
Synthesis of results	21	Present results of each meta-analysis done, including confidence intervals and measures of consistency.	NC
Risk of bias across studies	22	Present results of any assessment of risk of bias across studies (see Item 15).	NC
Additional analysis	23	Give results of additional analyses, if done (e.g., sensitivity or subgroup analyses, meta-regression [see Item 16]).	NC
<b>DISCUSSION</b>			
Summary of evidence	24	Summarize the main findings including the strength of evidence for each main outcome; consider their relevance to key groups (e.g., healthcare providers, users, and policy makers).	114
Limitations	25	Discuss limitations at study and outcome level (e.g., risk of bias), and at review-level (e.g., incomplete retrieval of identified research, reporting bias).	14-15
Conclusions	26	Provide a general interpretation of the results in the context of other evidence, and implications for future research.	20
<b>FUNDING</b>			
Funding	27	Describe sources of funding for the systematic review and other support (e.g., supply of data); role of funders for the systematic review.	21

From: Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 6(7): e1000097. doi:10.1371/journal.pmed1000097

For more information, visit: [www.prisma-statement.org](http://www.prisma-statement.org).

NC: non considered

**Disclosure-Conflict of Interest**

The authors report no conflict of interest concerning the materials or methods used in this study or the findings specified in this paper.

ACCEPTED MANUSCRIPT