



Association between early lead exposure and externalizing behaviors in adolescence: A developmental cascade

Mireille Desrochers-Couture, Yohann Courtemanche, Nadine Forget-Dubois, Richard E Bélanger, Olivier Boucher, Pierre Ayotte, Sylvaine Cordier, Joseph L Jacobson, Sandra W Jacobson, Gina Muckle

► To cite this version:

Mireille Desrochers-Couture, Yohann Courtemanche, Nadine Forget-Dubois, Richard E Bélanger, Olivier Boucher, et al.. Association between early lead exposure and externalizing behaviors in adolescence: A developmental cascade. *Environmental Research*, 2019, 178, pp.108679. 10.1016/j.envres.2019.108679 . hal-02280778

HAL Id: hal-02280778

<https://univ-rennes.hal.science/hal-02280778>

Submitted on 17 Feb 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1
2
3
4
5 1 **Association between early lead exposure and externalizing behaviors in adolescence:**
6 2 **a developmental cascade**
7
8 3
9 4 Mireille Desrochers-Couture^{1,a}, Yohann Courtemanche^{1,a}, Nadine Forget-Dubois¹,
10 5 Richard E. Bélanger^{1,2}, Olivier Boucher^{3,4}, Pierre Ayotte^{1,5,6}, Sylvaine Cordier^{1,7}, Joseph
11 6 L. Jacobson⁸, Sandra W. Jacobson⁸, Gina Muckle^{*1,9}
12
13
14 7
15
16 8 ¹Centre de recherche du CHU de Québec-Université Laval, Québec, Québec, Canada
17 9 ²Département de pédiatrie, Université Laval, Québec, Québec, Canada
18 10 ³Département de psychologie, Université de Montréal, Montréal, Québec, Canada
19 11 ⁴Service de psychologie, Centre hospitalier de l'Université de Montréal, Montréal,
20 12 Québec, Canada
21 13 ⁵Centre de toxicologie du Québec, Institut national de santé publique du Québec, Québec,
22 14 Québec, Canada
23 15 ⁶Département de médecine sociale et préventive, Université Laval, Québec, Québec,
24 16 Canada
25 17 ⁷Univ Rennes, Inserm, EHESP, Irset (Institut de recherche en santé, environnement et
26 18 travail) - UMR_S1085, F-35000 Rennes, France
27 19 ⁸Department of Psychiatry and Behavioral Neurosciences, Wayne State University
28 20 School of Medicine, Detroit, USA
29 21 ⁹École de psychologie, Université Laval, Québec, Québec, Canada
30 22
31 23 * Corresponding author: Gina Muckle (gina.muckle@psy.ulaval.ca), École de
32 24 psychologie, Université Laval, Pavillon Félix-Antoine-Savard, 2325 Rue des
33 25 Bibliothèques, Québec, G1V 0A6, Canada.
34 26
35 27 ^aEqual contributors.
36 28
37 29 The authors declare they have no actual or potential competing financial interests.
38 30
39 31
40 32 Keywords: lead; children and adolescents; externalizing behaviors; binge drinking;
41 33 cannabis use.
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

34 **Abstract**

35 **Background:** Lead (Pb) exposure is associated with adverse neurological development.
36 Most notably, it has been observed through externalizing behavior symptoms, as
37 observed among Inuit children from northern Québec. Evidence for a persistent
38 neurological impact of early Pb exposure later in life is however scarce. Pb exposure may
39 initiate a developmental cascade that increases the risk of long-term behavior problems.

40 **Objectives:** Testing for direct associations between childhood Pb concentrations and
41 adolescent externalizing symptoms and substance use, as well as indirect associations
42 through childhood behavior assessments.

43 **Methods:** The study sample is a longitudinal cohort of Inuit children (n=212) followed
44 since birth. Blood Pb concentrations were measured during childhood (median age = 11.4
45 years) and adolescence (median age=18.5 years). Externalizing/inattentive behavior were
46 teacher-assessed through the Teacher Report Form and the Disruptive Behavior Disorders
47 Rating Scale for children. At the adolescence follow-up, behavior problems were self-
48 reported by filling Achenbach's Youth Self-Report, the Barkley Adult ADHD-IV Rating
49 Scale, and the Diagnostics Interview Schedule for Children. Adolescent substance use
50 was also self-assessed through the DEP-ADO. Direct and indirect associations of child
51 Pb concentrations with adolescent outcomes were tested through mediation models.

52 **Results:** Child blood Pb concentrations were not directly associated with any adolescent
53 outcomes. On the contrary, childhood Pb exposure was indirectly associated, through
54 childhood externalizing behavior assessments, with adolescent externalizing behaviors,
55 binge drinking, and cannabis use. These indirect associations held after controlling for
56 adolescents' concurrent Pb blood concentrations.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

57 **Discussion:** Our results highlight the indirect but lasting effects of child Pb exposure on
58 adolescent behavior problems, and the importance of childhood externalizing behavior in
59 this relationship. Adverse early-life environment put children on a riskier developmental
60 trajectory, increasing their likelihood of lifelong psychological, social and health
61 problems.

62

Introduction

Lead (Pb) exposure in children has been associated with negative neurological outcomes for more than a century (Turner, 1897). In Western countries, policies implemented since the 1970s were largely successful in reducing Pb exposure. However, certain groups and subpopulations remain highly exposed to Pb through diverse human-linked sources such as aged water pipes, dust and paint chips in older neighborhoods, or Pb-based ammunitions.

Pb exposure in children is of particular concern considering its known influence on developing neurological structures, which may lead to long-lasting cognitive and behavioral impairments (Téllez-Rojo et al., 2006). Several studies conducted in a variety of populations, such as major inner cities citizen (Chiodo et al., 2007, 2004; Needleman et al., 2002) and representative samples in the US (Braun et al., 2006; Froehlich et al., 2009), South Africa (Naicker et al., 2012), and Inuit children from northern Quebec (Boucher et al., 2012c), have reported cross-sectional associations between Pb exposure and numerous externalizing behavior or Attention Deficit Hyperactivity Disorder (ADHD) symptoms and diagnoses in childhood and adolescence. However, only a handful of longitudinal studies have documented these associations later in adolescence or early adulthood (Beckley et al., 2018; Burns et al., 1999; Dietrich et al., 2001; Winter and Sampson, 2017; Wright et al., 2008). Consequences of early Pb exposure throughout the lifespan still remain largely unknown, in part because of complex interactions with other determinants of neurodevelopment.

The long-term effects of early Pb exposure could be explained as a perturbation of early neurodevelopment that changes the context of later development (Bellinger et al., 2016). Pb exposure during childhood may not directly cause later behavioral and cognitive problems; it could instead contribute to putting exposed children on an adverse developmental path. For this reason, Bellinger et al. (2016) predict that although Pb exposure is directly associated with childhood behavioral and cognitive outcomes, its association with adulthood outcomes could be indirect. Thus, instead of a direct association between Pb exposure during childhood and later problem behaviors, we could expect a process mediated by the continuation of behavior problems during childhood

1
2
3
4 93 into adolescence, corresponding to the concept of developmental cascade. In this
5
6 94 framework, early Pb exposure would play a role in fostering and maintaining behavior
7
8 95 problems from childhood to early adulthood, highlighting the importance of early-life
9
10 96 environment.

11
12 97 In order to incorporate a developmental perspective into the study of effects of
13
14 98 early neurotoxicant exposure, one should take into account the evolution of
15
16 99 manifestations of neurodevelopmental perturbations throughout development. In
17
18 100 adolescence and adulthood, externalizing behaviors manifest themselves as a spectrum of
19
20 101 mental disorders and behaviors, such as substance use and abuse (Krueger et al., 2005),
21
22 102 that children with behavior problems are more likely to develop later in life. A meta-
23
24 103 analysis of longitudinal studies reported that adolescents and young adults who were
25
26 104 diagnosed with ADHD as children were at an increased risk for alcohol use disorder and
27
28 105 cannabis use (Lee et al., 2011). Childhood oppositional defiant/conduct disorder has also
29
30 106 been prospectively associated with later substance use (Groenman et al., 2017). The only
31
32 107 exploratory study, with a limited sample size, investigating the association between
33
34 108 substance use and long-term Pb exposure did not find any significant association
35
36 109 (Fishbein et al., 2008). The mechanisms linking childhood externalizing behaviors with
37
38 110 adolescent or early adulthood substance use remain unclear. One potential mechanism
39
40 111 explaining the increased risk of later substance use may be a vulnerability to disinhibitory
41
42 112 behavior and impulse control (McGue et al., 2001). Accordingly, in this same cohort, our
43
44 113 team has previously published significant results showing an association between child
45
46 114 blood Pb concentrations and impairments in the child's ability to correctly inhibit a
47
48 115 response, resulting in increased impulsivity (Boucher et al., 2012a).

49
50 116 In a previous study by our group (Boucher et al., 2012b), we reported a cross-
51
52 117 sectional association between Pb exposure and externalizing symptoms in a sample of
53
54 118 Inuit children with a median age of 11.4 years. The goal of the present study is to expand
55
56 119 these results, which includes a follow-up of the same participants into adolescence. The
57
58 120 Inuit communities in northern Québec practice traditional activities central to their health
59
60 121 and well-being, but some of the practices expose them to multiple environmental
61
62 122 contaminants. The ingestion of Pb ammunition fragments in game meat is their primary
63
64
65

1
2
3
4 123 source of Pb exposure (Lévesque et al., 2003). Alongside symptoms of externalizing and
5
6 124 attention problems in adolescence, this study's outcomes include substance use, a
7
8 125 manifestation of externalizing problems in adolescence and adulthood that represent
9
10 126 major public health concern in many indigenous communities across Canada (Firestone
11
12 127 et al., 2015). Therefore, this paper will investigate two processes potentially leading to
13
14 128 the association between child Pb exposure and adolescent externalizing
15
16 129 behaviors/substance use: 1) direct association between child Pb concentrations and
17
18 130 adolescent outcomes and 2) indirect association in which childhood behavior problems
19
20 131 mediate the link between child Pb concentrations and adolescent outcomes, as expected
21
22 132 in a developmental cascade framework.
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Methods

Participants

The participants were Inuit children from the 14 coastal villages of Nunavik, in northern Québec. Between November 1993 and December 1996, 491 mothers were recruited as part of the Cord Blood Monitoring Program designed to document prenatal exposure to environmental contaminants and a range of nutrients in Arctic Québec (Dewailly et al., 1993). An additional 221 mothers were recruited between November 1995 and March 2002 as part of the National Institutes of Health (NIH) prospective infancy study (Jacobson et al., 2008; Muckle et al., 2001). At school age, a subsample of these children and their primary caregiver ($n=294$: 247 from the Cord Blood Monitoring Program and 47 from the NIH-infancy study) participated in the Nunavik Child Development Study (NCDS-childhood), a follow-up designed to examine effects of pre- and postnatal exposure to environmental contaminants on child behavior and cognitive abilities, which took place between September 2005 and February 2010. Recruitment methodologies for both the Cord Blood Monitoring Program and NCDS-childhood samples were reported in previous studies based on this cohort (Dallaire et al. 2014; Dewailly et al. 1993; Jacobson et al. 2008). The participants were met again during adolescence (NCDS-adolescence) between January 2013 and February 2016. Inclusion criteria for the NCDS-adolescence follow-up were participation in the previous two follow-ups, living in Nunavik, and ability to meet with the research team in one of Nunavik's three main villages. Those who were identified as suffering from severe health or neurological problems unrelated to exposure at the NCDS-childhood interview (epilepsy $n=2$; head trauma $n=1$; meningitis $n=1$; multiple sclerosis $n=1$) were excluded from the NCDS-adolescence follow-up. An additional 49 children of the NCDS-childhood study were not eligible for the follow-up because they were either deceased, incarcerated, had moved away or were unreachable. An additional 28 adolescents declined to participate in the follow-up. Thus, a remaining 212 adolescents participated in the NCDS-adolescence study (Fig. 1). As compensation, the adolescent participants received an electronic device worth \$50 USD.




Figure 1. Flow chart for recruitment and follow-up of study participants from November 1993 to February 2016 including reasons and number of excluded participants through follow-ups. Note: NIH, National Institutes of Health; NCDS, Nunavik Child Development Study.

The interviews and cognitive/behavioral assessments for the child and adolescent follow-ups were conducted in the three largest villages of the region. Participants from smaller communities were transported by plane to meet with the research team. Written informed consent was provided by the biological mother at recruitment, the primary caregiver at the child follow-up, and the participants themselves at the adolescent follow-up. The children also gave verbal consent at the childhood follow-up. The Université

Laval and Wayne State University ethics committees approved the Infancy and NCDS-childhood follow-ups, and the ethics committee of the Centre de recherche du CHU de Québec-Université Laval approved the consent and study procedures at the adolescence.

Biological samples

Contaminants and nutrients, including mercury (Hg), Pb and, docosahexaenoic acid (DHA, an omega-3 fatty acid) were measured in umbilical cord blood samples at birth (30 mL) and in venous blood samples of children (20 mL) and adolescents (30 mL). Analyses were performed at the Centre de Toxicologie, Institut National de Santé Publique du Québec (Québec, Canada) for all cord, child, and adolescent contaminant concentrations. Cord and child omega-3 fatty acid composition of plasma phospholipids were analyzed at the University of Guelph Lipid Analytical Laboratory (Guelph, Canada) and at the CHU de Québec for adolescent blood (see Supplemental Material for details of blood samples analytical procedures).

Child behavior assessment

Child externalizing. The Child Behavior Checklist (CBCL) from the Achenbach System of Empirically Based Assessments was used to evaluate externalizing behavior during childhood (Achenbach and Rescorla, 2001). Each participant's classroom teacher assessed symptoms through the Teacher Report Form obtained from the research team via the school principal. The Teacher Report Form includes 112 items rated from 0 to 2 (0 = not true, 1 = somewhat or sometimes true, 2 = very true or often true). It allows for computing various syndrome scores by summing the scores of specific items. The externalizing problems score was obtained by combining both the aggressive behavior and the rule-breaking behavior scores (sum of 32 of the 112 items). The Teacher Report Form had never been used with the Inuit population before the NCDS-childhood study (Boucher et al. 2012), but internal consistency of the externalizing subscale was high ($\alpha_{\text{ordinal}} = 0.97$). No normative data was available for Inuit children, thus raw scores were used in all statistical analyses as in the childhood study.

Child hyperactivity-impulsivity. The Disruptive Behavior Disorders Rating Scale (DBD), designed to be completed by parents and teachers, provides the information necessary for clinical diagnoses of disruptive behaviors in children (Pelham et al., 1992).

The questionnaire is composed of 45 behavioral descriptors rated on a four-point frequency scale ('never', 'sometimes', 'often', 'very often') based on the *Diagnostic and Statistical Manual of Mental Disorders, 4th edition* (DSM-IV). A symptom was considered present if reported as 'often' or 'very often' by the participant's classroom teacher, and the symptom score (continuous) was computed as the count of hyperactivity-impulsivity symptoms. This assessment tool has never been validated in the Inuit population; however, internal consistency was high ($\alpha_{\text{ordinal}} = 0.92$). Even though Boucher et al. (2012) used the diagnostic/dichotomous scores, we used the continuous scores to maximize statistical power in the context of a mediation model.

Child oppositional defiant and conduct disorder (OD/CD). The DBD was also used to evaluate symptoms of oppositional defiant disorder and conduct disorder in children. The DBD has two separate scales for oppositional defiant disorder (8 items) and conduct disorder (15 items; Pelham et al. 1992). A symptom was considered present if reported as 'often' or 'very often' by the participant's classroom teacher. Because most (88%) children identified as CD were also identified as OD, children meeting criteria for either of these diagnoses were grouped together in the statistical analyse (Boucher et al., 2012c). Internal consistency was high ($\alpha_{\text{ordinal}} = 0.95$) and the continuous score was retained in analyses.

Adolescent behavior assessment

Adolescent externalizing. The Youth Self-Report 2001-revision (YSR) from the Achenbach System of Empirically Based Assessments was used to evaluate externalizing behavior during adolescence (Achenbach and Rescorla, 2001). This screening tool is intended for use with youths up to 18 years of age. The interviewer asked the participant to rate the frequency of their symptoms from 0 to 2 (0 = not true, 1 = somewhat or sometimes true, 2 = very true or often true) for each of the 112 items included in the self-report form. The externalizing problems score was obtained by combining the aggressive behavior and the rule-breaking behavior scores (sum of 32 of the 112 items). The YSR has never been used in the Inuit population before, but internal consistency of the externalizing subscale was high ($\alpha_{\text{ordinal}} = 0.88$). No normative data was available for Inuit adolescents, thus raw score were used in all statistical analyses.

Adolescent hyperactivity-impulsivity. The Barkley Adult ADHD-IV Rating Scale (BAARS) is a questionnaire based on the DSM-IV-TR diagnostic criterion for ADHD, which also assesses ADHD symptoms and subtypes (inattention, hyperactivity-impulsivity) and total ADHD in American adults aged 18 to 81 years of age (Barkley, 2011). The subtype hyperactivity-impulsivity was used for its similarity with the childhood measure of hyperactivity-impulsivity. The interviewer asked participants to rate how often they exhibited each symptom ('never', 'sometimes', 'often', 'very often'). A symptom was considered present if reported as present 'often' or 'very often', and a symptom score was computed as the count of symptoms present for hyperactivity-impulsivity items (5 and 4 items, respectively). The BAARS has never been validated in the Inuit population, but the internal consistency of the selected subscale was acceptable ($\alpha_{\text{ordinal}} = 0.76$).

Adolescent conduct disorder. The Diagnostic Interview Schedule for Children (DISC-IV) is a structured diagnostic instrument originally developed to assess 34 psychiatric disorders in epidemiological studies on children and adolescents. Designed to collect information corresponding to the DSM-IV diagnostic criterion, it has since been used in clinical studies and service settings (Shaffer et al., 2000). The eight-item subsection of the DISC-IV on conduct disorder was included in the adolescent follow-up. We included it in our analyses for its similarity with the childhood measure of OD/CD. The participants were asked if the behavior described in each item applied to them (yes/no). The count of items present was used as an adolescent conduct disorder score. Neither the DISC-IV nor any subsection were validated in the Inuit population, but internal consistency in our sample was moderate ($\alpha_{\text{ordinal}} = 0.77$).

Adolescent substance use. The DEP-ADO 3.2 is a substance use evaluation grid originally developed and tested for use among Quebec's French-speaking adolescents (12 to 18 years old; Landry et al. 2004). The seven-question grid evaluates frequency, intensity and age of initiation of alcohol and drug use. Additional items were added to our questionnaire to describe the drinking environment (during a meal, at a party, etc.) and motives. Questions about age of substance use initiation were added at the third of four data collection trips, therefore age at first use data was available only for half of the

participants. The binge drinking score (continuous) was defined as the number of days, in the last year, where more than five/eight alcohol beverages were consumed in one occasion (for women/men). The cannabis use frequency was modeled at four levels: never, $\leq$ once a month, 1-6 times per week and daily.

Covariates

Paths in the mediation analysis (exposure to mediator, mediator to outcome, direct and indirect effect) are modeled by two regression analyses. Therefore, for each regression model, a different set of confounding variables was investigated. An *a priori* list of candidate variables was built for each relation based on previous knowledge and the Inuit population's specificities, including child age and sex (Claycomb et al., 2004), socioeconomic status (SES) (Froehlich et al., 2007), age of biological mother at delivery (Claycomb et al., 2004), maternal tobacco use during pregnancy (Huang et al., 2018), birth weight (Lim et al., 2018), gestational age (Bhutta et al., 2002), adoption status (Decaluwe et al., 2015), and house crowding (Solarig and Mare, 2012).

Confounders in the prediction of the child behavior scores. The same confounding variables as those selected in Boucher et al. (2012) were included in the prediction of childhood behaviors (mediators) from childhood Pb exposure: child age, sex, and birth weight, and childhood socioeconomic status, age of biological mother at delivery, maternal tobacco use during pregnancy, and cord blood Hg

Confounders in the prediction of adolescent outcomes. Age at adolescent testing, sex, and SES of principal care provider at the adolescence follow-up were treated as obligatory covariates. Adolescent schooling level was not considered because of its presence in the hypothetical causal pathway.

The potential covariates considered for the association between child blood Pb concentrations and adolescent outcomes, as well as for the association between child behavior (mediators) and adolescent outcomes were the same. They included a) participant characteristics: birth weight, adoption status (yes/no), and suicidal thoughts in adolescence (yes/no); b) maternal and family characteristics: gestational age, breast feeding duration, primary caregiver education level, age and parity of biological mother at delivery, marital status (single or not), maternal non verbal reasoning ability at child

1
2
3
4 290 follow-up (Raven et al., 1992), and childhood and adolescent domestic crowding (> 1
5
6 291 person per room; yes/no); c) contaminants and nutrients: cord, child and adolescent blood
7
8 292 Hg concentrations, and DHA expressed in percentage by weight of total fatty acids in
9
10 293 plasma phospholipids. Blood lead concentrations measured in cord blood and at
11
12 294 adolescent follow-up were also considered.

13 14 295 **Statistical analyses**

15
16 296 Descriptive statistics, t-test, Wilcoxon tests and Pearson's Chi-squared tests were
17
18 297 computed with the R 3.5.0 software (R Core Team, 2018). The *psych* and *GPArotation*
19
20 298 packages were used to obtain ordinal alphas, Pearson correlations and confidence
21
22 299 intervals (Bernaards and Jennrich, 2005; Revelle, 2018). The *RVAideMemoire* package
23
24 300 was used to compute Spearman correlations and confidence intervals (Hervé, 2018). The
25
26 301 *BaylorEdPsych* package was used to perform missing data pattern analysis and test the
27
28 302 missing completely at random (MCAR) assumption (Beaujean, 2012).

29
30 303 **Assumptions.** Continuous variables' distributions were visually checked for
31
32 304 normality and the following variables were log₂-transformed: contaminant variables,
33
34 305 mother's age at delivery, child externalizing problems scores and number of binge
35
36 306 drinking days in the last year. We analysed the missing value patterns to assess potential
37
38 307 bias. The most common missing data were adolescent domestic crowding (5.2%, *n*=11),
39
40 308 binge drinking (4.7%, *n*=10), cannabis use (4.2%, *n*=9) and pregnancy tobacco use
41
42 309 (2.8%, *n*=6). Little's MCAR test indicated no systematic bias arising from missingness
43
44 310 ($\chi^2 = 308.81$, *df* = 324, *p* = 0.72; Little 1988).

45
46 311 **Bivariate associations.** We computed either Pearson or Spearman correlations
47
48 312 depending on the distribution of the variables to document associations between
49
50 313 mediators and outcomes (Table 2). Correlations were also performed to document
51
52 314 associations between adolescent blood Pb concentrations and adolescent outcomes.

53
54 315 **Confounders selection.** Linear and ordinal regressions were performed prior to
55
56 316 mediation analysis to evaluate the final inclusion of confounding variables in models.
57
58 317 Covariates were included in a forward selection process: variables associated (*p* < 0.20)
59
60 318 with both the mediator and the outcome were entered in decreasing order of association
61
62 319 with the outcome and were retained if they altered the association by $\pm 10\%$ (Rothman et

al., 2008). When the correlation between two *a priori* potential confounders was greater than $p=0.4$ we only retained the one with the strongest association with the outcome. This selection process of confounding variables followed the same principles and criteria previously used by Boucher et al. (2012).

Mediation analyses. Mediation analyses were conducted to examine the direct and indirect association between child Pb exposure and adolescent externalizing behaviors/substance use by testing three different childhood behaviors scores as mediators based on the results of Boucher et al. (2012; Figure 2). Six mediation analyses were conducted with the three childhood behavior scores as mediators (externalizing, hyperactive-impulsive and OD/CD) to predict their corresponding adolescent behavioral outcomes. Three more were conducted to predict adolescent substance use outcomes (binge drinking and cannabis use) through the same mediators (Figure 2). The parameters were estimated using structural equation modeling in Mplus 8.1 (Muthén and Muthén, 2017).

Within the mediation models, linear regressions were used to model continuous outcomes, whereas probit regression was used for the ordinal outcome (cannabis use). The product of coefficients method was used to estimate the indirect effect in all models (MacKinnon et al., 2007). Full information maximum likelihood (FIML) or weighted least squares means variance adjusted (WLSMV, for the ordinal outcome) estimators were used to minimise the exclusion of participants because of missing values (Enders and Bandalos, 2001). The assumption of missing at random (for FIML) and missing at random with respect to independent variables (for WLSMV) were both met since our data were considered missing completely at random based on Little's MCAR test (Asparouhov and Muthén, 2010). The bias-corrected and accelerated 95% confidence intervals (95% CI) were estimated by bootstrap (10 000 resamples; Hayes & Scharkow, 2013). Non-parametric bootstrap methods were preferred to protect against the effects of departure from the normality assumption underlying parametric confidence intervals (Carpenter and Bithell, 2000). Model fit was assessed via three fit statistics: the model chi-square (χ^2), the comparative fit index (CFI) and the root mean square error of approximation (RMSEA; Hooper, Coughlan, & Mullen, 2008).




Figure 2. Path diagram showing the associations tested in mediation analyses. Direct effect refer to the association from child blood lead concentrations and externalizing behavior/substance use in the adolescence follow-up. Indirect effect refer to the association of child blood lead concentrations on adolescent's externalizing behaviors/substance use outcomes through childhood behaviors scores.

1
2
3
4 351 **Results**

5
6 352 *Descriptive analyses*

7
8
9 353 *Sample characteristics.* Participants ranged in age from 9-14 years (median 11.37
10 354 years) at the childhood follow-up visit and between 16-22 years at the adolescent follow-
11 355 up (median 18.48 years; see Table 1). A third of the participants were still in school at the
12 356 adolescent follow-up, whereas more than 35% were fully employed. The median blood
13 357 Pb concentrations were higher during childhood (2.07 µg/dL) than at the adolescence
14 358 (1.52 µg/dL). The majority of participants showed few symptoms of externalizing
15 359 behaviors at both the childhood and adolescent follow-ups. Substance use was very
16 360 common among adolescents, with over 40% of participants drinking alcohol at least
17 361 weekly and more than half using cannabis at least weekly. Tobacco was the most
18 362 widespread substance used, with nearly 8 of 10 participants (79.50%) smoking daily.

19
20
21
22
23
24
25
26
27 363 Of the 294 participants at the childhood follow-up, 212 participated in the
28 364 adolescent follow-up (72%). Men were more likely to be lost at follow-up (64.6% vs.
29 365 44.3%, $p=0.003$). Participants with higher scores on the OD/CD scale were more likely to
30 366 be lost at follow-up (mean of 2.33 for participants at the adolescent follow-up and of 3.73
31 367 for those lost).

Variables	<i>N</i>	Mean ± SD or <i>n</i> (%)	Median	Range
Family/birth characteristics				
Marital status (% married or living with someone)	211	156 (73.90)		
Parity before child birth	212	1.97 ± 1.79	2.00	0.00 – 8.00
Maternal tobacco smoking pregnancy (% yes)	206	177 (85.92)		
Maternal age at delivery (years)	212	23.69 ± 5.67	22.25	15.00 – 42.00
Gestational age (weeks)	212	39.20 ± 1.44	39.00	36.00 – 44.00
Birth weight (kg)	212	3.48 ± 0.46	3.50	2.44 – 4.74
Child sex (% girls)	212	118 (55.66)		
Breast-feeding status (% yes)	207			
None		52 (25.62)		
0 < 3 months		36 (17.73)		
3 < 6 months		24 (11.82)		
≥ 6 months		91 (44.83)		
Cord blood Pb, geomean ± GSD (µg/dL)	204	3.80 ± 1.84	3.73	0.83 – 17.80
Cord blood Hg, geomean ± GSD (µg/dL)	204	1.52 ± 2.15	1.53	0.18 – 9.93
Cord blood DHA (% total fatty acids)	203	1.60 ± 1.29	3.44	1.12 – 7.73
Childhood characteristics				
Age (years)	212	11.34 ± 0.71	11.37	9.32 – 13.97
Adoption status (% adopted)	212	33 (15.57)		
Primary caregiver education (years completed)	211	8.45 ± 2.61	9.00	0.00 – 16.00
SES score ^a	212	28.59 ± 11.38	28.25	8.00 – 66.00
Behaviors (teachers assessment)				
Externalizing - CBCL	205	14.23 ± 12.78	11.00	0.00 – 53.00
Hyperactivity-impulsivity - DBD	207	1.72 ± 2.51	0.00	0.00 – 9.00
Oppositional defiant/conduct disorder - DBD	207	2.33 ± 3.37	1.00	0.00 – 16.00
Blood Pb, geomean ± GSD (µg/dL)	210	2.34 ± 1.86	2.07	0.54 – 12.83
Blood Hg, geomean ± GSD (µg/dL)	210	3.36 ± 2.56	3.41	0.05 – 28.02
Blood DHA (% total fatty acids)	209	2.40 ± 0.97	2.21	0.60 – 4.96
Adolescent characteristics				
Age (years)	212	18.47 ± 1.11	18.48	16.01 – 21.88
Main language at home (% mainly Inuktitut)	210	200 (95.24)		
SES principal provider	207	28.59 ± 13.01	28.00	8.00 – 61.00
In a relationship (% yes)	209	95 (45.45)		
Education	210		3.00	0.00 – 5.00
Secondary 1 or less		46 (21.9)		
Secondary 2 or 3		105 (50.0)		
Secondary 4 or higher		59 (28.1)		
Occupational status (% yes)	210			
Working		75 (35.71)		
School		34 (16.19)		
Both		37 (17.62)		
None		64 (30.48)		
Behaviors (self-reported)				
Externalizing - CBCL	210	15.80 ± 7.46	15.00	0.00 – 40.00
Hyperactivity-impulsivity - BAARS	203	1.59 ± 1.52	1.00	0.00 – 6.00
Conduct disorder - DISC	212	0.94 ± 1.21	1.00	0.00 – 5.00
Substance use				
Age started drugs regular basis (years)	127	14.72 ± 1.93	15.00	7.00 – 19.00
Age started alcohol regular basis (years)	106	15.76 ± 1.63	16.00	10.00 – 19.00
Cannabis use (% , last 12 months)	203			
Never		63 (31.03)		
≤ 1 / month		34 (16.75)		
1-6 times / week		42 (20.69)		
Everyday		64 (31.53)		
Tobacco smoking (% , last 12 months)	200			
None		24 (12.00)		
Occasional		17 (8.50)		
Daily		159 (79.50)		
Alcohol use (% last 12 months)	203	177 (87.19)		
Binge drinking days (number in last year) ^b	176	36.40 ± 47.16	12.00	0.00 – 250.00
Blood Pb, geomean ± GSD (µg/dL)	212	1.63 ± 2.00	1.52	0.35 – 18.13
Blood Hg, geomean ± GSD (µg/dL)	212	3.82 ± 2.76	4.21	0.14 – 36.11
Blood DHA (% total fatty acids)	212	3.35 ± 1.21	3.32	0.60 – 7.12

Note: Pb, lead; Hg, mercury; DHA, docosahexaenoic acid; GSD, geomean standard deviation; CBCL, Child Behavior Checklist; DBD, Disruptive Behavior Disorders Rating Scale; BAARS, Barkley Adult ADHD-IV Rating Scale; DISC, Diagnostic Interview Schedule for Children.

^aAssessed on the Hollingshead Index, which represents a weighted score of parental occupation (type of work) and education level (Hollingshead, 1975).

^bBinge drinking, among drinkers, corresponds to consumption of ≥ 5 standard drinks per occasion for women and 8 for men; 1 standard drink ~ 0.5 oz absolute alcohol (~ 350 mL of beer (12 oz), 175 mL of wine (6 oz), or 44 mL of liquor (1.5 oz)).

Correlations between externalizing symptoms/behaviors. Associations between mediators and outcomes are shown in Table 2. All three childhood behavior scores were highly correlated (all correlation coefficients ≥ 0.65). The childhood externalizing problem and OD/CD scores were both significantly but weakly related to adolescent externalizing problem scores and cannabis use. Adolescent externalizing problem score was associated with all the adolescent behaviors; whereas child hyperactivity-impulsivity and the adolescent hyperactivity-impulsivity scores were poorly correlated. The child OD/CD score was poorly associated with adolescent conduct disorder score. Cannabis use was moderately related to adolescent conduct disorder score and binge drinking.

Correlations between blood Pb and adolescent behaviors scores. Associations between blood Pb concentrations and adolescent outcomes are summarized in Table 2. No significant association was observed between child blood Pb concentrations and adolescent outcomes nor between adolescent blood Pb concentrations and adolescent externalizing or hyperactivity-impulsivity symptoms, conduct disorder, or cannabis use. By contrast, significant associations were observed between adolescent blood Pb concentrations and binge drinking.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

392 Table 2 – Pearson and Spearman correlation coefficients (95% Confidence Intervals) between mediators and outcomes

	Childhood			Adolescence					
	Externalizing ^a	Hyperactivity-impulsivity ^a	OD/CD ^a	Blood Pb	Externalizing ^b	Hyperactivity-impulsivity ^b	Conduct disorder ^b	Binge drinking ^b	Cannabis use ^b
Childhood									
Blood Pb	0.26 (0.12, 0.39)	0.21^{sp} (0.07, 0.35)	0.20^{sp} (0.06, 0.32)	0.40 (0.28, 0.51)	0.08 (-0.04, 0.22)	0.10 (-0.04, 0.23)	0.08 ^{sp} (-0.06, 0.21)	-0.03 (-0.17,0.08)	0.12 ^{sp} (-0.02, 0.25)
Externalizing ^a	1	0.73^{sp} (0.66, 0.79)	0.82^{sp} (0.77, 0.87)	0.25 (0.13, 0.36)	0.20 (0.009, 0.32)	-0.003 (-0.13, 0.14)	0.09 ^{sp} (-0.05, 0.22)	0.14 (-0.02, 0.28)	0.20^{sp} (0.06, 0.33)
Hyperactivity-impulsivity ^a	-	1	0.65^{sp} (0.56, 0.72)	0.18^{sp} (0.04, 0.30)	0.11 ^{sp} (-0.02, 0.25)	-0.01 ^{sp} (-0.16, 0.14)	0.07 ^{sp} (-0.07, 0.21)	0.05 ^{sp} (-0.09, 0.19)	0.10 ^{sp} (-0.04, 0.23)
OD/CD ^a	-	-	1	0.21^{sp} (0.08, 0.33)	0.19^{sp} (0.05, 0.32)	-0.04 ^{sp} (-0.17, 0.10)	0.08 ^{sp} (-0.05, 0.22)	0.10 ^{sp} (-0.04, 0.24)	0.23^{sp} (0.08, 0.36)
Adolescence									
Blood Pb	-	-	-	1	0.01 (-0.12, 0.15)	-0.04 (-0.17, 0.09)	0.05 ^{sp} (-0.08, 0.19)	0.16 (0.03, 0.30)	0.13 ^{sp} (-0.02, 0.27)
Externalizing ^b	-	-	-	-	1	0.44 (0.32, 0.57)	0.34^{sp} (0.21, 0.45)	0.22 (0.10, 0.34)	0.32^{sp} (0.19, 0.43)
Hyperactivity-impulsivity ^b	-	-	-	-	-	1	0.16^{sp} (0.03, 0.29)	0.10 (-0.04, 0.24)	-0.02 ^{sp} (-0.15,0.12)
Conduct disorder ^b	-	-	-	-	-	-	1	0.09 ^{sp} (-0.05, 0.23)	0.18^{sp} (0.05, 0.31)
Binge drinking ^b	-	-	-	-	-	-	-	1	0.17^{sp} (0.03, 0.30)

393 Sample size for individual comparisons varies between 195 and 212 due to missing values.
394 Note: OD/CD, Oppositional defiant/conduct disorder;^a teachers assessment, ^b self-reported with assistance of interviewer, ^{sp}, Spearman coefficient.

Mediation analyses

Associations between child Pb and adolescent behavior scores. We investigated the effect of child blood Pb concentrations on adolescent behavior scores, directly and indirectly through their corresponding child behavior scores (Table 3). No direct association was observed between child Pb concentrations and any of the adolescent behavior scores. However, a significant indirect association was observed between child blood Pb concentrations and adolescent externalizing problem through child externalizing problem scores [0.32 (95% CI: 0.08, 0.72)]. Thus, a 2-fold increase in blood Pb concentrations units explained on average an increase of 0.25 symptoms on the adolescent externalizing problem score. No indirect association was observed with the adolescent hyperactivity-impulsivity or with adolescent conduct disorder scores. Fit statistics were good for the adolescent externalizing problem and hyperactivity-impulsivity models and acceptable for the adolescent conduct disorder model.

Association between child Pb and adolescent substance use. We examined the effect of child blood Pb concentrations on adolescent substance use, directly and indirectly through child behavior scores (Table 4). No direct association was observed between child Pb exposure and any of the adolescent substance use. Significant indirect associations were observed between child blood Pb concentrations and both binge drinking and cannabis use through child externalizing problem scores [0.09 (95% CI: 0.02, 0.23)] and [0.05 (95% CI: 0.002, 0.14)], respectively. A 2-fold increase in blood Pb concentrations explains on average an increase of 0.06 binge drinking episode per year. There is no straight-forward way to assess the effect size of an ordinal regression included in a mediation model. The 0.04 indirect association estimate in the cannabis use model can be interpreted as the probability to move from an inferior category of cannabis use frequency to a superior one for each 2-fold increase in blood Pb concentrations units. Fit statistics were acceptable for binge drinking models and good for cannabis use models.

422 1 Table 3 - Mediation analysis of childhood Pb exposure on adolescent externalizing behaviors, using childhood behavior scores as mediator. Regression coefficients (95% CI) and fit indices (n=212).

		Adolescent		
		Externalizing	Hyperactivity-impulsivity	Conduct disorder
		Estimate (95% CI)	Estimate (95% CI)	Estimate (95% CI)
5	Mediator			
7	Child externalizing (log ₂ -transformed)			
8	Child Pb → Child externalizing ^a	0.42 (0.15, 0.69)		
9	Child externalizing → Adolescent externalizing ^b	0.77 (0.15, 1.36)		
10	Direct effect	0.61 (-0.63, 1.96)		
11	Indirect effect			
12		0.32 (0.08, 0.72)		
13	χ ² (df), <i>p</i> -value	10.24(8), 0.25		
14	CFI	0.92		
15	RMSEA (90% CI)	0.04 (0.00, 0.09)		
17				
18	Child hyperactivity-impulsivity			
19	Child Pb → Child hyperactivity-impulsivity ^a		0.58 (0.17, 0.99)	
20	Child hyperactivity-impulsivity → Adolescent hyperactivity-impulsivity ^c		-0.03 (-0.12, 0.06)	
21	Direct effect		0.11 (-0.14, 0.37)	
22	Indirect effect		-0.02 (-0.09, 0.03)	
23	χ ² (df), <i>p</i> -value		5.01(8), 0.75	
24	CFI		1.00	
25	RMSEA (90% CI)		<0.01 (0.00, 0.06)	
27				
28	Child oppositional defiant/conduct disorder			
29	Child Pb → Child OD/CD ^a			0.67 (0.11, 1.25)
30	Child OD/CD → Adolescent conduct disorder ^d			0.02 (-0.03, 0.08)
31	Direct effect			0.02 (-0.20, 0.21)
32	Indirect effect			0.02 (-0.01, 0.07)
33	χ ² (df), <i>p</i> -value			7.29(7), 0.40
34	CFI			0.98
35	RMSEA (90% CI)			0.01 (0.002, 0.09)

423 37 Note : CI, Confidence interval; χ²(df), chi square (degrees of freedom); CFI, comparative fit index; RMSEA, root mean square error of approximation; OD/CD, Oppositional defiant/conduct disorder.

424 38 ^aAdjusted for child age and sex, SES, age of the biological mother at delivery, maternal tobacco smoking during pregnancy, and birth weight. Models with child hyperactivity-impulsivity and child OD/CD as mediator were also
425 39 adjusted for cord Hg (log₂-transformed).

426 40 ^bAdjusted for adolescent age and sex, provider SES, and child blood Hg (log₂-transformed).

427 41 ^cAdjusted for adolescent age and sex, provider SES, age of the biological mother at delivery and education level of the primary caregiver at child follow-up .

428 42 ^dAdjusted for adolescent age and sex, provider SES, age of the biological mother at delivery, maternal tobacco use during pregnancy, and house crowding at adolescence follow-up.

429 43

44

45

46

47

48

49

430	1	Table 4 - Mediation analysis of childhood Pb exposure on adolescent substance use, using childhood behavior scores as mediator. Regression coefficients (95% CI) and fit indices (<i>n</i> =212).		
2	2		Adolescent	
3	3			
4	4	Mediator	Binge drinking (log ₂ -transformed)	Cannabis use
5	5		Estimate (95% CI)	Estimate (95% CI)
6	6	Child externalizing (log ₂ -transformed)		
7	7	Child Pb → Child externalizing ^a	0.41 (0.14, 0.69)	0.45 (0.01, 0.85)
8	8	Child externalizing → Adolescent substance use ^b	0.22 (0.02, 0.43)	0.11 (0.01, 0.20)
9	9	Direct effect	-0.23 (-0.65,0.21)	0.04 (-0.24, 0.31)
10	10	Indirect effect	0.09 (0.02, 0.23)	0.05 (0.002, 0.14)
11	11	χ ² (df), <i>p</i> -value	12.39(9), 0.19	16.69(11), 0.12
12	12	CFI	0.92	0.98
13	13	RMSEA (90% CI)	0.04 (0.00, 0.09)	0.05 (0.00, 0.10)
14	14			
15	15			
16	16	Child hyperactivity-impulsivity		
17	17	Child Pb → Child hyperactivity-impulsivity ^a	0.57 (0.16, 0.99)	0.59 (0.09, 1.06)
18	18	Child hyperactivity-impulsivity → Adolescent substance use ^c	-0.001 (-0.14,0.14)	0.03 (-0.04, 0.09)
19	19	Direct effect	-0.15 (-0.58, 0.31)	0.10 (-0.15, 0.40)
20	20	Indirect effect	0.001 (-0.09, 0.09)	0.02 (-0.02, 0.08)
21	21	χ ² (df), <i>p</i> -value	13.06(10), 0.22	9.73.(11), 0.56
22	22	CFI	0.90	1.00
23	23	RMSEA (90% CI)	0.04 (0.00, 0.09)	0.00 (0.00, 0.07)
24	24			
25	25			
26	26	Child oppositional defiant/conduct disorder		
27	27	Child Pb → Child OD/CD ^a	0.66 (0.11, 1.25)	0.68 (-0.02, 1.33)
28	28	Child OD/CD → Adolescent substance use ^d	0.04 (-0.07, 0.15)	0.04 (-0.01, 0.08)
29	29	Direct effect	-0.17 (-0.59, 0.28)	0.09 (-0.18, 0.38)
30	30	Indirect effect	0.02 (-0.04, 0.15)	0.02 (-0.002, 0.08)
31	31	χ ² (df), <i>p</i> -value	20.05(12), 0.07	11.09(10), 0.35
32	32	CFI	0.74	0.99
33	33	RMSEA(90% CI)	0.06 (0.00, 0.10)	0.02 (0.00, 0.08)
34	34			
431	35	Note : CI, Confidence interval; χ ² (df), chi square (degrees of freedom); CFI, comparative fit index; RMSEA, root mean square error of approximation; OD/CD, Oppositional defiant/conduct disorder.		
432	36	^a Adjusted for child age and sex, SES, age of the biological mother at delivery, maternal tobacco use during pregnancy, and birth weight. Models with child hyperactivity-impulsivity and child OD/CD as mediator were also adjusted		
433	37	for cord Hg (log ₂ -transformed).		
434	38	^b Both models adjusted for adolescent age and sex, provider SES, and adolescent blood Pb (log ₂ -transformed). Model with binge drinking as outcome was also adjusted for birth weight, adoption status, and house crowding at		
435	39	adolescence follow-up. Model with cannabis use as outcome was also adjusted for breast feeding duration.		
436	40	^c Both models adjusted for adolescent age and sex, provider SES, and adolescent blood Pb(log ₂ -transformed). Model with binge drinking as outcome was also adjusted for birth weight, adoption status and house crowding at		
437	41	adolescence follow-up. Model with cannabis use as outcome was also adjusted for educational level of the primary caregiver at child follow-up and child blood Hg.		
438	42	^d Both models adjusted for adolescent age and sex, provider SES, and adolescent blood Pb (log ₂ -transformed). Model with binge drinking as outcome was adjusted for adoption status and house crowding at adolescence follow-up.		
439	43	Model with cannabis use as outcome was also adjusted for educational level of the primary caregiver at child follow-up		
44	44			
45	45			
46	46			
47	47			
48	48			
49	49			

440 Discussion

441 The main objective of this study was to test the developmental cascade linking child
442 blood Pb concentrations with adolescent behavior problems/substance use through
443 behavior problems assessed in childhood. Child blood Pb concentrations were associated
444 with behavioral challenges at the same age as previously reported by Boucher et al.
445 (2012), and these problems, in turn, predicted externalizing behaviors and substance use
446 in adolescence. Direct associations between child blood Pb concentrations and any of the
447 adolescent behavior assessments or with substance use were not significant. Thus, child
448 Pb concentrations were associated with some problem outcomes in adolescence but only
449 indirectly through child externalizing problem score. No indirect associations linked
450 childhood Pb concentrations with adolescent outcomes through child hyperactivity-
451 impulsivity or conduct disorder scores. While the presence of a direct effect was
452 previously considered as the first step in a mediation analysis, it is now recognized that
453 some relationships may be only indirect through a mediator. This is particularly relevant
454 in the longitudinal investigation of externalizing behavior given the persistence from
455 childhood to adolescence/adulthood of behaviors reported in other cohorts (Reef et al.,
456 2011).

457 The indirect associations observed in our study between child Pb concentrations
458 and adolescent outcomes are consistent with the developmental cascades hypothesized by
459 Bellinger et al. (2016). In a developmental framework, such a cascade would start with an
460 early-life neurotoxic exposure affecting some child characteristics, which in turn
461 modifies the neurocognitive development, and consequently cause lifelong impairments.
462 In this context, an intermediary endpoint is considered as a mediating factor between the
463 child's prior exposure and later end-points (Bellinger et al., 2016). The developmental
464 cascades framework predicts direct paths between early exposure and childhood
465 outcomes and indirect paths between the exposure and later impairments, as observed in
466 the present study with externalizing problem score, binge drinking and cannabis use. The
467 underlying mechanisms explaining the persistence of externalizing behaviors (and
468 manifestation of substance use) in adolescence remain unclear. In a previous study, our
469 team reported a significant association between child Pb exposure and increased

impulsivity showed by impairments in child's ability to correctly inhibit a response (Boucher et al. 2012b). These results were consistent with the proposition McGue and colleagues (2001) that a vulnerability to disinhibitory behavior and impulse control might explain those associations.

The few studies that investigated the effect of child blood Pb concentrations on adolescent behavior problems only reported total effects without decomposing them in direct and indirect paths. One such study reported that an increase of 1 µg/dL of the average child blood Pb concentrations was associated with an increase of 0.06 standard deviation on the adolescent impulsivity symptoms score (Winter and Sampson, 2017). In our study, no significant direct or indirect association was observed between child blood Pb on the adolescent hyperactivity-impulsivity scale. The inclusion of adolescent blood Pb concentrations did not modify the relationship between child Pb concentrations and adolescent outcomes even if it was associated with substance use outcomes. However, we observed indirect associations – through child externalizing behaviors – showing that a 2-fold increase in blood Pb concentrations can explain 0.25 symptoms on the adolescent externalizing score and an increase of 0.06 binge drinking episode. As pointed out by Bellinger et al. (2016), longitudinal studies are able to provide a more complete picture of the burden of child Pb exposure on later impairments if direct/indirect effects are considered instead of the traditional approach of only considering the total effect. The small magnitude of the associations we reported was expected given that adolescent behavior problems are complex outcomes with multiple predictors; we would not expect a single contaminant exposure to explain a large proportion of these behaviors, particularly in Inuit communities where historical and social determinants, coupled with limited educational and social services, may play a predominant role.

Strengths

The major strength of this paper is its longitudinal design, which documented exposure and behavior beyond the critical child developmental period and allowed the use of mediation analyses as a novel analytic approach in the investigation of Pb exposure on future impairments, yet well-grounded in Bellinger et al.'s (2016) developmental cascade framework. Second, externalizing behaviors at the childhood and

adolescent follow-up were assessed by different raters, namely the school teachers and the participants themselves, ensuring independence of behavioral measures. Additionally, a longitudinal design reduces the potential reverse causation between externalizing behaviors and substance use; the overwhelming majority of our participants started to consume alcohol or drugs regularly after the childhood assessment of externalizing behaviors.

Limitations

One of the key limitations of our study is its relatively small sample size. The social precariousness in Inuit communities may explain why many participants were not eligible for the adolescence follow-up. We had a retention rate of 72% between the child and adolescent follow-up. Most participants ineligible to participate to the adolescent follow-up were unreachable because of incarceration, hospitalization or death. All of these are known to be related to gender and oppositional defiant/conduct disorder, which may explain differences in the adolescent follow-up sample (Begg et al., 1999; Malakieh, 2018; Mordre et al., 2011; Sorenson, 2011). Furthermore, because of limited statistical power, we were not able to conduct sex effect modification analyses even though externalizing behaviors and substance use manifestations are likely to differ between boys and girls (Hammerslag and Gulley, 2016). Additionally, nine mediation models were tested, which raises the risk of type I error. However, it is important to note that significant results were not randomly distributed: the indirect associations were consistently present in models mediated by externalizing behaviors. Next, our sample was comprised of Inuit children from northern Québec, who are more exposed to Pb than children from the general Canadian population (2.34 µg/dL compared to 0.8-0.9 µg/dL, respectively; Health Canada 2010). This and the major historical, cultural and societal differences of Inuit population with the southern Canadian population, mean that the generalization of our results is not straightforward. We have consider many potential covariates measured at different follow-ups, but it is always possible that some were omitted. Also, lead exposure is usually highest around age 1 to 3, while our measure of Pb concentration was taken a later age. Using early-life Pb exposure might have shown stronger associations. Still we were able to observed significant association between later childhood exposure and behavioral outcomes. Further cohort studies on neurobehavioral

1
2
3
4 531 outcomes should measure blood Pb concentrations at an earlier age, during the peak of
5
6 532 exposition. Finally, the behavior measurement tools used in this study were all developed
7
8 533 for, validated in non-indigenous populations, and administered by non-Inuit interviewers.
9
10 534 Therefore, they may not fully capture behaviors that are considered as problematic by the
11
12 535 Inuit themselves. However, internal consistencies were adequate in our sample (ordinal
13
14 536 alphas higher than 0.76).

15 16 537 ***Conclusion***

17
18 538 Children exposed to Pb have a higher rate of developmental problems, but uncertainties
19
20 539 remain about what role Pb exposure might play in the relation between childhood
21
22 540 developmental problems and persisting impairments. Our study highlights the indirect but
23
24 541 persistent effects of child Pb exposure on adolescent behavior problems and the
25
26 542 importance of childhood externalizing behavior in this relationship; these early-life
27
28 543 impairments may put children on a impeded developmental trajectory, increasing their
29
30 544 likelihood of lifelong psychological, social and health problems.. Despite the small
31
32 545 contribution of Pb exposure in the prediction of adolescent behavior problems, our results
33
34 546 offer a new perspective in the investigation of the Pb burden on human capital. Further
35
36 547 studies should document the long-term effect of Pb exposure and other neurotoxicants
37
38 548 during adolescence or early adulthood by modelling developmental processes reflecting
39
40 549 the risk of developmental cascades instead of documenting mere associations.

41 550 ***Acknowledgements***

42
43 551 The authors gratefully thank the Nunavik population, all the people involved in this study
44
45 552 and the Nunavik Nutrition and Health Committee for their useful comments on this
46
47 553 manuscript.

48
49
50 554 This longitudinal research was funded by the Institute of Indigenous People's
51
52 555 Health from the Canadian Institutes of Health Research (NRF 130242), NIH/National
53
54 556 Institute of Environmental Health Sciences (R01-ES007902); the Northern Contaminants
55
56 557 Program from Government Canada; the Lycaki-Young, Sr., Fund from the State of
57
58 558 Michigan.

References

- Achenbach, T.M., Rescorla, L.A., 2001. Manual for the ASEBA School-Age Forms & Profiles. University of Vermont, Research Centre for Children, Youth, & Families, Burlington, VT.
- Asparouhov, T., Muthén, B., 2010. Weighted least squares estimation with missing data. Muthén & Muthén, Los Angeles.
- Barkley, R.A., 2011. Barkley Adult ADHD Rating Scale-IV (BAARS-IV). The Guilford Press, New York.
- Beaujean, A.A., 2012. BaylorEdPsych: R Package for Baylor University Educational Psychology Quantitative Courses. R package version 0.5.
- Beckley, A.L., Caspi, A., Broadbent, J., Harrington, H., Houts, R.M., Poulton, R., Ramrakha, S., Reuben, A., Moffitt, T.E., 2018. Association of childhood blood lead levels with criminal offending. *JAMA Pediatr.* 172, 166–173.
<https://doi.org/10.1001/jamapediatrics.2017.4005>
- Begg, D.J., Langley, J.D., Williams, S.M., 1999. A longitudinal study of lifestyle factors as predictors of injuries and crashes among young adults. *Accid. Anal. Prev.* 31, 1–11. [https://doi.org/https://doi.org/10.1016/S0001-4575\(98\)00039-6](https://doi.org/10.1016/S0001-4575(98)00039-6)
- Bellinger, D.C., Matthews-bellinger, J.A., Kordas, K., 2016. A developmental perspective on early-life exposure to neurotoxicants. *Environ. Int.* 94, 103–112.
<https://doi.org/10.1016/j.envint.2016.05.014>
- Bernaards, C.A., Jennrich, R.I., 2005. Gradient Projection Algorithms and Software for Arbitrary Rotation Criteria in Factor Analysis. *Educ. Psychol. Mes.* 65, 676–696.
- Bhutta, A.T., Cleves, M.A., Casey, P.H., Cradock, M.M., Anand, K.J.S., 2002. Cognitive and Behavioral Outcomes of School-Aged Children Who Were Born Preterm. *JAMA - J. Am. Med. Assoc.* 288.
- Boucher, O., Burden, M.J., Muckle, G., Saint-Amour, D., Ayotte, P., Dewailly, É., Nelson, C.A., Jacobson, S.W., Jacobson, J.L., 2012a. Response inhibition and error

586 monitoring during a visual go/no-go task in Inuit children exposed to lead,
 587 polychlorinated biphenyls, and methylmercury. *Environ. Health Perspect.* 120, 608.
 588 <https://doi.org/10.1289/ehp.1103828>

589 Boucher, O., Jacobson, S.W., Plusquellec, P., Dewailly, E., Ayotte, P., Forget-Dubois,
 590 N., Jacobson, J.L., Muckle, G., 2012b. Prenatal methylmercury, postnatal lead
 591 exposure, and evidence of attention deficit/hyperactivity disorder among Inuit
 592 children in Arctic Québec. *Environ. Health Perspect.* 120, 1456.
 593 <https://doi.org/10.1289/ehp.1204976>

594 Boucher, O., Jacobson, S.W., Plusquellec, P., Dewailly, É., Ayotte, P., Forget-Dubois,
 595 N., Jacobson, J.L., Muckle, G., 2012c. Prenatal methylmercury, postnatal lead
 596 exposure, and evidence of attention deficit/hyperactivity disorder among Inuit
 597 children in Arctic Québec. *Environ. Health Perspect.* 120, 1456–1461.
 598 <https://doi.org/10.1289/ehp.1204976>

599 Braun, J.M., Kahn, R.S., Froehlich, T., Auinger, P., Lanphear, B.P., 2006. Exposures to
 600 Environmental Toxicants and Attention Deficit Hyperactivity Disorder in U.S.
 601 Children. *Environ. Health Perspect.* 114, 1904–1909.
 602 <https://doi.org/10.1289/ehp.9478>

603 Burns, J., Baghurst, P., Sawyer, M.G., McMichael, A.J., Tong, S.-L., 1999. Lifetime
 604 Low-Level Exposure to Environmental Lead and Children's Emotional and
 605 Behavioral Development at Ages 11-13 Years. *Public Health* 149, 740–749.

606 Carpenter, J., Bithell, J., 2000. Bootstrap confidence intervals: when, which, what? A
 607 practical guide for medical statisticians. *Stat. Med.* 19, 1141–1164.

608 Chiodo, L.M., Covington, C., Sokol, R.J., Hannigan, J.H., Jannise, J., Ager, J.,
 609 Greenwald, M., Delaney-Black, V., 2007. Blood lead levels and specific attention
 610 effects in young children. *Neurotoxicol. Teratol.* 29, 538–546.
 611 <https://doi.org/10.1016/j.ntt.2007.04.001>

612 Chiodo, L.M., Jacobson, S.W., Jacobson, J.L., 2004. Neurodevelopmental effects of
 613 postnatal lead exposure at very low levels. *Neurotoxicol. Teratol.* 26, 359–371.

- 614 <https://doi.org/10.1016/j.ntt.2004.01.010>
- 615 Claycomb, C.D., Ryan, J.J., Miller, L.J., Schnakenberg-Ott, S.D., 2004. Relationships
616 among attention deficit hyperactivity disorder, induced labor, and selected
617 physiological and demographic variables. *J. Clin. Psychol.* 60, 689–693.
618 <https://doi.org/10.1002/jclp.10238>
- 619 Dallaire, R., Dewailly, É., Ayotte, P., Forget-Dubois, N., Jacobson, S.W., Jacobson, J.L.,
620 Muckle, G., 2014. Growth in Inuit children exposed to polychlorinated biphenyls
621 and lead during fetal development and childhood. *Environ. Res.* 134, 17–23.
622 <https://doi.org/10.1016/j.envres.2014.06.023>
- 623 Decaluwe, B., Jacobson, S.W., Poirier, M.-A., Forget-Dubois, N., Jacobson, J.L.,
624 Muckle, G., 2015. Impact of Inuit customary adoption on behavioral problems in
625 school-age Inuit children. *Am J Orthopsychiatry* 85, 250–258.
626 <https://doi.org/doi:10.1037/ort0000071>
- 627 Dewailly, E., Bruneau, S., Ayotte, P., Laliberté, C., Gingras, S., Bélanger, D., Ferron, L.,
628 1993. Health status at birth of inuit newborn prenatally exposed to organochlorines.
629 *Chemosphere* 27, 359–366. [https://doi.org/10.1016/0045-6535\(93\)90313-T](https://doi.org/10.1016/0045-6535(93)90313-T)
- 630 Dietrich, K.N., Douglas, R.M., Succop, P.A., Berger, O.G., Bornschein, R.L., 2001. Early
631 exposure to lead and juvenile delinquency. *Neurotoxicol. Teratol.* 23, 511–518.
632 [https://doi.org/10.1016/S0892-0362\(01\)00184-2](https://doi.org/10.1016/S0892-0362(01)00184-2)
- 633 Enders, C.K., Bandalos, D.L., 2001. The Relative Performance of Full Information
634 Maximum Likelihood Estimation for Missing Data in Structural Equation Models.
635 *Struct. Equ. Model. A Multidiscip. J.* 8, 430–457.
636 https://doi.org/10.1207/S15328007SEM0803_5
- 637 Firestone, M., Tyndall, M., Fischer, B., 2015. Substance Use and Related Harms among
638 Aboriginal People in Canada: A Comprehensive Review. *J. Health Care Poor*
639 *Underserved* 26, 1110–1131. <https://doi.org/10.1353/hpu.2015.0108>
- 640 Fishbein, D.H., Todd, A.C., Ricketts, E.P., Semba, R.D., 2008. Relationship between lead
641 exposure, cognitive function, and drug addiction: Pilot study and research agenda.

- Environ. Res. 108, 315–319. <https://doi.org/10.1016/j.envres.2008.07.012>
- Froehlich, T.E., Lanphear, B.P., Auinger, P., Hornung, R., Epstein, J.N., Braun, J., Kahn, R.S., 2009. Association of Tobacco and Lead Exposures With Attention-Deficit/Hyperactivity Disorder. *Pediatrics* 124, e1054–e1063. <https://doi.org/10.1542/peds.2009-0738>
- Froehlich, T.E., Lanphear, B.P., Epstein, J.N., Barbaresi, W.J., Katusic, S.K., Kahn, R.S., Objective., 2007. Prevalence, recognition, and treatment of attention-deficit/hyperactivity disorder in a national sample of US children. *Arch. Pediatr. Adolesc. Med.* 161, 857–864. <https://doi.org/10.1001/archpedi.161.9.857> LK - <http://vu.on.worldcat.org/atoztitles/link?sid=EMBASE&issn=10724710&id=doi:10.1001%2Farchpedi.161.9.857&atitle=Prevalence%2C+recognition%2C+and+treatment+of+attention-deficit%2Fhyperactivity+disorder+in+a+national+sample+of+US+children&stitle=Arch.+Pediatr.+Adolesc.+Med.&title=Archives+of+Pediatrics+and+Adolescent+Medicine&volume=161&issue=9&spage=857&epage=864&aulast=Froehlich&aufirst=Tanya+E.&auinit=T.E.&aufull=Froehlich+T.E.&coden=APAME&isbn=&pages=857-864&date>
- Groenman, A.P., Janssen, T.W.P., Oosterlaan, J., 2017. Childhood Psychiatric Disorders as Risk Factor for Subsequent Substance Abuse: A Meta-Analysis. *J. Am. Acad. Child Adolesc. Psychiatry* 56, 556–569. <https://doi.org/10.1016/j.jaac.2017.05.004>
- Hammerslag, L.R., Gulley, J.M., 2016. Sex differences in behavior and neural development and their role in adolescent vulnerability to substance use. *Behav. Brain Res.* 298, 15–26. <https://doi.org/10.1016/j.bbr.2015.04.008>
- Hayes, A.F., Scharkow, M., 2013. The Relative Trustworthiness of Inferential Tests of the Indirect Effect in Statistical Mediation Analysis: Does Method Really Matter? *Psychol. Sci.* 24, 1918–1927. <https://doi.org/10.1177/0956797613480187>
- Health Canada, 2010. Report on Human Biomonitoring of Environmental Chemicals in Canada - Results of the Canadian Health Measures Survey Cycle 1 (2007–2009). Ottawa.

- 1
- 2
- 3
- 4 671 Hervé, M., 2018. RVAideMemoire: Testing and Plotting Procedures for Biostatistics. R
- 5
- 6 672 Packag. version 0.9-69-03.
- 7
- 8
- 9 673 Hollingshead, A., 1975. Four Factor Index of Social Status. Yale University Department
- 10 674 of Sociology, New Haven.
- 11
- 12
- 13 675 Hooper, D., Coughlan, J., Mullen, M., 2008. Structural Equation Modelling: Guidelines
- 14 676 for Determining Model Fit. *Electron. J. Buisness Res.* 6, 53–60.
- 15
- 16
- 17 677 Huang, L., Wang, Y., Zhang, L., Zheng, Z., 2018. Maternal Smoking and Attention-
- 18 678 Deficit / Hyperactivity Disorder in Offspring : A Meta-analysis. *Pediatrics* 141.
- 19
- 20
- 21 679 Jacobson, J.L., Jacobson, S.W., Muckle, G., Kaplan-Estrin, M., Ayotte, P., Dewailly, É.,
- 22 680 2008. Beneficial Effects of a Polyunsaturated Fatty Acid on Infant Development :
- 23 681 Evidence from the Inuit of Artic Quebec. *J. Pediatr.* 152, 356–364.
- 24
- 25
- 26
- 27 682 Krueger, R.F., Markon, K.E., Patrick, C.J., Iacono, W.G., 2005. Externalizing
- 28 683 psychopathology in adulthood: A dimensional-spectrum conceptualization and its
- 29 684 implications for DSM-V. *J. Abnorm. Psychol.* 114, 537–550.
- 30 685 <https://doi.org/10.1037/0021-843X.114.4.537>
- 31
- 32
- 33
- 34
- 35
- 36 686 Landry, M., Tremblay, J., Guyon, L., Bergeron, J., Brunelle, N., 2004. La Grille de
- 37 687 dépistage de la consommation problématique d'alcool et de drogues chez les
- 38 688 adolescents et les adolescentes (DEP-ADO) : développement et qualités
- 39 689 psychométriques. *Drog. santé société* 3, 20–37. <https://doi.org/10.7202/010517ar>
- 40
- 41
- 42
- 43
- 44 690 Lee, S.S., Humphreys, K.L., Flory, K., Liu, R., Glass, K., 2011. Prospective association
- 45 691 of childhood attention-deficit/hyperactivity disorder (ADHD) and substance use and
- 46 692 abuse/dependence: A meta-analytic review. *Clin. Psychol. Rev.* 31, 328–341.
- 47 693 <https://doi.org/10.1016/j.cpr.2011.01.006>
- 48
- 49
- 50
- 51
- 52 694 Lévesque, B., Duchesne, J.-F., Gariépy, C., Rhainds, M., Dumas, P., Scheuhammer,
- 53 695 A.M., Proulx, J.-F., Déry, S., Muckle, G., Dallaire, F., Dewailly, É., 2003.
- 54 696 Monitoring of umbilical cord blood lead levels and sources assessment among the
- 55 697 Inuit. *Occup. Environ. Med.* 60, 693. <https://doi.org/10.1136/oem.60.9.693>
- 56
- 57
- 58
- 59
- 60 698 Lim, K.X., Liu, C.-Y., Schoeler, T., Cecil, C.A.M., Barker, E.D., Viding, E., Greven,
- 61
- 62
- 63
- 64
- 65

- 699 C.U., Pingault, J.-B., 2018. The role of birth weight on the causal pathway to child
700 and adolescent ADHD symptomatology: a population-based twin differences
701 longitudinal design. *J. Child Psychol. Psychiatry.* 59, 1036–1043.
702 <https://doi.org/10.1111/jcpp.12949>
- 703 Little, R.J.A., 1988. A Test of Missing Completely at Random for Multivariate Data with
704 Missing Values. *J. Am. Stat. Assoc.* 83, 1198–1202.
705 <https://doi.org/10.1080/01621459.1988.10478722>
- 706 MacKinnon, D.P., Lockwood, C.M., Brown, C.H., Wang, W., Hoffman, J.M., 2007. The
707 intermediate endpoint effect in logistic and probit regression. *Clin. Trials* 4, 499–
708 513. <https://doi.org/10.1177/1740774507083434>
- 709 Malakieh, J., 2018. Adult and youth correctional statistics, 2016/2017.
- 710 Mcgue, M., Iacono, W.G., Legrand, L.N., Malone, S., Elkins, I., 2001. Origins and
711 Consequences of Age at First Drink. I. Associations With Substance- Use
712 Disorders, Disinhibitory Behavior and Psychopathology, and P3 Amplitude.
713 *Alcohol. Clin. Exp. Res.* 25, 1156–1165. [https://doi.org/10.1111/j.1530-](https://doi.org/10.1111/j.1530-0277.2001.tb02330.x)
714 [0277.2001.tb02330.x](https://doi.org/10.1111/j.1530-0277.2001.tb02330.x)
- 715 Mordre, M., Groholt, B., Kjelsberg, E., Sandstad, B., Myhre, A.M., 2011. The impact of
716 ADHD and conduct disorder in childhood on adult delinquency: A 30 years follow-
717 up study using official crime records. *BMC Psychiatry* 11, 57.
718 <https://doi.org/10.1186/1471-244X-11-57>
- 719 Muckle, G., Ayotte, P., Dewailly, E., Jacobson, S.W., Jacobson, J.L., 2001. Prenatal
720 Exposure of the Northern Quebec Inuit Infants to Environmental Contaminants.
721 *Environ. Health Perspect.* 109, 1291–1299. <https://doi.org/10.2307/3454753>
- 722 Muthén, L.K., Muthén, B.O., 2017. *Mplus User's Guide*, Eighth Edi. ed. Los Angeles,
723 CA.
- 724 Naicker, N., Richter, L., Mathee, A., Becker, P., Norris, S.A., 2012. Environmental lead
725 exposure and socio-behavioural adjustment in the early teens: The birth to twenty
726 cohort. *Sci. Total Environ.* 414, 120–125.

727 <https://doi.org/10.1016/j.scitotenv.2011.11.013>
 728 Needleman, H.L., McFarland, C., Ness, R.B., Fienberg, S.E., Tobin, M.J., 2002. Bone
 729 lead levels in adjudicated delinquents: A case control study. *Neurotoxicol. Teratol.*
 730 24, 711–717. [https://doi.org/10.1016/S0892-0362\(02\)00269-6](https://doi.org/10.1016/S0892-0362(02)00269-6)
 731 Pelham, W.E., Gnagy, E.M., Greenslade, K.E., Milich, R., 1992. Teacher Ratings of
 732 DSM-III-R Symptoms for the Disruptive Behavior Disorders. *J. Am. Acad. Child*
 733 *Adolesc. Psychiatry* 31, 210–218. [https://doi.org/10.1097/00004583-199203000-](https://doi.org/10.1097/00004583-199203000-00006)
 734 00006
 735 R Core Team, 2018. R: A Language and Environment for Statistical Computing.
 736 Raven, JC, Court, J., Raven, J, 1992. Manual for Raven’s Progressive Matrices and
 737 Vocabulary Scales: Standard Progressive Matrices, 192nd ed. Oxford.
 738 Reef, J., Diamantopoulou, S., Van Meurs, I., Verhulst, F.C., Van Der Ende, J., 2011.
 739 Developmental trajectories of child to adolescent externalizing behavior and adult
 740 DSM-IV disorder: Results of a 24-year longitudinal study. *Soc. Psychiatry*
 741 *Psychiatr. Epidemiol.* 46, 1233–1241. <https://doi.org/10.1007/s00127-010-0297-9>
 742 Revelle, W., 2018. psych: Procedures for Personality and Psychological Research,
 743 Northwestern University. Northwest. Univ. Evanston, Illinois, USA.
 744 Rothman, K.J., Greenland, S., Lash, T.L., 2008. Modern Epidemiology, 3rd ed.
 745 Lippincott Williams & Wilkins, Philadelphia.
 746 Shaffer, D., Fisher, P., Lucas, C.P., Dulcan, M.K., Schwab-Stone, M.E., 2000. NIMH
 747 Diagnostic Interview Schedule for Children Version IV (NIMH DISC-IV):
 748 Description, Differences From Previous Versions, and Reliability of Some Common
 749 Diagnoses. *J. Am. Acad. Child Adolesc. Psychiatry* 39, 28–38.
 750 <https://doi.org/10.1097/00004583-200001000-00014>
 751 Solarig, C.D., Mare, R.D., 2012. HOUSING CROWDING EFFECTS ON CHILDREN’S
 752 WELLBEING. *Soc Sci Res* 41, 464–476.
 753 <https://doi.org/10.1016/j.ssresearch.2011.09.012>

- 1
2
3
4 754 Sorenson, S.B., 2011. Gender disparities in injury mortality: consistent, persistent, and
5
6 755 larger than you'd think. *Am. J. Public Health* 101 Suppl, S353–S358.
7
8 756 <https://doi.org/10.2105/AJPH.2010.300029>
9
10 757 Téllez-Rojo, M.M., Bellinger, D.C., Arroyo-Quiroz, C., Lamadrid-Figueroa, H.,
11
12 758 Mercado-García, A., Schnaas-Arrieta, L., Wright, R.O., Hernández-Avila, M., Hu,
13
14 759 H., 2006. Longitudinal associations between blood lead concentrations lower than
15
16 760 10 microg/dL and neurobehavioral development in environmentally exposed
17
18 761 children in Mexico City. *Pediatrics* 118, e323–e330.
19
20 762 <https://doi.org/10.1542/peds.2005-3123>
21
22 763 Turner, A., 1897. Lead poisoning among Queensland Children. *Australas. Med. Gaz.* 16,
23
24 764 475–479.
25
26 765 Winter, A.S., Sampson, R.J., 2017. From lead exposure in early childhood to adolescent
27
28 766 health: A Chicago birth cohort. *Am. J. Public Health* 107, 1496–1501.
29
30 767 <https://doi.org/10.2105/AJPH.2017.303903>
31
32 768 Wright, J.P., Dietrich, K.N., Ris, M.D., Hornung, R.W., Wessel, S.D., Lanphear, B.P.,
33
34 769 Ho, M., Rae, M.N., 2008. Association of Prenatal and Childhood Blood Lead
35
36 770 Concentrations with Criminal Arrests in Early Adulthood. *PLoS Med.* 5, 732–740.
37
38 771 <https://doi.org/10.1371/journal.pmed.0050101>
39
40
41 772
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Supplemental Material - Analytical procedures for contaminant and nutrient analyses

Total mercury (Hg) concentrations in umbilical cord blood samples were determined using cold vapour atomic absorption spectrometry (Pharmacia Model 120; Pharmacia, Piscataway, NJ, USA). Cord blood lead (Pb) levels were determined by graphite furnace atomic absorption with Zeeman background correction (Perkin Elmer model ZL 4100; Perkin Elmer, Norwalk, CT, USA). Concentrations of docosahexaenoic acid (DHA) in cord blood were expressed as percentages of the total area of all fatty acid peaks from C14:0 to C24:1 (percent weight). All sample were above the limit of detection for Hg, Pb and DHA.

Total Pb and Hg concentrations in child blood samples were determined by ICP-MS (Perkin Elmer Sciex Elan 6000 ICP-MS instrument for Pb; PE DRC II instrument for Hg). Concentrations of DHA were expressed as percentages of the total area of all fatty acid peaks from C14:0 to C24:1 (percent weight). Limits of detection for child blood sample analyses were 0.1 µg/L for Hg, 0.002 µg/dL for Pb and 0.09 µmol/L for DHA. A value equal to half the limit of detection of the analytical method was entered in the database whenever a substance was not detected (child Hg: n = 1; other substances were detected in all samples).

All blood samples from the adolescent follow-up were analyzed using inductively coupled plasma mass spectrometry (PerkinElmer ELAN ICP-MS DRC II) at the Centre de Toxicologie, Institut National de Santé Publique du Québec (Quebec City, QC, Canada) for contaminants and at the CHU de Québec (Québec City, QC, Canada) for DHA. All samples were above the limits of detection for Pb, Hg and DHA.