Periodontal pathogens and clinical parameters in chronic periodontitis

Journal:	<i>Molecular Oral Microbiology</i>
Manuscript ID	MOM-07-19-1075.R3
Wiley - Manuscript type:	Original Article
Date Submitted by the Author:	n/a
Complete List of Authors:	<p>Boyer, Emile; INSERM, INRA, Univ Rennes 1, Univ Bretagne Loire, Nutrition Metabolisms and Cancer</p> <p>Martin, Bénédicte; Univeristé de Rennes 1, Microbiology - EA1254</p> <p>Le Gall-David, Sandrine; INSERM, INRA, Univ Rennes 1, Univ Bretagne Loire, Nutrition Metabolisms and Cancer</p> <p>Fong, Shao Bing; INSERM, INRA, Univ Rennes 1, Univ Bretagne Loire, Nutrition Metabolisms and Cancer</p> <p>Deugnier, Yves; INSERM, INRA, Univ Rennes 1, Univ Bretagne Loire, Nutrition Metabolisms and Cancer</p> <p>Bonnaure-Mallet, Martine; Université de Rennes 1, Equipe de Microbiologie UPRES-EA 1254</p> <p>Meuric, Vincent; INSERM, INRA, Univ Rennes 1, Univ Bretagne Loire, Nutrition Metabolisms and Cancer, ;</p>
Keywords:	Genetics < Oral Microbiology, Periodontal Disease, Ecology < Oral Microbiology, Dysbiosis
Summary	<p>Background: The use of Next Generation Sequencing and bioinformatics have revealed the complexity and richness of the human oral microbiota. While some species are well known for their periodontal pathogenicity, the molecular-based approaches for bacterial identification have raised awareness about new putative periodontal pathogens. Although they are found increased in case of periodontitis, there is currently a lack of data on their interrelationship with the periodontal measures.</p> <p>Methods: We processed the sequencing data of the subgingival microbiota of 75 patients with haemochromatosis and chronic periodontitis in order to characterize well-described and newly identified subgingival periodontal pathogens. We used correlation tests and statistical models to assess the association between the periodontal pathogens and mean pocket depth, and to determine the most relevant bacterial biomarkers of periodontitis severity.</p> <p>Results: Based on correlation test results, nine taxa were selected and included in the statistical models. The multiple linear regression models adjusted for systemic and periodontal clinical variables showed that mean pocket depth was negatively associated with <i>Aggregatibacter</i> and <i>Rothia</i>, and positively associated with <i>Porphyromonas</i>. Furthermore, a bacterial ratio that was previously described as a signature of dysbiosis in periodontitis, $(\%Porphyromonas + \%Treponema + \%Tannerella) / (\%Rothia + \%Corynebacterium)$, was the most significant predictor.</p> <p>Conclusion: In this specific population, we found that the best model in</p>

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

	predicting the mean pocket depth was microbial dybiosis using the dysbiosis ratio taxa formula. While further studies are needed to assess the validity of these results on the general population, such a dysbiosis ratio could be used in the future to monitor the subgingival microbiota.

Periodontal pathogens and clinical parameters in chronic periodontitis

Running Title: Dysbiosis in periodontitis

Emile Boyer¹; Bénédicte Martin¹; Sandrine Le Gall-David¹; Shao Bing Fong¹; Yves Deugnier²;
Martine Bonnaure-Mallet¹; Vincent Meuric¹

¹ INSERM, INRA, Univ Rennes 1, CHU de Rennes, Nutrition Metabolisms and Cancer, Rennes,
France

² CIC 1414, Inserm, Rennes, 35033, France

Corresponding author:

Vincent Meuric

2, avenue du Professeur Léon Bernard, 35043 Rennes, France

Telephone number: +33 2 23 23 49 19 Fax number: +33 2 23 23 49 13

E-mail: vincent.meuric@univ-rennes1.fr

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abstract

Background: The use of Next Generation Sequencing and bioinformatics have revealed the complexity and richness of the human oral microbiota. While some species are well known for their periodontal pathogenicity, the molecular-based approaches for bacterial identification have raised awareness about new putative periodontal pathogens. Although they are found increased in case of periodontitis, there is currently a lack of data on their interrelationship with the periodontal measures.

Methods: We processed the sequencing data of the subgingival microbiota of 75 patients with haemochromatosis and chronic periodontitis in order to characterize well-described and newly identified subgingival periodontal pathogens. We used correlation tests and statistical models to assess the association between the periodontal pathogens and mean pocket depth, and to determine the most relevant bacterial biomarkers of periodontitis severity.

Results: Based on correlation test results, nine taxa were selected and included in the statistical models. The multiple linear regression models adjusted for systemic and periodontal clinical variables showed that mean pocket depth was negatively associated with *Aggregatibacter* and *Rothia*, and positively associated with *Porphyromonas*. Furthermore, a bacterial ratio that was previously described as a signature of dysbiosis in periodontitis, $(\%Porphyromonas + \%Treponema + \%Tannerella) / (\%Rothia + \%Corynebacterium)$, was the most significant predictor.

Conclusion: In this specific population, we found that the best model in predicting the mean pocket depth was microbial dybiosis using the dysbiosis ratio taxa formula. While further studies are needed to assess the validity of these results on the general population, such a dysbiosis ratio could be used in the future to monitor the subgingival microbiota.

37 **Keywords**

38 Deep sequencing; potential periodontal pathogens; chronic periodontitis; dysbiosis.

For Review Only

1
2
3 39 **1 Introduction**
4
5
6 40

7
8 41 Periodontitis is a worldwide health problem and the 6th most common disease globally
9
10 42 (Frencken et al., 2017). It is a chronic inflammation produced in response to a disease-
11
12 43 associated multispecies bacterial community with intermittent episodes of remission and
13
14
15 44 relapse, characterized by alveolar bone loss. The current model of periodontitis pathogenesis
16
17 45 implies a synergistic polymicrobial community and a dysregulated host inflammatory response
18
19 46 (G. Hajishengallis & Lamont, 2012). Indeed, a dysbiosis of the complex and diverse oral
20
21 47 microbiota (Paster et al., 2001) that inhabits the sulcus is concomitant with the occurrence of a
22
23 48 periodontal pocket. This pathognomonic sign of the periodontitis can be quantified as pocket
24
25 49 depth (PD) and clinical attachment loss (CAL) (Lamont, Koo, & Hajishengallis, 2018).
26
27
28

29 50 Subgingival colonization by specific species, including the red complex bacterial species,
30
31 51 i.e., *Porphyromonas gingivalis*, *Tannerella forsythia* and *Treponema denticola*, has been reported
32
33 52 to be related to PD and bleeding as determined by DNA-DNA checkerboard analysis (Socransky,
34
35 53 Haffajee, Cugini, Smith, & Kent, 1998). *P. gingivalis* is also considered as the keystone pathogen
36
37 54 of the disease, **that is** able to disrupt the immune system allowing for dysbiosis to arise (George
38
39 55 Hajishengallis, Darveau, & Curtis, 2012; Jiao, Hasegawa, & Inohara, 2014). The development of
40
41 56 contemporary molecular methods (Next Generation of Sequencing – NGS) has expanded the list
42
43 57 of potential periodontal pathogens. **The review by** Pérez-Chaparro *et al.* identified in the literature
44
45 58 29 species and one genus putatively implicated in chronic periodontitis that have been cited **by** at
46
47 59 least two different authors (Pérez-Chaparro et al., 2014). However, these results are usually based
48
49 60 on health *versus* disease comparisons. The relationship between periodontitis measures and these
50
51 61 taxa are lacking. Using NGS data from the literature, we have previously defined a bacterial ratio
52
53
54
55
56
57
58
59
60

as a signature of periodontitis that could score the dysbiosis of the subgingival microbiota (Meuric, Le Gall-David, et al., 2017). However, its relationship with the global periodontal status remains to be tested.

We have worked on a cohort of patients with chronic periodontitis and hemochromatosis, that allowed us to show worsened periodontal measures in case of iron overload (Meuric, Lainé, et al., 2017). An early study (with 66 patients) of their subgingival microbiota showed a potential impact of iron through correlations between transferrin saturation and the abundance of *Porphyromonas* and *Treponema* (Boyer et al., 2018). Nevertheless, an extensive characterization of their microbiota (including dysbiosis) has not been performed yet, regarding occurrence, abundance and relationship of well-known and newly identified potential periodontal pathogens with periodontal measures.

The purpose of this study is three-fold. 1: to characterize all the well-described and newly identified periodontal pathogens (bacterial complexes described by Socransky *et al.* and reviewed by Pérez-Chaparro *et al.*) in the deepest pocket using the NGS in patients with chronic periodontitis (Pérez-Chaparro et al., 2014; Socransky et al., 1998). Two genera usually cited as implicated in health, *Corynebacterium* and *Rothia*, are also analyzed, as a part of the bacterial dysbiosis ratio. 2: to determine the association between the well-described, newly discovered periodontal pathogens and the periodontitis severity through continuous measures (mean PD, sites with $PD \geq 4$ mm). 3: to propose multiple regression linear models to refine the bacterial biomarkers of periodontitis severity adjusted for demographic and clinical data (including iron overload).

2 Materials and methods

Participants and collection of clinical data

The participants corresponded to a group of 75 patients with hemochromatosis and chronic periodontitis previously studied by Meuric *et al.* (Meuric, Lainé, et al., 2017). This case study series was approved by the local ethical committee (CPP Ouest V - 10/02-744). The informed consent for study participation has been obtained from all patients and all research was performed in accordance with relevant guidelines and regulations. These patients were recruited at the University Hospital of Rennes and benefited from a full-mouth periodontal examination, including PD, CAL, papillary bleeding index, gingival index, plaque score (Barnett, Ciancio, & Mather, 1980; Lobene, Weatherford, Ross, Lamm, & Menaker, 1986; Silness & Loe, 1964) and evaluation of periodontitis severity according to the case definitions from the Centers for Disease Control and Prevention and the American Association of Periodontology (CDC/AAP) (Eke, Page, Wei, Thornton-Evans, & Genco, 2012). For the 2018 classification, the algorithms considered the interdental CAL on two non-adjacent teeth and tooth loss (Graetz et al., 2019; Tonetti, Greenwell, & Kornman, 2018). The exclusion criteria were pregnancy, periodontal therapy within the last 12 months and treatment with either systemic antibiotics or drugs known to cause gingival hyperplasia within the last 3 months. Clinical data recorded at the time of dental examination included smoking status, alcohol consumption (non-drinkers, former drinkers, occasional drinkers and regular drinkers evaluated quantitatively in dose/day), number of dental visits per year and transferrin saturation. The assays of serum iron and transferrin were performed on an automated analyzer AU2700 Olympus (Beckman Coulter). Serum iron levels were measured by a standard colorimetric method that used TPTZ[2,4,6-Tri-(2-pyridyl)-5-triazine] as a chromogen and serum transferrin concentrations were quantified by immunoturbidimetry. Test procedures were conducted as described in the manufacturer's standard operating manual. Total iron binding capacity (TIBC) was calculated using the formula: (transferrin in g/l x 25) and transferrin saturation is calculated as follows: (iron in $\mu\text{mol/l}$ / TIBC) $\times$ 100.

110 Sample collection

111 After supra-gingival plaque removal, sterile endodontic paper points (Henry Schein, France) were
112 inserted into one of the deepest periodontal pockets for 30s. The material was transferred into a
113 sterile tube with 100 µL of sterile distilled water and kept at 4°C overnight before DNA
114 extraction.

115 DNA extraction and sequencing

116 DNA extraction from supernatants was performed using QIAamp DNA Mini Kit (Qiagen,
117 France) according to the manufacturer's recommendations. DNA was then kept frozen at -80°C
118 until amplification. The V3-V4 regions of the 16S rRNA gene were amplified with the primers
119 338F (5'-ACTCCTACGGGAGGCAGCAG-3') (Huse et al., 2008) and 802R (5'-
120 TACNVGGGTATCTAATCC-3') (Claesson et al., 2009) using 25 amplification cycles with an
121 annealing temperature of 45°C. PCR products were sequenced with the Illumina MiSeq of the
122 Get-PlaGe platform (Toulouse, France). The sequence data were submitted to the NCBI
123 Sequence Read Archive ([https://www.ncbi.nlm.nih.gov/Traces/](https://www.ncbi.nlm.nih.gov/Traces/study/) study/) under BioProject
124 accession number PRJNA416501 and PRJNA517671.

125 Microbiological analysis

126 Taxonomy at the genus level was assigned using the "Visualization and Analysis of Microbial
127 Population Structures" (VAMPS) analysis pipeline web tool (Huse et al., 2014). Default
128 parameters were used for assignment to the genus level with the Ribosomal Database Project
129 (RDP) classification (Cole et al., 2014). Through the Global Alignment for Sequence Taxonomy
130 (GAST) process of VAMPS, the best taxonomic hit was assigned for each read. Reads identified
131 as Archaea, Eukarya, Organelle and unknown were excluded from further analysis. Taxonomy at

the species level was assigned using the "Quantitative Insights Into Microbial Ecology" (QIIME 1.9.1) package software (Caporaso et al., 2010) with the default parameters (script: pick_open_reference_otus.py, uclust as the clustering method) and the Human Oral Microbiome Database (HOMD, version 15.1) as the reference database (Chen et al., 2010). The mean unassigned reads per sample was 6.4 % ± 6.2 %. These reads have been excluded from further species analyses. To challenge the assignment capacities used in this study, the V3V4 sequences of each taxon were extracted from the HOMD and entered in the previously described pipeline. Only three species (*Mogibacterium timidum*, *Anaeroglobus geminatus* and *Enterococcus faecalis*) could not be correctly classified by the pipeline, among all the putative periodontal pathogens described.

Statistics

Data were studied using R (version 3.3.3) (R. Core Team, 2015) with the RStudio software (RStudio Team, 2016). Statistical tests were considered significant for $p < 0.05$. Results of the descriptive analysis are presented as mean ± standard error (SE) and with prevalence (%). Distribution of continuous variables was assessed using the Shapiro-Wilk test. The mean PD was analyzed according to the demographic and clinical data using the Mann-Whitney test. Correlations between taxa relative abundance and clinical parameters were calculated using Spearman correlation coefficient. Multiple linear regression models were used to study the association between periodontal parameters (i.e. mean PD and percentage of site with PD ≥ 4mm) and bacterial taxa presenting a significant correlation ($r \geq 0.20$, $p < 0.05$) or the dysbiosis ratio ($(\% \textit{Porphyromonas} + \% \textit{Treponema} + \% \textit{Tannerella}) / (\% \textit{Rothia} + \% \textit{Corynebacterium})$) described in Meuric et al. 2017 (Meuric, Le Gall-David, et al., 2017), adjusted for confounding factors including age, gender, smoking status, alcohol intake, plaque index, transferrin saturation,

gingival and bleeding index. The stepwise AIC algorithm was performed using the ‘MASS’ package for R (Venables & Ripley, 2002). Final regression coefficients, standard errors (SE) and statistics for each variable adjusted for confounding factors, with the characteristics of the models (adjusted R^2 , p -value) are presented.

3 Results

- Clinical data

Based on the CDC/AAP case definition (Eke et al., 2012), all patients presented periodontitis; 44.3 % and 54.4 % had moderate and severe periodontitis, respectively. On average, 50.0 ± 1.96 % of sites per mouth had $CAL \geq 3$ mm, 10.7 ± 1.43 % of sites had $CAL \geq 5$ mm, and 12.5 ± 1.15 % of sites had $PD \geq 4$ mm. The full description of the periodontal measures is available in Table S1. According to the 2018 classification, and using CAL and tooth loss as the main available criteria, 3/75 (4%) patients were classified as stage II, 59/75 (79%) as stage III and 13/75 (17%) as stage IV (Tonetti et al., 2018). A minority (9/75, 12%) was classified as localized. The average age was 50.6 ± 8 years. Males represented 56 %, and former or current smokers and drinkers were 62.6 % and 80 %, respectively. Most of the patients visited their dentist at least once a year (78.6 %). Table 1 presents the statistics exploring the potential impact of each demographic on the mean PD. The mean PD was increased in former or current drinkers, compared to patients who had never experienced alcohol abuse ($p=0.028$).

- Prevalence and proportion of bacteria

The average of reads per sample was $14\,980 \pm 740$ (minimum = 5 298 reads). From the 74 genera and species related to periodontitis and described by Socransky *et al.*, Pérez-Chaparro *et al.* and

Meuric *et al.*, 47 taxa (64.4 %) have been identified in this study (Table 2). All the bacteria belonging to the orange and red complexes described by Socransky *et al.* were found. The lowest prevalence was *Fusobacterium periodonticum* (6.4 %), while *Fusobacterium nucleatum* and *T. forsythia* were found in all patients. For the 8 species which have been reported as highly implicated in periodontitis (reported by ≥ 4 studies according to the systematic review of the literature by Pérez-Chaparro *et al.*), five of them were found: i.e. *Porphyromonas endodontalis* (prevalence = 88.6 %), *Treponema lecithinolyticum* (62 %), *Treponema medium* (55.7 %), *Filifactor alocis* (40.5 %) and *Eubacterium saphenum* (11.4 %), while two were not detected (*Fretibacterium OT360* and *Selenomonas sputigena*). For the taxa cited 3 times, the following were not detected: *Bacteroidales OT274*, *Peptostreptococcus stomatis*, *Desulfobulbus OT041* and phylum TM7. Species with lower evidence of implication with periodontitis (cited only by two studies) or belonging to the purple or green complex (*Actinomyces odontolyticus*, *Veillonella parvula* and *Eikenella corrodens* respectively) were not, or hardly detected (Table 2). *Mogibacterium timidum*, *Anaeroglobus geminatus* and *Enterococcus faecalis* were not found due to methodological limitations.

The most abundant genus was *Fusobacterium* (mean = 26.1 ± 1.7 %) followed by *Prevotella* (12 ± 1 %), *Porphyromonas* (7 ± 0.8 %), *Streptococcus* (5.8 ± 0.7 %), *Treponema* (5.6 ± 0.6 %) and *Tannerella* (5.4 ± 0.6 %). The most abundant species were *F. nucleatum* (15.3 ± 1.4 %) followed by *T. forsythia* (4.5 ± 0.5 %), *P. endodontalis* (3.1 ± 0.4 %), and *P. gingivalis* (2.8 ± 0.6 %).

- Microbiological correlation with periodontal data

Analysis showed that the species exhibiting significant correlations with periodontal clinical data were *P. gingivalis* ($r=0.33$ and 0.41 , with mean PD and percentage of sites with $PD \geq 4$ mm, respectively), *T. forsythia* ($r=0.25$ and 0.30 , respectively), and *E. nodatum* ($r=0.35$ and 0.34 ,

respectively). *Desulfobulbus*, *Treponema* and *Porphyromonas* genera were also found significantly correlated with these periodontal data. Significant inverse correlations were found for the taxa of *Campylobacter*, *Aggregatibacter* and *Rothia*. No correlation was found between the sampled pocket depth and any of the bacterial taxa. Finally, the calculated dysbiosis ratio corresponding to $(\%Porphyromonas + \%Treponema + \%Tannerella) / (\%Rothia + \%Corynebacterium)$ showed the highest correlation with mean PD ($r=0.37$, $p=0.001$) and with the percentage of sites with $PD \geq 4$ mm ($r=0.35$, $p=0.0017$), as compared with individual taxa. Table 3 presents all the significant results of the correlation between bacterial/clinical data and mean PD, which were further used in the following statistical experiments. As shown in Table 3, the periodontal indices rating the plaque, gingival inflammation and bleeding on probing, were found significantly correlated with the mean PD. Therefore, they were also used in the statistical models as confounding variables.

- Multiple linear regression analyses

The multiple linear regression was performed with the stepwise method which selects the most significant variables affecting the mean pocket depth. After adjustment for age, sex, smoking status, alcohol status, plaque index, transferrin saturation, bleeding index and gingival index, the mean PD was found independently associated with *Aggregatibacter*, *Rothia* and *Porphyromonas* genera ($\beta=-0.053$, -0.063 and 0.051 , respectively, $p<0.05$) in the first model (adjusted $R^2 = 0.3798$, $p<0.001$). The most significant data were obtained in the second model using the dysbiosis ratio and *Aggregatibacter* as independent variables with $\beta=0.062$ ($p<0.001$) and $\beta=-0.053$ ($p<0.01$) respectively and an adjusted $R^2 = 0.4868$ ($p<0.00001$) (Table 4). By using another

periodontal measure as the outcome variable – the percentage of sites with $PD \geq 4$ mm –, the dysbiosis ratio ($p < 0.001$ – Table S2) remained as the most significant predictor.

4 Discussion

Owing to decades of oral microbiology studies, some species are well known for their periodontal pathogenicity. Culture-independent and molecular-based approaches to bacterial identification has highlighted new putative periodontal pathogens, that were found increased in periodontitis patients. However, few studies assess the correlation between the severity of periodontal disease and the abundance of these taxa. The identification of taxa that are significantly associated with the periodontal status is the first step in the search for biomarkers that could predict the onset of the disease.

Most of the periodontal pathogens were identified in the analyses due to the sequencing depth. However, six species were not detected. *Bacteroidales* OT274, *Peptostreptococcus stomatis*, *Desulfobulbus* OT041, *Fretibacterium* sp., TM7 and *Selenomonas Sputigena* could not be found while they are highly cited (≥ 3 times) in the literature as being implicated in periodontitis. Only three (*Mogibacterium timidum*, *Anaeroglobus geminatus* and *Enterococcus faecalis*) could not be identified through the pipeline of assignation, due to methodological limitations. The lack of detection of *Fretibacterium* is surprising since it was proposed as a diagnostic bacterial biomarker for periodontitis screening in a Japanese population (Khemwong et al., 2019). Indeed, *Fretibacterium* was not detected in our Caucasian population. As for the other five remaining taxa, we have confirmed that they were not detected in this population. Thus, different population could carry different taxon associated with periodontitis. However, previous works on periodontitis microbiota analysis through NGS have used a binary approach (healthy or disease) to determine in which condition taxon are increased (Abusleme et al., 2013;

Kirst et al., 2015). In this study, we have performed correlation analyses between the abundance of species considered as periodontal pathogens (from previous studies) and continuous periodontal measures, in haemochromatosis patients with periodontitis to assess the current weight of their implication in the disease. Adjusted regression models were then used to determine the best bacterial biomarkers of periodontitis severity. Taxa already well described as implicated in the disease were found correlated with the mean PD or the percentage of sites with $PD \geq 4$ mm: *P. gingivalis* and *T. forsythia* (red complex) and *E. nodatum* (orange complex) (Socransky et al., 1998). From the newly identified putative periodontal pathogen, only *Desulfobulbus* genus (Pérez-Chaparro et al., 2014) was found correlated with the mean pocket depth. Correlation coefficients between periodontal data and bacterial taxa remained inferior to 0.4, similarly to other microbiota analyses in chronic periodontitis (Khemwong et al., 2019; Lourenço et al., 2014; Silva-Boghossian, Cesário, Leão, & Colombo, 2018). Finding associations is especially more difficult when looking at the species level. However, it is not surprising as this disease is a multifactorial pathology, as each taxon might be partially implicated in the disease. For instance, *T. denticola* which belongs to the red complex (Socransky et al., 1998), was not correlated with the mean PD. Nevertheless, the *Treponema* genus, which contains *T. denticola* and other species highly suspected to be periodontal pathogens (i.e. *T. vincentii*, *T. lecithinolyticum* and *T. medium*, as reported in the systematic review by Pérez-Chaparro et al.), was found correlated with the PD measures.

Surprisingly, *Campylobacter* was negatively associated with the mean PD, while *Campylobacter* species are mainly classified as periodontal pathogens (i.e.: *C. showae*, *C. gracilis*, and *C. rectus* belonging to the orange complex and *C. concisus* to the green complex (Socransky et al., 1998)). However, this repartition of species is controversial: *C. gracilis* belongs to the orange complex, though it was found increased in healthy subjects (Henne, Fuchs, Kruth,

Horz, & Conrads, 2014) and decreased in severe periodontitis (Macuch & Tanner, 2000). Finally, the *Campylobacter* genus was still negatively associated with percentage of PD $\geq$ 4mm in the regression model.

This analytical approach is also interesting when focused on the prevalence or occurrence of periodontal pathogens in patients. Taxa are not found in all patients. Even the prevalence of *P. gingivalis*, which is well described as one of the keystone pathogens in the disease (George Hajishengallis *et al.*, 2012), was only found at 62% in this study. Moreover, *P. gingivalis* has been detected in 25% of healthy Caucasian (Griffen, Becker, Lyons, Moeschberger, & Leys, 1998), up to 40% in healthy Japanese (Kato, Imai, Ochiai, & Ogata, 2013) and not higher than 79% in chronic periodontitis (Griffen *et al.*, 1998). The chronic nature and clinical course of the disease, with alternating episodes of exacerbation and remission (Goodson, Tanner, Haffajee, Sornberger, & Socransky, 1982), could explain that the bacteria are more difficult to detect. The single sampling could also be a potential modifier of the microbiota and a limitation, as different periodontal pockets in the same patient could show various microbiota composition. Long-term follow-up with regular and multi-site sampling is certainly needed, which could address these issues. As such studies are difficult to implement, a ratio of taxa was developed (Meuric, Le Gall-David, *et al.*, 2017) to explore the periodontal dysbiosis responsible for chronic periodontitis (Kilian *et al.*, 2016). This dysbiosis ratio showed the highest and the most significant correlation with mean PD in this study ($p=0.001$) and could be calculated for each patient, whereas *P. gingivalis* could only be detected in 62% of the patients. Thereafter, in the first multiple regression model, each taxon found significantly correlated with mean PD was entered, while *Porphyromonas*, *Treponema*, *P. gingivalis*, *T. forsythia* and *Rothia* were replaced in the second model by the dysbiosis ratio which is computed with their relative abundance. The models were

refined with the stepwise method (Venables & Ripley, 2002) adjusted with the clinical and demographic data as for example, alcohol intake, which is considered as a risk factor for periodontitis (Wang, Lv, Wang, & Jiang, 2016) and a worsened mean PD in our population. Since the patients come from a sub-cohort of a previous study where high transferrin saturation (specifically for the hemochromatosis disease) was implicated in explaining clinical severity of periodontitis, we have added the transferrin saturation into the multiple linear regression models as a confounding factor. As described previously, periodontitis through mean PD was associated with transferrin saturation in the two models (Meuric, Lainé, et al., 2017). For each dependent variable (mean PD or percentage of sites with $PD \geq 4$ mm), the most significant result, appeared in the second model with the dysbiosis ratio ($p < 0.001$) which is compiled with majority of the periopathogens (i.e.: *Porphyromonas*, *Treponema* and *Tannerella* genera) and the genera usually found in health (Meuric, Le Gall-David, et al., 2017). This result is independent from the transferrin saturation (reflecting the iron status of the hemochromatosis disease). Thus, the dysbiosis ratio confirmed its utility in exploring the microbial dysbiosis to explain periodontal disease (Meuric, Le Gall-David, et al., 2017). Similar microbial dysbiosis ratios have also been calculated in Crohn's disease (Gevers et al., 2014) and in peri-implantitis (Kröger et al., 2018).

These results are based on a specific population suffering from haemochromatosis. Extrapolations outside of this population need to be validated, though the iron status of the patients has been included as a confounding variable in statistical models. Nonetheless, these results are in accordance with the existing view that periodontitis or peri-implantitis are orchestrated by a bacterial consortium rather than a single pathogen (Darveau, 2010).

1
2
3 318 **5 Conclusion**
4

5 319 In this study, we have confirmed that most of the well-described and newly identified
6
7 320 periodontal pathogens can be found in the deepest periodontal pocket of patients suffering from
8
9 321 haemochromatosis and periodontitis. Yet, only few taxa were correlated to mean pocket depth.
10
11 322 The relative abundance of these taxa was therefore used as a predictor, with confounding
12
13 323 variables, in multiple linear regression models that allowed for the prediction of mean pocket
14
15 324 depth. While *Porphyromonas*, *Aggregatibacter* and *Rothia* returned significant results, the
16
17 325 previously described dysbiosis ratio – (% *Porphyromonas* + % *Treponema* + % *Tannerella*) / (%
18
19 326 *Rothia* + % *Corynebacterium*) – was the most reliable biomarker to predict periodontitis severity
20
21 327 in our patients. Longitudinal studies in patients with no systemic disease, as well as several
22
23 328 sampling sites and time points during the episodes of disease-active periodontitis should be
24
25 329 considered to further validate and generalize the findings of this study.
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Acknowledgments: The authors are indebted to the medical and nursing staff for participating in the recruitment and follow-up of patients. We thank the GeT-PlaGe facility and the NED team for sequencing the samples. We also thank Adina Pascu (INSERM U1241) for her contribution to this work and the “Direction de la Recherche et de l’Innovation” at the University hospital of Rennes for their support. The study was funded by the Comité de la Recherche Clinique of the University Hospital (2009-A00998-49) of Rennes and the University of Rennes 1. The authors report no conflicts of interest related to this study.

Data availability: The sequence data were deposited to the NCBI Sequence Read Archive (<https://www.ncbi.nlm.nih.gov/Traces/study/>) under BioProject accession numbers PRJNA416501 and PRJNA517671. The detailed periodontal and demographic data for every patient can be accessed in Table S3.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

Abusleme, L., Dupuy, A. K., Dutzan, N., Silva, N., Burleson, J. A., Strausbaugh, L. D., ... Diaz, P. I. (2013). The subgingival microbiome in health and periodontitis and its relationship with community biomass and inflammation. *The ISME Journal*, 7(5), 1016–1025. <https://doi.org/10.1038/ismej.2012.174>

Barnett, M., Ciancio, S., & Mather, M. (1980). The Modified Papillary Bleeding Index—Comparison with Gingival Index During the Resolution of Gingivitis. *Journal of Preventive Dentistry*, 6(2), 135–138.

Boyer, E., Gall-David, S. L., Martin, B., Fong, S. B., Loréal, O., Deugnier, Y., ... Meuric, V. (2018). Increased transferrin saturation is associated with subgingival microbiota dysbiosis and severe periodontitis in genetic haemochromatosis. *Scientific Reports*, 8(1), 15532. <https://doi.org/10.1038/s41598-018-33813-0>

Caporaso, J. G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F. D., Costello, E. K., ... others. (2010). QIIME allows analysis of high-throughput community sequencing data. *Nature Methods*, 7(5), 335–336.

Chen, T., Yu, W.-H., Izard, J., Baranova, O. V., Lakshmanan, A., & Dewhirst, F. E. (2010). The Human Oral Microbiome Database: A web accessible resource for investigating oral microbe taxonomic and genomic information. *Database*, 2010, baq013. <https://doi.org/10.1093/database/baq013>

Claesson, M. J., O’Sullivan, O., Wang, Q., Nikkilä, J., Marchesi, J. R., Smidt, H., ... O’Toole, P. W. (2009). Comparative Analysis of Pyrosequencing and a Phylogenetic Microarray for Exploring Microbial Community Structures in the Human Distal Intestine. *PLOS ONE*, 4(8), e6669. <https://doi.org/10.1371/journal.pone.0006669>

Cole, J. R., Wang, Q., Fish, J. A., Chai, B., McGarrell, D. M., Sun, Y., ... Tiedje, J. M. (2014). Ribosomal Database Project: Data and tools for high throughput rRNA analysis. *Nucleic Acids Research*, 42(Database issue), D633-642. <https://doi.org/10.1093/nar/gkt1244>

Darveau, R. P. (2010). Periodontitis: A polymicrobial disruption of host homeostasis. *Nature Reviews Microbiology*, 8(7), 481–490. <https://doi.org/10.1038/nrmicro2337>

Eke, P. I., Page, R. C., Wei, L., Thornton-Evans, G., & Genco, R. J. (2012). Update of the case definitions for population-based surveillance of periodontitis. *Journal of Periodontology*, 83(12), 1449–1454. <https://doi.org/10.1902/jop.2012.110664>

Frencken, J. E., Sharma, P., Stenhouse, L., Green, D., Lavery, D., & Dietrich, T. (2017). Global epidemiology of dental caries and severe periodontitis – a comprehensive review. *Journal of Clinical Periodontology*, 44(S18), S94–S105. <https://doi.org/10.1111/jcpe.12677>

Gevers, D., Kugathasan, S., Denson, L. A., Vázquez-Baeza, Y., Van Treuren, W., Ren, B., ... Xavier, R. J. (2014). The treatment-naïve microbiome in new-onset Crohn’s disease. *Cell Host &*

- 378 *Microbe*, 15(3), 382–392. <https://doi.org/10.1016/j.chom.2014.02.005>
- 379 Goodson, J. M., Tanner, A. C., Haffajee, A. D., Sornberger, G. C., & Socransky, S. S. (1982).
 380 Patterns of progression and regression of advanced destructive periodontal disease. *Journal of*
 381 *Clinical Periodontology*, 9(6), 472–481.
- 382 Graetz, C., Mann, L., Krois, J., Sälzer, S., Kahl, M., Springer, C., & Schwendicke, F. (2019).
 383 Comparison of periodontitis patients' classification in the 2018 versus 1999 classification.
 384 *Journal of Clinical Periodontology*. <https://doi.org/10.1111/jcpe.13157>
- 385 Griffen, A. L., Becker, M. R., Lyons, S. R., Moeschberger, M. L., & Leys, E. J. (1998).
 386 Prevalence of Porphyromonas gingivalis and periodontal health status. *Journal of Clinical*
 387 *Microbiology*, 36(11), 3239–3242.
- 388 Hajishengallis, G., & Lamont, R. J. (2012). Beyond the red complex and into more complexity:
 389 The polymicrobial synergy and dysbiosis (PSD) model of periodontal disease etiology.
 390 *Molecular Oral Microbiology*, 27(6), 409–419. [https://doi.org/10.1111/j.2041-](https://doi.org/10.1111/j.2041-1014.2012.00663.x)
 391 [1014.2012.00663.x](https://doi.org/10.1111/j.2041-1014.2012.00663.x)
- 392 Hajishengallis, George, Darveau, R. P., & Curtis, M. A. (2012). The keystone-pathogen
 393 hypothesis. *Nature Reviews. Microbiology*, 10(10), 717–725.
 394 <https://doi.org/10.1038/nrmicro2873>
- 395 Henne, K., Fuchs, F., Kruth, S., Horz, H.-P., & Conrads, G. (2014). Shifts in Campylobacter
 396 species abundance may reflect general microbial community shifts in periodontitis progression.
 397 *Journal of Oral Microbiology*, 6, 25874. <https://doi.org/10.3402/jom.v6.25874>
- 398 Huse, S. M., Dethlefsen, L., Huber, J. A., Welch, D. M., Relman, D. A., & Sogin, M. L. (2008).
 399 Exploring Microbial Diversity and Taxonomy Using SSU rRNA Hypervariable Tag Sequencing.
 400 *PLOS Genet*, 4(11), e1000255. <https://doi.org/10.1371/journal.pgen.1000255>
- 401 Huse, S. M., Mark Welch, D. B., Voorhis, A., Shipunova, A., Morrison, H. G., Eren, A. M., &
 402 Sogin, M. L. (2014). VAMPS: A website for visualization and analysis of microbial population
 403 structures. *BMC Bioinformatics*, 15, 41. <https://doi.org/10.1186/1471-2105-15-41>
- 404 Jiao, Y., Hasegawa, M., & Inohara, N. (2014). The Role of Oral Pathobionts in Dysbiosis during
 405 Periodontitis Development. *Journal of Dental Research*, 93(6), 539–546.
 406 <https://doi.org/10.1177/0022034514528212>
- 407 Kato, A., Imai, K., Ochiai, K., & Ogata, Y. (2013). Higher Prevalence of Epstein–Barr Virus
 408 DNA in Deeper Periodontal Pockets of Chronic Periodontitis in Japanese Patients. *PLoS ONE*,
 409 8(8), e71990. <https://doi.org/10.1371/journal.pone.0071990>
- 410 Khemwong, T., Kobayashi, H., Ikeda, Y., Matsuura, T., Sudo, T., Kano, C., ... Izumi, Y. (2019).
 411 Fretibacterium sp. Human oral taxon 360 is a novel biomarker for periodontitis screening in the
 412 Japanese population. *PloS One*, 14(6), e0218266. <https://doi.org/10.1371/journal.pone.0218266>
- 413 Kilian, M., Chapple, I. L. C., Hannig, M., Marsh, P. D., Meuric, V., Pedersen, A. M. L., ...
 414 Zaura, E. (2016). The oral microbiome – an update for oral healthcare professionals. *British*

- 415 *Dental Journal*, 221(10), 657–666. <https://doi.org/10.1038/sj.bdj.2016.865>
- 416 Kirst, M. E., Li, E. C., Alfant, B., Chi, Y.-Y., Walker, C., Magnusson, I., & Wang, G. P. (2015).
 417 Dysbiosis and alterations in predicted functions of the subgingival microbiome in chronic
 418 periodontitis. *Applied and Environmental Microbiology*, 81(2), 783–793.
 419 <https://doi.org/10.1128/AEM.02712-14>
- 420 Kröger, A., Hülsmann, C., Fickl, S., Spinell, T., Hüttig, F., Kaufmann, F., ... Kebschull, M.
 421 (2018). The severity of human peri-implantitis lesions correlates with the level of submucosal
 422 microbial dysbiosis. *Journal of Clinical Periodontology*, 45(12), 1498–1509.
 423 <https://doi.org/10.1111/jcpe.13023>
- 424 Lamont, R. J., Koo, H., & Hajishengallis, G. (2018). The oral microbiota: Dynamic communities
 425 and host interactions. *Nature Reviews. Microbiology*, 16(12), 745–759.
 426 <https://doi.org/10.1038/s41579-018-0089-x>
- 427 Lobene, R. R., Weatherford, T., Ross, N. M., Lamm, R. A., & Menaker, L. (1986). A modified
 428 gingival index for use in clinical trials. *Clinical Preventive Dentistry*, 8(1), 3–6.
- 429 Lourenço, T. G. B., Heller, D., Silva-Boghossian, C. M., Cotton, S. L., Paster, B. J., & Colombo,
 430 A. P. V. (2014). Microbial signature profiles of periodontally healthy and diseased patients.
 431 *Journal of Clinical Periodontology*, 41(11), 1027–1036. <https://doi.org/10.1111/jcpe.12302>
- 432 Macuch, P. J., & Tanner, A. C. (2000). Campylobacter species in health, gingivitis, and
 433 periodontitis. *Journal of Dental Research*, 79(2), 785–792.
 434 <https://doi.org/10.1177/00220345000790021301>
- 435 Meuric, V., Lainé, F., Boyer, E., Le Gall-David, S., Oger, E., Bourgeois, D., ... Deugnier, Y.
 436 (2017). Periodontal status and serum biomarker levels in HFE hemochromatosis patients. A case
 437 series study. *Journal of Clinical Periodontology*, 44(9), 892–897.
 438 <https://doi.org/10.1111/jcpe.12760>
- 439 Meuric, V., Le Gall-David, S., Boyer, E., Acuña-Amador, L., Martin, B., Fong, S. B., ...
 440 Bonnaure-Mallet, M. (2017). Signature of Microbial Dysbiosis in Periodontitis. *Applied and*
 441 *Environmental Microbiology*, 83(14). <https://doi.org/10.1128/AEM.00462-17>
- 442 Paster, B. J., Boches, S. K., Galvin, J. L., Ericson, R. E., Lau, C. N., Levanos, V. A., ...
 443 Dewhirst, F. E. (2001). Bacterial diversity in human subgingival plaque. *Journal of Bacteriology*,
 444 183(12), 3770–3783. <https://doi.org/10.1128/JB.183.12.3770-3783.2001>
- 445 Pérez-Chaparro, P. J., Gonçalves, C., Figueiredo, L. C., Faveri, M., Lobão, E., Tamashiro, N., ...
 446 Feres, M. (2014). Newly identified pathogens associated with periodontitis: A systematic review.
 447 *Journal of Dental Research*, 93(9), 846–858. <https://doi.org/10.1177/0022034514542468>
- 448 R. Core Team. (2015). R: A language and environment for statistical computing. *R Foundation*
 449 *for Statistical Computing, Vienna, Austria*. Retrieved from <https://www.R-project.org/>
- 450 RStudio Team. (2016). RStudio: Integrated development for R. *RStudio, Inc., Boston, MA*. URL
 451 <Http://Www.Rstudio.Com>.

- 452 Silness, J., & Loe, H. (1964). Periodontal disease in pregnancy. II. Correlation between oral
453 hygiene and periodontal condition. *Acta Odontologica Scandinavica*, 22, 121–135.
- 454 Silva-Boghossian, C. M., Cesário, P. C., Leão, A. T. T., & Colombo, A. P. V. (2018).
455 Subgingival microbial profile of obese women with periodontal disease. *Journal of*
456 *Periodontology*, 89(2), 186–194. <https://doi.org/10.1002/JPER.17-0236>
- 457 Socransky, S. S., Haffajee, A. D., Cugini, M. A., Smith, C., & Kent, R. L. (1998). Microbial
458 complexes in subgingival plaque. *Journal of Clinical Periodontology*, 25(2), 134–144.
- 459 Tonetti, M. S., Greenwell, H., & Kornman, K. S. (2018). Staging and grading of periodontitis:
460 Framework and proposal of a new classification and case definition. *Journal of Clinical*
461 *Periodontology*, 45 Suppl 20, S149–S161. <https://doi.org/10.1111/jcpe.12945>
- 462 Venables, W. N., & Ripley, B. D. (2002). *Modern Applied Statistics with S* (4th ed.). Retrieved
463 from <https://www.springer.com/gp/book/9780387954578>
- 464 Wang, J., Lv, J., Wang, W., & Jiang, X. (2016). Alcohol consumption and risk of periodontitis: A
465 meta-analysis. *Journal of Clinical Periodontology*, 43(7), 572–583.
466 <https://doi.org/10.1111/jcpe.12556>

Demographics	Mean PD (mm)	<i>p</i> -value
<i>Gender</i>		
Male (<i>n</i> = 42)	2.32 ± 0.06	0.863
Female (<i>n</i> = 33)	2.29 ± 0.06	
<i>Smoking status</i>		
Never (<i>n</i> = 28)	2.29 ± 0.04	0.81
Former (<i>n</i> = 30) and Current (<i>n</i> = 17)	2.31 ± 0.06	
<i>Alcohol status</i>		
Never (<i>n</i> = 15)	2.12 ± 0.07	0.028
Former (<i>n</i> = 2) and Current (<i>n</i> = 58)	2.35 ± 0.05	
<i>Frequency of dentist visits</i>		
Once or twice a year (<i>n</i> = 59)	2.32 ± 0.06	0.535
Less than once a year (<i>n</i> = 16)	2.27 ± 0.09	

Table 1: Mean probing depth (PD) according to demographic data. Data are presented as mean ± standard error (SE). Bold *p*-values indicate significant Mann-Whitney test.

Phylum	Genera	Species	HOT	Author	Year	Identified in	Mean % genera	Mean % species	+/-SE	Occurrence (%)
Actinobacteria	<i>Corynebacteria</i>			Meuric	2017	Health	0.143		0.028	83.5
Actinobacteria	<i>Rothia</i>			Meuric	2017	Health	0.215		0.050	73.4
Firmicutes	<i>Streptococcus</i>			Socransky	1998	yellow complex	5.782		0.728	100
Firmicutes	<i>Veillonella</i>	<i>parvula</i>	161	Socransky	1998	purple complex	ND			
Actinobacteria	<i>Actinomyces</i>						0.237		0.054	77.2
Actinobacteria	<i>Actinomyces</i>	<i>odontolyticus</i>	701	Socransky	1998	purple complex	ND			
Proteobacteria	<i>Eikenella</i>	<i>corrodens</i>	577	Socransky	1998	green complex	ND			
Bacteroidetes	<i>Capnocytophaga</i>						1.667		0.262	98.7
Bacteroidetes	<i>Capnocytophaga</i>	<i>gingivalis</i>	337	Socransky	1998	green complex		0.002	0.001	11.4
Bacteroidetes	<i>Capnocytophaga</i>	<i>sputigena</i>	775	Socransky	1998	green complex		0.509	0.087	83.5
Bacteroidetes	<i>Capnocytophaga</i>	<i>ochracea</i>	700	Socransky	1998	green complex		0.002	0.001	19.0
Bacteroidetes	<i>Campylobacter</i>						4.279		0.365	100.0
Bacteroidetes	<i>Campylobacter</i>	<i>concisus</i>	575	Socransky	1998	green complex		0.190	0.041	77.2
Bacteroidetes	<i>Porphyromonas</i>						7.054		0.833	98.7
Bacteroidetes	<i>Porphyromonas</i>	<i>gingivalis</i>	619	Socransky	1998	red complex		2.769	0.626	62.0
Spirochaetes	<i>Treponema</i>						5.591		0.584	100
Spirochaetes	<i>Treponema</i>	<i>denticola</i>	584	Socransky	1998	red complex		1.061	0.170	87.3
Bacteroidetes	<i>Tannerella</i>						5.368		0.584	100
Bacteroidetes	<i>Tannerella</i>	<i>forsythia</i>	613	Socransky	1998	red complex		4.522	0.531	100
Firmicutes	<i>Eubacterium</i>						1.415		0.200	97.6
Firmicutes	<i>Eubacterium</i>	<i>nodatum</i>	694	Socransky	1998	orange complex		0.177	0.043	62.0
Fusobacteria	<i>Fusobacterium</i>						26.140		1.733	100
Fusobacteria	<i>Fusobacterium</i>	<i>nucleatum</i>	200. 202. 420. 698	Socransky	1998	orange complex		15.292	1.458	100
Fusobacteria	<i>Fusobacterium</i>	<i>periodonticum</i>	201	Socransky	1998	orange complex		0.0003	0.0002	6.3
Firmicutes	<i>Parvimonas</i>						1.019		0.186	92.4
Firmicutes	<i>Parvimonas</i>	<i>micra</i>	111	Socransky	1998	orange complex		1.015	0.176	91.1
Bacteroidetes	<i>Prevotella</i>						12.006		1.043	100
Bacteroidetes	<i>Prevotella</i>	<i>intermedia</i>	643	Socransky	1998	orange complex		0.774	0.228	58.2
Bacteroidetes	<i>Prevotella</i>	<i>nigrescens</i>	693	Socransky	1998	orange complex		1.558	0.290	82.3
Firmicutes	<i>Streptococcus</i>	<i>constellatus</i>	576	Socransky	1998	orange complex		0.789	0.177	82.3
Proteobacteria	<i>Campylobacter</i>	<i>gracilis</i>	623	Socransky	1998	orange complex		1.363	0.202	92.4
Proteobacteria	<i>Campylobacter</i>	<i>rectus</i>	748	Socransky	1998	orange complex		1.740	0.240	94.9
Proteobacteria	<i>Campylobacter</i>	<i>showae</i>	763	Socransky	1998	orange complex		0.012	0.002	53.2
Proteobacteria	<i>Aggregatibacter</i>						0.851		0.167	86.1
Proteobacteria	<i>Aggregatibacter</i>	<i>actinomycetemcomitans</i>	531	Socransky	1998			0.027	0.012	17.7
Bacteroidetes	<i>Prevotella</i>	<i>denticola</i>	291	Perez-Chapparo	2014	2 studies		0.868	0.226	75.9
Bacteroidetes	<i>Alloprevotella</i>	<i>tanneriae</i>	466	Perez-Chapparo	2014	2 studies	ND			

1				124. 125. 126. 130. 133. 134.								
2				136. 137. 138. 139. 146. 149.								
3				388. 442. 478. 479. 481. 501.								
4	Firmicutes	<i>Selenomonas</i>	genus	639. 892. 919. 920. 936. 937	Perez-Chapparo	2014	2 studies	ND				
5												
6	Firmicutes	<i>Johnsonella</i>	oral taxon 166	166	Perez-Chapparo	2014	2 studies	ND				
7	Firmicutes	<i>Eubacterium</i> [X1] [G-3]	brachy	557	Perez-Chapparo	2014	2 studies		0.005	0.001	30.4	
8	Firmicutes	<i>Peptostreptococcaceae</i> [XIII] [G-1]	oral taxon 113	113	Perez-Chapparo	2014	2 studies	ND				
9	Firmicutes	<i>Lachnospiraceae</i> [G-8]	oral taxon 500	500	Perez-Chapparo	2014	2 studies	ND				
10	Firmicutes	<i>Dialister</i>						1.518		0.197	94.9	
11	Firmicutes	<i>Dialister</i>	pneumosintes	736	Perez-Chapparo	2014	2 studies	ND				
12	Proteobacteria	<i>Acinetobacter</i>						0.081		0.030	45.6	
13	Proteobacteria	<i>Acinetobacter</i>	baumannii	554	Perez-Chapparo	2014	2 studies	ND				
14	Proteobacteria	<i>Escherichia</i>						0.065		0.016	55.7	
15	Proteobacteria	<i>Escherichia</i>	coli	574	Perez-Chapparo	2014	2 studies	ND				
16	Spirochaetes	<i>Treponema</i>	phylogroup II	584. 743	Perez-Chapparo	2014	2 studies		1.106	0.194	86.7	
17	Synergistetes	<i>Fretibacterium</i>	oral taxon 359	359	Perez-Chapparo	2014	2 studies	ND				
18	TM7	TM7 [G-1]	oral taxon 346. 349	346. 349	Perez-Chapparo	2014	2 studies	ND				
19	SR1	SR1 [G-1]	oral taxon 345	345	Perez-Chapparo	2014	2 studies	ND				
20	Bacteroidetes	<i>Bacteroidales</i> [G-2]	oral taxon 274	274	Perez-Chapparo	2014	3 studies	ND				
21	Firmicutes	<i>Mogibacterium</i>						0.293		0.037	93.7	
22	Firmicutes	<i>Mogibacterium</i>	timidum	0 42	Perez-Chapparo	2014	3 studies	ND				
23	Firmicutes	<i>Peptostreptococcus</i>						0.192		0.040	77.2	
24	Firmicutes	<i>Peptostreptococcus</i>	stomatis	112	Perez-Chapparo	2014	3 studies	ND				
25	Firmicutes	<i>Anaeroglobus</i>	geminatus	121	Perez-Chapparo	2014	3 studies	ND				
26	Proteobacteria	<i>Desulfobulbus</i>						1.515		0.285	81.0	
27	Proteobacteria	<i>Desulfobulbus</i>	sp. oral taxon 041	0 41	Perez-Chapparo	2014	3 studies	ND				
28	Spirochaetes	<i>Treponema</i>	vincentii	0 29	Perez-Chapparo	2014	3 studies		0.034	0.009	41.8	
29	Synergistetes	<i>Fretibacterium</i>	oral taxon 362	362	Perez-Chapparo	2014	3 studies	ND				
30	Synergistetes	<i>Fretibacterium</i>	fastidiosum	363	Perez-Chapparo	2014	3 studies	ND				
31	TM7	TM7 [G-5]	oral taxon 356	356	Perez-Chapparo	2014	3 studies	ND				
32	Bacteroidetes	<i>Porphyromonas</i>	endodontalis	273	Perez-Chapparo	2014	4 studies		3.105	0.421	88.6	
33	Firmicutes	<i>Enterococcus</i>	faecalis	604	Perez-Chapparo	2014	4 studies	ND				
34	Spirochaetes	<i>Treponema</i>	lecithinolyticum	653	Perez-Chapparo	2014	4 studies		0.696	0.158	62	
35	Synergistetes	<i>Fretibacterium</i>	oral taxon 360	360	Perez-Chapparo	2014	4 studies	ND				
36	Firmicutes	<i>Eubacterium</i> [XI] [G-5]	saphenum	759	Perez-Chapparo	2014	5 studies		0.002	0.001	11.4	
37	Firmicutes	<i>Filifactor</i>						2.202		0.305	97.5	
38	Firmicutes	<i>Filifactor</i>	alocis	539	Perez-Chapparo	2014	5 studies		0.041	0.011	40.5	
39	Firmicutes	<i>Selenomonas</i>	sputigena	151	Perez-Chapparo	2014	5 studies	ND				
40	Spirochaetes	<i>Treponema</i>	medium	667	Perez-Chapparo	2014	5 studies		0.044	0.010	55.7	
41	Dysbiosis Ratio	(% Porphyromonas + % Treponema + % Tannerella) / (% Corynebacterium + % Rothia)			Meuric	2017		Mean %	+/-SE		Occurrence	
42								61 816	+/- 19 257		100	

Table 2: Taxonomy and dysbiosis ratio (PTT/RC) implicated in chronic periodontitis, with the mean, standard error (SE) and occurrence of the bacterial taxa and of the dysbiosis ratio in the present study. ND: not detected

Correlation with mean PD	Rho	p-value	Mean ± SE	Occurrence (%)
<i>Bacterial data</i>				
Dysbiosis ratio	0.37	0.001	3 573 ± 202	100
<i>E. nodatum</i>	0.35	0.01	0.18 ± 0.04	62
<i>P. gingivalis</i>	0.33	0.01	2.77 ± 0.63	62
<i>Desulfobulbus</i>	0.28	0.05	1.52 ± 0.29	81
<i>Treponema</i>	0.26	0.05	5.59 ± 0.58	100
<i>T. forsythia</i>	0.25	0.05	4.52 ± 0.53	100
<i>Porphyromonas</i>	0.23	0.05	7.05 ± 0.83	98.7
<i>Aggregatibacter</i>	-0.31	0.01	0.85 ± 0.17	86.1
<i>Campylobacter</i>	-0.27	0.05	4.28 ± 0.36	100
<i>Rothia</i>	-0.26	0.05	0.22 ± 0.05	73.4
<i>Clinical data</i>				
Plaque index	0.25	0.03	0.19 ± 0.03	
Gingival index	0.35	0.002	0.41 ± 0.03	
Papillary bleeding index	0.36	0.002	0.4 ± 0.03	

Table 3: Correlation between mean probing depth (PD) and bacterial or clinical data. Only significant correlations are presented.

Dysbiosis ratio: $(\%Porphyromonas + \%Treponema + \%Tannerella) / (\%Rothia + \%Corynebacterium)$; SE: standard error. Plaque index from Silness & Loe. 1964, Gingival index from Lobene *et al.* 1986 and Papillary bleeding index from Barnett *et al.* 1980.

Coefficients:	Estimate	Std. Error	t value	p-value
Model 1				
(Intercept)	1.484	0.221	6.730	3.54e-08
Transferrin saturation (binary)	0.198	0.076	2.612	0.05
Plaque index	-0.049	0.195	-0.252	NS
Papillary bleeding index	0.301	0.161	1.872	NS
log(<i>Aggregatibacter</i>)	-0.053	0.021	-2.492	0.05
log(<i>Rothia</i>)	-0.063	0.025	-2.482	0.05
log(<i>Porphyromnas</i>)	0.051	0.025	2.035	0.05
Model 2				
(Intercept)	1,280	0.147	8.686	4.39e-12
Log(Dysbiosis ratio)	0.062	0,016	3.791	0.001
Log(<i>Aggregatibacter</i>)	-0.053	0.018	-2.895	0.01
Papillary bleeding index	0.349	0.128	2.732	0.01
Transferrin saturation (binary)	0.159	0.069	2.309	0.05
Male	0.171	0.069	2.457	0.05
Alcohol	0.121	0.088	1.373	NS

Table 4: Multiple linear adjusted regression models with bacteria positively associated before or dysbiosis ratio and *Aggregatibacter* as predictors of **mean periodontal pocket depth**, adjusted for age, sex, smoking status, alcohol intake, transferrin saturation, plaque index, gingival index and bleeding index.

Characteristics of the model 1: multiple $R^2 = 0.4573$, adjusted $R^2 = 0.3798$, F-statistic = 5.899, P -value = 0.0001569. (42 degrees of freedom)

Characteristics of the model 2: multiple $R^2 = 0.5349$, adjusted $R^2 = 0.4868$, F-statistic = 11.12, P -value = 3.219e-08. (58 degrees of freedom)