Looking into each other's eyes makes it better: eye-to-eye contact enhances sexual interactions in wild geladas

Anna Zanolli, Marco Gamba, Alban Lemasson, Elisabetta Palagi, Ivan Norscia

► To cite this version:

Anna Zanolli, Marco Gamba, Alban Lemasson, Elisabetta Palagi, Ivan Norscia. Looking into each other's eyes makes it better: eye-to-eye contact enhances sexual interactions in wild geladas. *Animal Behaviour*, 2021, 177, pp.269-276. 10.1016/j.anbehav.2021.05.011 . hal-03264410

HAL Id: hal-03264410

<https://hal-univ-rennes1.archives-ouvertes.fr/hal-03264410>

Submitted on 29 Jun 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Looking into each other's eyes makes it better: Eye-to-Eye Contact enhances sexual interactions in geladas

Anna Zanolì^a, Marco Gamba^a, Alban Lemasson^b, Elisabetta Palagi^{c,d,*}, & Ivan Norscia^{a,c,*}

^aUniversity of Turin, Department of Life Sciences and Systems Biology, Italy

^bUniversity of Rennes, University of Normandie, CNRS, EthoS (Éthologie animale et humaine), France

^cNatural History Museum, University of Pisa, Calci, Italy

^dUnit of Ethology, Department of Biology, University of Pisa, Italy

***Corresponding authors:**

Ivan Norscia:

E-mail: ivan.norscia@unito.it

Postal Address: University of Turin, Department of Life Sciences and Systems Biology, Via Accademia Albertina 13, 10123 Torino (Italy)

Telephone: +39 0116704547

Elisabetta Palagi:

E-mail: elisabetta.palagi@unipi.it

Postal Address: Unit of Ethology, Department of Biology, University of Pisa, Via A. Volta 6, 56126 Pisa (Italy)

Telephone: +39 0502211385

In human and non-human primates Eye-to-Eye Contact (EEC), a face-to-face communication component, can promote emotional/attentional engagement, and prolong affiliative interactions. Due to its direct impact on fitness, the reproductive context is perhaps the most critical context for investigating EEC's importance. However, the presence of this phenomenon around mating and its functions in primates is still understudied. In this work, we investigated whether EEC was present during copulations and influenced the copula duration and post-copulation grooming occurrence in the wild gelada (*Theropithecus gelada*), an Old World monkey species. We found that the previous presence of the male 'look-at' triggered the female 'look-at'. Moreover, copulations were most likely to last longer in the presence of EEC. In addition, the occurrence of post-copulation grooming between partners - most frequently initiated by females - increased when copulations included EEC. Females' engagement in EEC with the male may be a form of continuation of female pre-copulatory proceptivity and facilitate males' copulatory activity. EEC by prolonging sexual contacts, may also increase the chances of ejaculation. By grooming their partners after mating, female geladas may attempt to reduce male arousal and prolong the social interaction with them, possibly strengthening their social bond. These results provide the first quantitative evidence that EEC is an effective mechanism for prolonging mating interactions and enhancing post-mating affiliation in a Papionini species. On a broader perspective, the presence of EEC in an Old-World monkey species suggests that EEC may have been favoured by natural selection to promote reproductive advantages during human evolution.

KEYWORDS: Eye-to-Eye Contact; facial communication; mutual gaze; reciprocal looking; sexual behaviour; social bonding; *Theropithecus gelada*; visual communication

62 In evolutionary terms, the measure of an individual's success is the amount of its genes present in
63 subsequent generations (Smith & Maynard-Smith, 1978). Being the critical point of an individual's
64 existence, reproduction is the central theme around which all other aspects of biology revolve
65 (Dunbar, 2014). Among the different forms of reproduction, sexual reproduction is the most common
66 in nature, and it depends on effective communication between senders and receivers (Bell, 1982).
67 Courtship and mating involve the production of sexual signals that transmit crucial information about
68 the senders' identity, quality, social status, and motivation (Bradbury & Vehrencamp, 1998).
69 Depending on the species, the exchange of information in the reproductive context can occur via
70 different sensory modalities (Partan & Marler, 1999; 2005). Although hearing and/or smell can be
71 crucial in primates, vision is pivotal for communication, especially in anthropoids. For example, their
72 relatively large, forward-facing eyes give rise to binocular eyesight fields, enabling stereoscopic
73 vision (Ravosa & Savakova, 2004; Fleagle, 2013). Despite the importance of visual signals in
74 primates (Higham et al., 2011; 2012), eye-gaze behaviour in the reproductive context has not received
75 much attention so far (Dixson, 2012; Liebal, Waller, Slocombe, & Burrows, 2014).

76 In anthropoids, face-to-face communication is important in regulating social interactions such as
77 competition, affiliation, and socio-sexual contacts (Gothard, Erickson, & Amaral, 2004; Parr, Waller,
78 Vick, & Bard, 2007; Micheletta, Whitehouse, Parr, & Waller, 2015; Annicchiarico, Bertini, Cordoni,
79 & Palagi, 2020; for a review see: Waller & Micheletta, 2013). In human and other non-human
80 primates, specific forms of face-to-face communication such as rapid facial mimicry and yawn
81 contagion are associated with enhanced affiliative behaviour and social bonding (Mancini, Ferrari, &
82 Palagi, 2013a; Norscia & Palagi, 2011).

83 Eye-to-Eye Contact (EEC) is a crucial component of face-to-face communication (Kret, Fischer, &
84 De Dreu, 2015; Schino & Sciarretta, 2016). The Cooperative Eye Hypothesis (CEH) predicts that
85 EEC in humans has evolved to maintain cooperative behaviours (Tomasello, Hare, Lehmann, & Call,
86 2007). In non-human primates, EEC can also be an effective way to convey essential information
87 about the subjects' motivation when they engage in social interactions (Wrangham, 1993; Kobayashi

88 & Hashiya, 2011). In this respect, EEC may be a means to promote emotional/attentional engagement,
89 thus prolonging affiliative interactions (Cordell & McGahan, 2004; Prochazkova & Kret, 2017;
90 Annicchiarico et al., 2020).

91 EEC can become particularly critical when reproduction is at stake (Dixon, 2012). In many primate
92 species, spanning monkeys and apes, females can turn to look back and establish eye-to-eye contact
93 with males during copulation, as a possible continuation of pre-copulatory, eye-contact proceptivity
94 (Dixon, 2012; Chevalier-Skolnikoff, 1975). According to previous reports, this visual contact can
95 facilitate male's copulatory activity, enhance its arousal, and prolong the sexual contact, thus
96 improving ejaculation chances (Dixon, 2012; Palagi, Bertini, Annicchiarico, & Cordoni, 2020a).
97 Therefore, sex - due to its direct impact on fitness - is perhaps the most critical context to evaluate
98 the importance of EEC (Dixon, 2012; Palagi et al., 2020a). However, despite its importance, facial
99 communication around mating – that includes EEC - and its implications for social bonding in
100 primates remains understudied (Dixon, 2012).

101 Here, we focused on *Theropithecus gelada* (Hill, 1970) to understand whether EEC was present
102 during copulations and, if so, how it influenced mating interactions. Geladas possess a rich repertoire
103 of facial expressions (Dunbar & Dunbar, 1975) that they use to communicate in different contexts:
104 playful context (*play face* – *full play face*: Palagi & Mancini, 2011; Mancini et al., 2013a; Mancini,
105 Ferrari, & Palagi, 2013b); affiliative context (*yawns*: Leone, Ferrari, & Palagi, 2014; Palagi, Leone,
106 Mancini, & Ferrari, 2009; *lip-smacking*: Gustison, le Roux, & Bergman, 2012); and agonistic context
107 (*yawns*: Leone et al., 2014; *lip-flip*: Lazow & Bergman, 2020). In the mating context both male and
108 female geladas can emit different vocalizations around copulation (Aich, Moos-Heilen, &
109 Zimmermann, 1990; Gustison et al., 2012; Gustison & Bergman, 2017; Gustison, Johnson, Beehner,
110 & Bergman, 2019), but little it is known about the eye-gaze behaviour in this context.

111 To fill this gap, this study aimed at testing the following hypotheses:

112 (1) If visual communication has a role in managing the mating interaction in geladas, we expect that
113 males and females seek EEC with the partner.

114 (2) If EEC contributes to increasing the probability of the ongoing copula's success, we expect the
115 longest copulas to be characterized by the presence of EEC.

116 (3) If EEC enhances post-mating affiliation probability, we expect that grooming (the primary form
117 of affiliation in primates; Dunbar, 1991) between mates is widespread after copulations including
118 EEC.

119

120 METHODS

121

122 *Study Subjects and Data Collection*

123 We conducted this research on the Kundi highland (North Shewa Zone, Amhara Region, Ethiopia
124 N9°40.402' E39°45.060'), regularly frequented by 18 One-Male Units (OMUs) of geladas. Data were
125 collected from January to May 2019, and from December 2019 to February 2020. From two to four
126 observers (A.Z. and three field assistants) observed the visible OMUs every day from 0930 hours to
127 1700 hours, for a total of 658 hours of observation. By using the all-occurrences sampling method
128 (Altmann 1974), all copulations (including possible post-copulation grooming between mates)
129 performed by the visible animals were audio- and video-recorded. Copulations were easily
130 predictable thanks to clearly detectable visual and acoustic sexual invitations (present-rear, genital
131 inspection, and female pre-copulation calls; Dunbar & Dunbar, 1975). Hence, the observers were able
132 to anticipate impending copulations and to record each mating before it began. We made video
133 recordings by using HC-V180 Full HD Panasonic video cameras (optical zoom 50×). We recorded
134 sounds using Zoom H5, OLYMPUS-LS100 and Marantz PMD661 solid-state digital audio recorders
135 built up with Sennheiser ME64 and Sennheiser ME66 microphones with a sampling rate of 96kHz
136 (16-bit depth). We recorded a total of 443 mating events, but, for this study, we could only use a
137 subset of 244 copulations performed by 145 dyads from 18 One-Male units (18 alpha males and 142
138 adult females). The high-quality resolution (1920x1080 Pixel) and the optical zoom (50×) allowed to
139 obtain optimal frames of faces and eyes of the mating subjects. Nevertheless, we had to exclude from

140 the complete dataset all the cases ($N=199$) in which it was impossible to see the interacting
141 individuals' eyes due to distance, limited visibility (e.g., foggy weather), and/or animal position.

142

143 *Operational Definitions and Data Processing*

144 The copulation videos were analysed frame-by-frame via the freeware VideoLAN Client 3.0.11.1
145 (2020; with the extension Jump to Time) whereas the audio-recordings were analysed by using Praat
146 6.0.56 (Boersma & Weenink, 2008). Copulation started when the genital areas of the male and the
147 female entered in contact and ended when one of them spontaneously interrupted the contact. We
148 assigned each copulation to one of the conditions described here below. We defined the condition
149 "no-look" when: (a.1) the male turned its face (and gaze) away from the female, (a.2) the male
150 oriented its face frontally without lowering the head, and (b.1) the female did not turn its head back,
151 (b.2) the female turned its head, but its gaze was not directed at the male. In the condition "male look-
152 at" (a) the male lowered its head and had its gaze directed towards the female, but (b.1) the female
153 did not turn its head back, or (b.2) the female turned its head, but its gaze was not directed at the male.
154 We defined the condition "female look-at" when (a) the female turned its head back and had its gaze
155 directed towards the male, but (b.1) the male turned its face (and gaze) away from the female, (b.2)
156 the male oriented its face frontally without lowering the head. We defined the condition "Eye-to-Eye
157 Contact" (EEC) when the look-at was reciprocated, with male and female looking into each other's
158 eyes. So, the look-at conditions could become an EEC interaction only if one subject looked its
159 partner back. The conditions assigned to each copulatory event were based on the presence/absence
160 of look-at or EEC, not on the gaze duration. Examples of each condition are shown in Figure 1.

161 Both "male look-at" and "female look-at" conditions started when one of the mating subjects looked
162 at the other and ended when one of the subjects interrupted the visual contact. EEC conditions started
163 when both sexes looked into each other eyes and ended when one of the subjects interrupted the visual
164 contact. If a copulation included both look-at and EEC, such copulation fell into the EEC condition.

165 This methodology avoided data pseudo-replication. Since the mean duration of a copulatory event
166 was 10.18 ($\pm$ SD 4.15) seconds, we defined as "post-copulation grooming" each
167 grooming session occurring within 10s of the end of the copulation.

168 Following Roberts, Lu, Bergman, & Beehner (2017), we classified the female status as "oestrus" and
169 "non-oestrus" based on the chest vesicle coverage and turgidity, the chest colour, and the presence of
170 paracallosal vesicles.

171 A.Z. analysed all the videos. Twenty-four randomly selected copulation events (10% of the total
172 sample) were assigned to another observer, expert in gelada behaviour and unaware of the study's
173 aim, to check for inter-observer agreement and reliability over scoring. For each category in which
174 we divided our sample Cohen's kappa values were: no-look = 1 male look-at = 0.95, female look-at
175 = 0.90, and EEC = 1.

176 From each copulation video we extracted the following data: (1) identity of the mating dyad, (2)
177 copula duration, (3) the second when look-at and EEC occurred, (4) occurrence of post-copulatory
178 grooming, and (5) female oestrus status. We used the audio recordings to extract (1) presence/absence
179 of male copulation calls (Aich et al., 1990), (2) the second when each subject started the emission of
180 copulation calls, and (3) presence/absence of male post-copulation call sequences. We extracted a
181 behavioural string for each copulatory event, including the temporal sequence of all behaviours and
182 vocalizations.

183

184 *Statistical Analysis*

185 Preliminarily, we conducted a sequential analysis to evaluate the temporal association of the target
186 behavioural patterns and vocalizations (hereafter "items") during and after copulatory events. We
187 created a string for each copulation, including the items separated by a break symbol. The resulting
188 string represented the ordered concatenation of items as they occurred during copulation. Using the
189 software Behatrix 0.9.11 (Friard, & Gamba, 2020), we generated the flow diagram with the transitions
190 from one item to the next, with the percentage values of transition relative occurrences. Then, we ran

191 a permutation test based on observed counts of the behavioural transitions (“Run random permutation
192 test” Behatrix-function). We permuted the strings 10 000 times (allowing us to achieve an accuracy
193 of 0.001 of the probability values) and we obtained *P*-values for each behavioural transition.

194 The sequential analysis showed that the male look-at occurred more frequently before the female
195 look-at. For this reason, we ran a Generalized Linear Mixed Model (GLMM; “lme4” package: Bates,
196 Mächler, Bolker, & Walker, 2015) in R (R Core Team, 2020; version 4.0.2) to verify which variables
197 could affect the occurrence of the female look-at during copulations. This model included the female
198 look-at (presence/absence) as a binomial response variable. The occurrence of male look-at (presence
199 /absence), male copulation calls (presence/absence), and the female oestrus status (oestrus/non-
200 oestrus) were entered as binomial fixed factors, whereas the dyad identity was entered as a random
201 factor.

202 We ran a second model to investigate whether the presence of EEC affected the copula duration
203 (LMM, family = “gaussian”). The log-transformed copula duration (in seconds) was the response
204 variable, whereas EEC (presence/absence) and male look-at (presence/absence) were the fixed
205 factors, and the dyad identity was the random factor. For this model, we verified the normal
206 distribution and homogeneity of the residuals by looking at the qq-plot and plotting the residuals
207 against the fitted values (Estienne, Mundry, Köhl, & Boesch, 2016).

208 Finally, to verify whether EEC's presence influenced the occurrence of post-copulation grooming, we
209 ran a third GLMM. The occurrence of post-copulation grooming (presence/absence) was the binomial
210 response variable. EEC (presence/absence) and male post-copulation call sequence
211 (presence/absence), and the copula duration were the fixed factors, whereas the dyad identity was the
212 random factor.

213 For all models, we computed multicollinearity with generalised variance inflation factors (GVIF; Fox
214 & Monette, 1992) in R (“vif” function; Fox & Weisberg, 2011). The GVIF revealed no collinearity
215 between fixed factors (< 1.02 in all cases). To test the significance of the models, we compared each
216 full model with a null model including only the random factor (Forstmeier & Schielzeth, 2011), using

217 a likelihood ratio test (Anova with the "Chisq" test argument; Dobson, 2002). Then, we estimated p-
218 values for each predictor based on likelihood ratio tests between the full model and the respective
219 null model (R-function "drop1"; Barr, Levy, Scheepers, & Tily, 2013).

220

221 *Ethical Note*

222 This is a non-invasive research compliant with the ASAB/ABS Guidelines for the Use of Animals in
223 Research, the current Ethiopian Italian and French law and University regulations. Thus, no permit
224 from the Bio-Ethical Committee was needed.

225

226 RESULTS

227

228 *Behavioural Transitions During and After Copulation*

229 The sequential analysis on the behaviours/vocalizations revealed that, during copulations, both males
230 and females emitted copulation calls before looking at each other (transition male copulation calls □
231 male look-at : percentage of occurrence = 9.22%; $P = 0.040$; transition male copulation calls □ female
232 look-at: percentage of occurrence = 14.89%; $P = 0.007$; transition female copulation calls □ male
233 look-at: percentage of occurrence = 13.63%; $P = 0.008$; transition female copulation calls □ female
234 look-at: percentage of occurrence = 24.24%; $P < 0.001$). In addition, most frequently the male was
235 the first to look at the female (transition male look-at □ female look-at: percentage of occurrence
236 = 43.75%; $P < 0.001$). Finally, during copulations EEC was followed by grooming (started by the
237 female) in the 70.58% of the cases ($P < 0.001$). A flow diagram with the significant behavioural
238 transitions is reported in Figure 2a.

239

240 *EEC Presence and Effects During and After Copulation*

241 When investigating which variables affected the female look-at occurrence, we found that the full
242 model significantly differed from the null model ($\chi^2 = 27.519$, $df = 5$, $P < 0.001$; Table 1). The previous

243 presence of male look-at was associated with an increased likelihood of female-look at (Figure 1b),
244 whereas the main effect of male copulation calls did not reach statistical significance. Likewise, the
245 female oestrus status did not affect the occurrence of female look-at.

246 The full model that we built to check whether EEC affected the copula duration significantly differed
247 from the null model ($\chi^2=7.211$, $df=5$, $P=0.027$; Table 2). We found that copulations in which EEC
248 was present lasted significantly more (mean [s] $\pm$ SD = 13.203 ± 4.659) than copulations in which
249 EEC was absent (mean [s] $\pm$ SD = 8.390 ± 2.624) (Figure 1c).

250 Finally, we built a model to investigate whether EEC during copulations influenced the occurrence
251 of post-copulation grooming. The full model significantly differed from the null model
252 ($\chi^2=9.206$, $df=5$, $P=0.026$; Table 3). We found that EEC's presence during copulations was
253 associated with an increased likelihood of post-copulation grooming (Figure 1d). In contrast, male
254 post-copulation call sequences and the copula duration did not have a significant main effect on the
255 target variable.

256

257 DISCUSSION

258 This study investigated whether Eye-to-Eye Contact (EEC) was present during copulation and
259 affected copula duration and post-copulation grooming in wild geladas. We found that during
260 copulations, female look-at was influenced by the previous presence of male look-at but not by the
261 previous emission of male copulation calls (Figure 2a, 2b – in line with Prediction 1). Moreover,
262 copulations were most likely to last longer when EEC was present (Figure 2c – in line with Prediction
263 2), but not when only male look-at occurred. Finally, the probability of post-copulation grooming
264 between partners increased (with grooming most likely started by females) when copulations included
265 EEC (Figure 2a, 2d - in agreement with Prediction 3).

266 These results provide the first quantitative evidence of EEC's presence during copulations in geladas
267 and allow inferences on its potential functions in favouring positive social interactions. As predicted,
268 partners looked at each other, with females being most likely to look at the male after being looked

269 by the male independently from the presence of male copulation calls. Although geladas possess an
270 elaborate vocal repertoire used in the mating context (Aich et al., 1990; Gustison et al., 2012; Gustison
271 & Bergman, 2017; Gustison et al., 2019), in this case, male copulation calls do not seem to be the
272 main trigger of the visual contact (Table 1; Figure 2b). This result suggests that the female look-at
273 was not a simple reaction to male copulation calls, but that the females probably sought for EEC with
274 males. Previous studies showed that in all major radiations of anthropoid primates, including New
275 World monkeys (e.g. *Callimico goeldii*: Heltne, Wojcik, & Pook, 1981; *Callithrix jacchus*: Kendrick
276 & Dixon, 1984; *Leontopithecus rosalia*: Kleiman, Hoage, & Green, 1988; *Brachyteles arachnoides*:
277 Milton, 1985), Old World monkeys (e.g. *Macaca* spp: Hinde & Rowell, 1962; Zumpe & Michael,
278 1968; Dixon, 1977; Wolfe, 1984; Slob & Nieuwenhuijsen, 1980; Slob et al., 1986; Chevalier-
279 Skolnikoff, 1975; in *Lophocebus albigena*: Wallis, 1983; *Papio ursinus*: Saayman,
280 1970; *Miopithecus talapoin*: Dixon, Scruton, & Herbert, 1975) and apes (*Pan paniscus*: Tutin &
281 McGinnis, 1981; Palagi et al., 2020a; *Pan troglodytes*: Goodall, 1986; *Gorilla beringei beringei*:
282 Harcourt, 1981), EEC between partners possibly occurred also during dorso-ventral sexual
283 interactions. As reported for other primate species, gelada females may seek the males' eye-contact
284 to assess males' intent and communicate their engagement. In this respect, females seeking eye
285 contact can, therefore, be interpreted as a form of a possible continuation of pre-copulatory, eye-
286 contact proceptivity (Dixon, 2012).

287 EEC was present during copulations and associated with more prolonged sexual interactions (Figure
288 2c – Prediction 2 supported). Besides, we found that the male look-at's presence did not *per se* affect
289 the copula duration (Table 2). This result allowed us to exclude the possibility that copulation lasted
290 longer because males were generally more “attentive”. In a general perspective, this result is in line
291 with previous findings on the possible function of EEC in prolonging social interactions in humans
292 and apes under different contexts (*Homo sapiens*: Cordell & McGahan, 2004; Prochazkova & Kret,
293 2017; *Pan paniscus*: Annicchiario et al., 2020). More specifically, our findings support the previous,
294 few studies on the possible effect of EEC on mating. Savage-Rumbaugh & Wilkerson (1978)

described that in bonobos, the success of sexual interactions, estimated by their duration, could be associated with maintaining mutual gaze during sexual contacts. More recently, Palagi et al. (2020a) reported that the presence of rapid facial mimicry (a facial mirror response occurring within a second after the perception of other facial expressions; Mancini et al., 2013a; Palagi, Celeghin, Tamietto, Winkielman, & Norscia, 2020b) increased the duration of bonobo hetero-sexual contacts. Female look-at during mating may trigger male pelvic thrusting, which ends with ejaculation (*Brachyteles arachnoides*; Milton, 1985). Thus, we can suppose that also in geladas EEC may facilitate the copulatory activity of males, enhance their sexual arousal and, by prolonging the sexual contact, increase ejaculation chances.

Finally, we found that EEC's presence was associated with an increased occurrence of post-copulation grooming, especially started by females (Figure 2a, 2d). The duration of copulas (a possible proxy of the copulation's success; Milton, 1985) and male post-copulation call sequences did not significantly affect the subsequent occurrence of grooming. Hence, it is unlikely that these two factors were the primary triggers of the post-copulation grooming increase (Table 3). However, we cannot exclude that the co-occurrence of EEC and grooming may be a by-product of the possible link between EEC and ejaculation. Our result supports our third prediction and can be discussed on two levels. In the short term, if EEC's presence during copulations increased the levels of male arousal, females - by grooming the partner - may attempt to reduce such arousal to favour affiliative interactions. Previous findings reported that grooming is effective in reducing arousal-related anxiety in non-human primates, from lemurs to apes (e.g., *Lemur catta*: Sclafani, Norscia, Antonacci, & Palagi, 2012; *Macaca fascicularis*: Schino, Scucchi, Maestripieri, & Turillazzi, 1988; *Pan paniscus*: Palagi & Norscia, 2013; *Pan troglodytes*: De Waal & van Roosmalen, 1979; for a review see: Dunbar, 2010). Similarly, in humans, mutual-grooming may serve to reduce relationship-related anxiety and favour bonding (Nelson & Geher, 2007).

In the longer run, gelada females may try to prolong the social interaction with males and possibly reinforce their social bond with them. In primates, grooming is the predominant form of affiliation

321 used to establish, maintain and strengthen social bonds (Dunbar, 1991). Moreover, in previous studies
322 on human and non-human primates, EEC has been described as an effective mechanism that has
323 evolved to maintain cooperative behaviours and prolong affiliative interactions by promoting
324 emotional/attentional engagement (Tomasello et al., 2007; Cordell & McGahan, 2004; Prochazkova
325 & Kret, 2017; Annicchiarico et al., 2020). This explanation may be especially valid in the light of the
326 characteristics of geladas. In this species, females can benefit from male protection, especially in
327 relation to reproduction, considering that high levels of infanticides have been observed in case of
328 takeover attempts (Mori, Shimizu, & Hayashi, 2003; Beehner & Bergman, 2008; Roberts, Lu,
329 Bergman, & Beehner, 2012; Pallante, Stanyon, & Palagi, 2016). By prolonging the social interaction
330 with males by grooming them after mating, females may reinforce social bonding and increase male
331 protection.

332 The impossibility of evaluating the quality of the relationship between the mating partners may be a
333 limitation of this study. This factor could affect the gaze behaviour during copulations and the
334 grooming rate between the partners and could lead to more comprehensive results. Although long-
335 term studies are necessary to assess EEC's function in strengthening social bonding between male
336 and female geladas, we provided reliable support that EEC represents an effective mechanism to
337 prolong mating interactions (possibly increasing chances of success) and enhance post-copula
338 affiliation in a species of Papionini. More generally, this study confirms that visual communication
339 can function as an aid to reproduction (Liebal et al., 2014). Finally, by focusing on an Old-World
340 monkey species (which separated from the human lineage around 18-22 million years ago; Pozzi et
341 al., 2014), this study suggests that EEC may have long been favoured by natural selection to promote
342 reproductive advantages over the course of human evolution.

343

344 DATA AVAILABILITY

345 Data are available at <https://doi.org/10.5281/zenodo.4434496>.

346 REFERENCES

- 347
348 Aich, H., Moos-Heilen, R., & Zimmermann, E. (1990). Vocalizations of adult gelada baboons
349 (Theropithecus gelada): acoustic structure and behavioural context. *Folia primatologica*, 55(3-4),
350 109-132. <https://doi.org/10.1159/000156508>.
- 351 Altmann, J. (1974). Observational study of behavior: sampling methods. *Behaviour* 49, 227–267.
352 <https://doi.org/10.1163/156853974X00534>.
- 353 Annicchiarico, G., Bertini, M., Cordoni, G., & Palagi, E. (2020). Look at me while having sex!
354 Eye-to-eye contact affects homosexual behaviour in bonobo females. *Behaviour*, 1(aop), 1-22.
355 <https://doi.org/10.1163/1568539X-bja10034>.
- 356 Barr, D. J., Levy, R., Scheepers, C., & Tily, H. J. (2013). Random effects structure for confirmatory
357 hypothesis testing: keep it maximal. *Journal of Memory and Language*, 68, 255–278.
358 <https://doi.org/10.1016/j.jml.2012.11.001>.
- 359 Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed- Effects Models
360 using lme4. *Journal of Statistical Software*, 67, 1–48. <https://doi.org/10.18637/jss.v067.i01>.
- 361 Beehner, J. C., & Bergman, T. J. (2008). Infant mortality following male takeovers in wild
362 geladas. *American Journal of Primatology: Official Journal of the American Society of*
363 *Primatologists*, 70(12), 1152-1159. <https://doi.org/10.1002/ajp.20614>.
- 364 Bell, G. (1987). Two theories of sex and variation. In *The evolution of sex and its consequences* (pp.
365 117-134). Birkhäuser, Basel. https://doi.org/10.1007/978-3-0348-6273-8_5.
- 366 Boersma, P., & Weenink, D. (2008). Praat: Doing Phonetics by Computer (Computer Program).
367 Retrieved from <https://www.fon.hum.uva.nl/praat/>. Accessed December 22, 2020

368 Bradbury, J. W., & Vehrencamp, S. L. (1998). Principles of animal communication. Sinauer
 369 Associates.

370 Brunson, J. C., Read, Q. D. (2020). ggalluvial: Alluvial Plots in 'ggplot2'. R package version 0.12.3.
 371 Retrieved from <http://corybrunson.github.io/ggalluvial/>. Accessed December 28, 2020

372 Chevalier-Skolnikoff, S. (1975). Heterosexual copulatory patterns in stumptail macaques (*Macaca*
 373 *arctoides*) and in other macaque species. *Archives of sexual behavior*, 4(2), 199-220.
 374 <https://doi.org/10.1007/BF01541083>.

375 Cordell, D. M., & McGahan, J. R. (2004). Mutual Gaze Duration as a Function of Length of
 376 Conversation in Male—Female Dyads. *Psychological reports*, 94(1), 109-114.
 377 <https://doi.org/10.2466%2Fpr0.94.1.109-114>.

378 De Waal, F. B., & van Roosmalen, A. (1979). Reconciliation and consolation among
 379 chimpanzees. *Behavioral Ecology and Sociobiology*, 5(1), 55-66.
 380 <https://doi.org/10.1007/BF00302695>.

381 Dixon, A. F. (1977). Observations on the displays, menstrual cycles and sexual behaviour of the
 382 “Black ape” of Celebes (*Macaca nigra*). *Journal of Zoology*, 182(1), 63-84.
 383 <https://doi.org/10.1111/j.1469-7998.1977.tb04141.x>.

384 Dixon, A. F., Scruton, D. M., & Herbert, J. (1975). Behaviour of the talapoin monkey
 385 (*Miopithecus talapoin*) studied in groups, in the laboratory. *Journal of Zoology*, 176(2), 177-210.
 386 <https://doi.org/10.1111/j.1469-7998.1975.tb03192.x>.

387 Dixon, A. F., (2012). *Primate Sexuality: Comparative Studies of the Prosimians, Monkeys, Apes,*
 388 *and Humans*. Oxford University Press.

389 Dobson, A. J. (2002). *An Introduction to Generalized Linear Models*. (2nd ed.). Boca Raton, FL:
 390 Chapman and Hall/CRC Press.

391 Dunbar, R. I. (1991). Functional significance of social grooming in primates. *Folia*
 392 *primatologica*, 57(3), 121-131. <https://doi.org/10.1159/000156574>.

393 Dunbar, R. I. (2010). The social role of touch in humans and primates: behavioural function and
 394 neurobiological mechanisms. *Neuroscience & Biobehavioral Reviews*, 34(2), 260-268.
 395 <https://doi.org/10.1016/j.neubiorev.2008.07.001>.

396 Dunbar, R. I. M. (2014). *Reproductive decisions: an economic analysis of gelada baboon social*
 397 *strategies*. Princeton, New Jersey: Princeton University Press.

398 Dunbar, R. I. M. & Dunbar, P. (1975). Social dynamics of gelada baboons. In H. Kuhn, W. P.
 399 Luckett, C. R. Noback, A. H. Schultz, D. Starck & F. S. Szalay (Eds.) *Contributions to Primatology*
 400 (pp. 1-157). Basel: S. Karger.

401 Estienne, V., Mundry, R., Köhl, H. S., & Boesch, C. (2017). Exploitation of underground bee nests
 402 by three sympatric consumers in Loango National Park, Gabon. *Biotropica*, 49(1), 101-109.
 403 <https://doi.org/10.1111/btp.12354>.

404 Fleagle, J. G. (2013). *Primate adaptation and evolution*. Academic press.

405 Forstmeier, W., & Schielzeth, H. (2011). Cryptic multiple hypotheses testing in linear models:
 406 overestimated effect sizes and the winner's curse. *Behavioral Ecology and Sociobiology*, 65, 47–55.
 407 <https://doi.org/10.1007/s00265-010-1038-5>.

408 Fox, J., & Monette, G. (1992). Generalized collinearity diagnostics. *Journal of the American*
 409 *Statistical Association*, 87(417), 178-183. <https://doi.org/10.1080/01621459.1992.10475190>.

410 Fox, J., & Weisberg, S. (2011). *An R Companion to Applied Regression*. (2nd ed). Thousand Oaks,
 411 CA: SAGE Publications, Inc.

412 Friard, O., Gamba, M., (2020). Behatrix: Behavioral sequences analysis with permutations test.
 413 Retrieved from <http://www.boris.unito.it/pages/behatrix>. Accessed January 04, 2021

414 Goodall, J. (1986). *The chimpanzees of Gombe: Patterns of behavior*. Cambridge Mass.

415 Gothard, K. M., Erickson, C. A., & Amaral, D. G. (2004). How do rhesus monkeys (*Macaca*
 416 *mulatta*) scan faces in a visual paired comparison task?. *Animal cognition*, 7(1), 25-36.
 417 <https://doi.org/10.1007/s10071-003-0179-6>.

418 Gustison, M. L., & Bergman, T. J. (2017). Divergent acoustic properties of gelada and baboon
 419 vocalizations and their implications for the evolution of human speech. *Journal of Language*
 420 *Evolution*, 2(1), 20-36. <https://doi.org/10.1093/jole/lzx015>.

421 Gustison, M. L., le Roux, A., & Bergman, T. J. (2012). Derived vocalizations of geladas
 422 (*Theropithecus gelada*) and the evolution of vocal complexity in primates. *Philosophical*
 423 *Transactions of the Royal Society B: Biological Sciences*, 367(1597), 1847-1859.
 424 <https://doi.org/10.1098/rstb.2011.0218>.

425 Gustison, M. L., Johnson, E. T., Beehner, J. C., & Bergman, T. J. (2019). The social functions of
 426 complex vocal sequences in wild geladas. *Behavioral Ecology and Sociobiology*, 73(1), 14.
 427 <https://doi.org/10.1007/s00265-018-2612-5>.

428 Harcourt, A. H. (1981). Gorilla reproduction in the wild. *Reproductive biology of the great apes*.

429 Heltne, P. G., Wojcik, J. F., & Pook, A. G. (1981). Goeldi's monkey, genus *Callimico*. In *Ecology*
 430 *and Behaviour of Neotropical Primates* (Vol. I, pp. 169–221). Rio de Janeiro: Academia Brasileira
 431 de Ciências

432 Higham, J. P., Hughes, K. D., Brent, L. J., Dubuc, C., Engelhardt, A., Heistermann, M., ... &
 433 Stevens, M. (2011). Familiarity affects the assessment of female facial signals of fertility by free-
 434 ranging male rhesus macaques. *Proceedings of the Royal Society B: Biological*
 435 *Sciences*, 278(1723), 3452-3458. <https://doi.org/10.1098/rspb.2011.0052>.

436 Higham, J. P., Heistermann, M., Saggau, C., Agil, M., Perwitasari-Farajallah, D., & Engelhardt, A.
 437 (2012). Sexual signalling in female crested macaques and the evolution of primate fertility
 438 signals. *BMC evolutionary biology*, 12(1), 1-10. <https://doi.org/10.1186/1471-2148-12-89>.

439 Hill, W. C. O. (1970). Primates, comparative anatomy and taxonomy. Vol. VIII. Cynopithecinae,
 440 Papio, Mandrillus, Theropithecus.

441 Hinde, R. A., & Rowell, T. E. (1962, January). Communication by postures and facial expressions
 442 in the rhesus monkey (*Macaca mulatta*). In *Proceedings of the Zoological Society of London*(Vol.
 443 138, No. 1, pp. 1-21). Oxford, UK: Blackwell Publishing Ltd. [https://doi.org/10.1111/j.1469-](https://doi.org/10.1111/j.1469-7998.1962.tb05684.x)
 444 [7998.1962.tb05684.x](https://doi.org/10.1111/j.1469-7998.1962.tb05684.x).

445 Kendrick, K. M., & Dixson, A. F. (1984). A quantitative description of copulatory and associated
 446 behaviors of captive marmosets (*Callithrix jacchus*). *International journal of primatology*, 5(3),
 447 199-212. <https://doi.org/10.1007/BF02735757>.

448 Kleiman, D. G., Hoage, R. J., & Green, K. M. I., (1988). Description of species. In *Ecology and*
 449 *Behavior of Neotropical Primates* (pp. 2-299).

450 Kret, M. E., Fischer, A. H., & De Dreu, C. K. (2015). Pupil mimicry correlates with trust in in-
 451 group partners with dilating pupils. *Psychological science*, 26(9), 1401-1410.
 452 <https://doi.org/10.1177%2F0956797615588306>.

453 Kobayashi, H., & Hashiya, K. (2011). The gaze that grooms: contribution of social factors to the
 454 evolution of primate eye morphology. *Evolution and Human Behavior*, 32(3), 157-165.
 455 <https://doi.org/10.1016/j.evolhumbehav.2010.08.003>.

456 Lazow, S. P., & Bergman, T. J. (2020). The structural and motivational role of the unique lip-flip
 457 movement in the gelada (*Theropithecus gelada*) facial display repertoire. *American Journal of*
 458 *Physical Anthropology*, 172(2), 280-290. <https://doi.org/10.1002/ajpa.24031>.

459 Leone, A., Ferrari, P. F., & Palagi, E. (2014). Different yawns, different functions? Testing social
 460 hypotheses on spontaneous yawning in *Theropithecus gelada*. *Scientific reports*, 4, 4010.
 461 <https://doi.org/10.1038/srep04010>.

462 Liebal, K., Waller, B. M., Slocombe, K. E., & Burrows, A. M. (2014). *Primate communication: a*
 463 *multimodal approach*. Cambridge University Press.

464 Mancini, G., Ferrari, P. F., & Palagi, E. (2013a). Rapid facial mimicry in geladas. *Scientific*
 465 *reports*, 3(1), 1-6. <https://doi.org/10.1038/srep01527>.

466 Mancini, G., Ferrari, P. F., & Palagi, E. (2013b). In play we trust. Rapid facial mimicry predicts the
 467 duration of playful interactions in geladas. *PloS one*, 8(6), e66481.
 468 <https://doi.org/10.1371/journal.pone.0066481>.

469 Micheletta, J., Whitehouse, J., Parr, L. A., & Waller, B. M. (2015). Facial expression recognition in
 470 crested macaques (*Macaca nigra*). *Animal Cognition*, 18(4), 985-990.
 471 <https://doi.org/10.1007/s10071-015-0867-z>.

472 Milton, K. (1985). Mating patterns of woolly spider monkeys, *Brachyteles arachnoides*:
 473 implications for female choice. *Behavioral Ecology and Sociobiology*, 17(1), 53-59.
 474 <https://doi.org/10.1007/BF00299429>.

475 Mori, T., Shimizu, K., & Hayashi, M. (2003). Levels of serum brain-derived neurotrophic factor in
 476 primates. *Primates*, 44(2), 167-169. <https://doi.org/10.1007/s10329-002-0015-7>.

477 Nelson, H., & Geher, G. (2007). Mutual grooming in human dyadic relationships: an ethological
 478 perspective. *Current Psychology*, 26(2), 121-140. <https://doi.org/10.1007/s12144-007-9009-3>.

479 Norscia, I., & Palagi, E. (2011). Yawn contagion and empathy in Homo sapiens. *PloS one*, 6(12),
 480 e28472. <https://doi.org/10.1371/journal.pone.0028472>.

481 Palagi, E., Leone, A., Mancini, G., & Ferrari, P. F. (2009). Contagious yawning in gelada baboons
 482 as a possible expression of empathy. *Proceedings of the National Academy of Sciences*, 106(46),
 483 19262-19267. <https://doi.org/10.1073/pnas.0910891106>.

484 Palagi, E., & Mancini, G. (2011). Playing with the face: Playful facial “chattering” and signal
 485 modulation in a monkey species (*Theropithecus gelada*). *Journal of Comparative*
 486 *Psychology*, 125(1), 11. <https://psycnet.apa.org/doi/10.1037/a0020869>.

487 Palagi, E., & Norscia, I. (2013). Bonobos protect and console friends and kin. *PloS One*, 8(11),
 488 e79290. <https://doi.org/10.1371/journal.pone.0079290>.

489 Palagi, E., Bertini, M., Annicchiarico, G., & Cordoni, G. (2020a). Mirror replication of sexual facial
 490 expressions increases the success of sexual contacts in bonobos. *Scientific reports*, 10(1), 1-11.
 491 <https://doi.org/10.1038/s41598-020-75790-3>.

492 Palagi, E., Celeghin, A., Tamietto, M., Winkielman, P., & Norscia, I. (2020b). The neuroethology
 493 of spontaneous mimicry and emotional contagion in human and non-human animals. *Neuroscience*
 494 *& Biobehavioral Reviews*, 111, 149-165. <https://doi.org/10.1016/j.neubiorev.2020.01.020>.

495 Pallante, V., Stanyon, R., & Palagi, E. (2016). Agonistic support towards victims buffers aggression
 496 in geladas (*Theropithecus gelada*). *Behaviour*, 153(9-11), 1217-1243.
 497 <https://doi.org/10.1163/1568539X-00003369>.

498 Parr, L. A., Waller, B. M., Vick, S. J., & Bard, K. A. (2007). Classifying chimpanzee facial
 499 expressions using muscle action. *Emotion*, 7(1), 172. [https://psycnet.apa.org/doi/10.1037/1528-](https://psycnet.apa.org/doi/10.1037/1528-3542.7.1.172)
 500 [3542.7.1.172](https://psycnet.apa.org/doi/10.1037/1528-3542.7.1.172).

501 Partan, S., & Marler, P. (1999). Communication goes multimodal. *Science*, 283(5406), 1272-1273.
 502 <https://doi.org/10.1126/science.283.5406.1272>.

503 Partan, S. R., & Marler, P. (2005). Issues in the classification of multimodal communication
 504 signals. *The American Naturalist*, 166(2), 231-245. <https://doi.org/10.1086/431246>.

505 Pozzi, L., Hodgson, J. A., Burrell, A. S., Sterner, K. N., Raaum, R. L., & Disotell, T. R. (2014).
 506 Primate phylogenetic relationships and divergence dates inferred from complete mitochondrial
 507 genomes. *Molecular phylogenetics and evolution*, 75, 165-183.
 508 <https://doi.org/10.1016/j.ympev.2014.02.023>.

509 Prochazkova, E., & Kret, M. E. (2017). Connecting minds and sharing emotions through mimicry:
 510 A neurocognitive model of emotional contagion. *Neuroscience & Biobehavioral Reviews*, 80, 99-
 511 114. <https://doi.org/10.1016/j.neubiorev.2017.05.013>.

512 R Core Team (2020). R: A language and environment for statistical computing. R Foundation for
 513 Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.

514 Ravosa, M. J., & Savakova, D. G. (2004). Euprimate origins: the eyes have it. *Journal of human*
 515 *evolution*, 46(3), 357-364.

516 Roberts, E. K., Lu, A., Bergman, T. J., & Beehner, J. C. (2012). A Bruce effect in wild
 517 geladas. *Science*, 335(6073), 1222-1225. <https://doi.org/10.1126/science.1213600>.

518 Roberts, E. K., Lu, A., Bergman, T. J., & Beehner, J. C. (2017). Female reproductive parameters in
 519 wild geladas (*Theropithecus gelada*). *International Journal of Primatology*, 38(1), 1-20.
 520 <https://doi.org/10.1007/s10764-016-9939-4>.

521 Saayman, G. S. (1970). The menstrual cycle and sexual behaviour in a troop of free ranging chacma
 522 baboons (*Papio ursinus*). *Folia Primatologica*, 12(2), 81-110. <https://doi.org/10.1159/000155283>.

523 Savage-Rumbaugh, E. S., & Wilkerson, B. J. (1978). Socio-sexual behavior in *Pan paniscus* and
 524 *Pan troglodytes*: a comparative study. *Journal of Human Evolution*, 7(4), 327-IN6.
 525 [https://doi.org/10.1016/S0047-2484\(78\)80074-8](https://doi.org/10.1016/S0047-2484(78)80074-8).

526 Schino, G., Scucchi, S., Maestripieri, D., & Turillazzi, P. G. (1988). Allogrooming as a tension-
 527 reduction mechanism: a behavioral approach. *American journal of primatology*, 16(1), 43-50.
 528 <https://doi.org/10.1002/ajp.1350160106>.

529 Schino, G., & Sciarretta, M. (2016). Patterns of social attention in mandrills, *Mandrillus*
 530 *sphinx*. *International Journal of Primatology*, 37(6), 752-761. [https://doi.org/10.1007/s10764-016-](https://doi.org/10.1007/s10764-016-9936-7)
 531 [9936-7](https://doi.org/10.1007/s10764-016-9936-7).

532 Sclafani, V., Norscia, I., Antonacci, D., & Palagi, E. (2012). Scratching around mating: factors
 533 affecting anxiety in wild *Lemur catta*. *Primates*, 53(3), 247-254. [https://doi.org/10.1007/s10329-](https://doi.org/10.1007/s10329-012-0294-6)
 534 [012-0294-6](https://doi.org/10.1007/s10329-012-0294-6).

535 Slob, A. K., & Nieuwenhuijsen, K. (1980). Heterosexual interactions of pairs of laboratory-housed
 536 stump-tail macaques (*Macaca arctoides*) under continuous observation with closed-circuit video
 537 recording. *International Journal of Primatology*, 1(1), 63-80. <https://doi.org/10.1007/BF02692258>.

538 Smith, J. M., & Maynard-Smith, J. (1978). *The evolution of sex* (Vol. 4). Cambridge: Cambridge
 539 University Press. <https://doi.org/10.1017/S0016672300018693>.

540 Tomasello, M., Hare, B., Lehmann, H., & Call, J. (2007). Reliance on head versus eyes in the gaze
 541 following of great apes and human infants: the cooperative eye hypothesis. *Journal of human*
 542 *evolution*, 52(3), 314-320. <https://doi.org/10.1016/j.jhevol.2006.10.001>.

543 Tutin, C. E., & McGinnis, P. R. (1981). Chimpanzee reproduction in the wild. *Reproductive biology*
 544 *of the great apes: Comparative and biomedical perspectives*, 239-264.

545 Video Lan, (2020). VLC media player. <https://www.videolan.org/vlc/index.html>

546 Waller, B. M., & Micheletta, J. (2013). Facial expression in nonhuman animals. *Emotion*
 547 *Review*, 5(1), 54-59. <https://doi.org/10.1177%2F1754073912451503>.

548 Wallis, S. J. (1983). Sexual behavior and reproduction of *Cercocebus albigena johnstonii* in Kibale
 549 forest, Western Uganda. *International journal of primatology*, 4(2), 153-166.
 550 <https://doi.org/10.1007/BF02743755>.

551 Wilke, C. O. (2018). Ggribes: Ridgeline plots in 'ggplot2'. *R package version 0.5, 1*.

552 Wolfe, L. D. (1984). Female rank and reproductive success among Arashiyama B Japanese
 553 macaques (*Macaca fuscata*). *International Journal of Primatology*, 5(2), 133-143.
 554 <https://doi.org/10.1007/BF02735737>.

555 Wrangham, R. W. (1993). The evolution of sexuality in chimpanzees and bonobos. *Human*
 556 *Nature*, 4(1), 47-79. <https://doi.org/10.1007/BF02734089>.

557 Zumpe, D., & Michael, R. P. (1968). The clutching reaction and orgasm in the female rhesus
 558 monkey (*Macaca mulatta*). *Journal of Endocrinology*, 40(1), 117-123.
 559 <https://doi.org/10.1677/joe.0.0400117>.

560

561 **Table 1.** Results of the GLMM showing which variables affected the occurrence of female look-at
 562 during copulations.

Fixed Effects	Estimate	SE	df	z	P
(Intercept)	-2.199	0.693	a	-3.187	a
Male look-at (Presence) ^{b,c}	2.285	0.475	1	4.837	0.000
Male copulation call (Presence) ^{b,c}	-0.174	0.622	1	-0.384	0.782
Female status (oestrus) ^{b,c}	0.649	0.446	1	1.466	0.141

563

*a*Not shown as not having a meaningful interpretation.

564

*b*Estimate ± SE refer to the difference of the response between the reported level of this categorical predictor and the reference category of the same predictor.

565

566

*c*These predictors were dummy coded, with the “Male look-at (Absence)”, “Male copulation call (Absence)”, and “Female status (non oestrus)” being the reference categories.

567

568

569 **Table 2.** Results of the LMM showing the effect of EEC and male look-at on the copula duration.

Fixed Effects	Estimate	SE	df	t	P
(Intercept)	2.212	0.030	a	74.022	a
Eye-to-eye Contact (Presence) ^{b,c}	0.230	0.085	1	2.700	0.007
Male look-at (Presence) ^{b,c}	0.038	0.105	1	0.363	0.722

570

*a*Not shown as not having a meaningful interpretation.

571

*b*Estimate ± SE refer to the difference of the response between the reported level of this categorical predictor and the reference category of the same predictor.

572

573

*c*These predictors were dummy coded, with the “Eye-to-eye Contact (Absence)” and “Male look-at (Absence)” being the reference category.

574

575 **Table 3.** Results of the GLMM showing which variables influenced the occurrence of post-copulation
 576 grooming.

Fixed Effects	Estimate	SE	df	z	P
(Intercept)	-1.084	0.554	a	-1.956	a
Eye-to-eye Contact (Presence) ^{b,c}	1.062	0.498	1	2.134	0.028
Copula duration	0.06	0.038	1	1.577	0.102
Male post-copulation call seq. (Presence) ^{b,c}	-0.127	0.451	1	-0.282	0.779

577

*a*Not shown as not having a meaningful interpretation.

578

*b*Estimate ± SE refer to the difference of the response between the reported level of this categorical predictor and the reference category of the same predictor.

579

580

*c*These predictors were dummy coded, with the “Eye-to-eye Contact (Absence)” and “Male post-copulation call seq. (Absence)” being the reference categories.

581

582

583

584 **Figure 1.** [2-column fitting image] Pictures showing the four gaze conditions. (a): no-look condition;

585 (b): male look-at condition; (c): female look-at condition; (d): EEC condition.

Figure 2. (a): Flow diagram representing the transitions and the percentage of occurrence between each behaviour/vocalization and the proceeding one. Asterisks indicate significance values ($P \leq 0.001 = ***$; $P < 0.01 = **$; $P < 0.05 = *$). Round arrowheads indicate that previous behaviours can be part of the following behaviour. Dashed line indicates the non-significant transition between EEC and Male initiated grooming ($P=0.129$). (b): Percentage of the presence of Female look-at in relation with Male look-at occurrence. Dark-grey bars indicate the absence of Female look-at; pink bars indicate the presence of Female look-at. (c): Raincloud ridge plot, drawn with the R package “ggridges” (Wilke, 2018), showing the copula duration (s) when EEC was present (orange density curves) and when it was absent (blue density curves) in the 18 OMUs studied. Individual observations are presented under the density curves with pipe symbols. (d): Alluvial plot (R package “ggalluvial”; Brunson & Read, 2020) showing the percentage of presence of post-

598 copulation grooming in the presence of EEC during copulation (orange bars) and absence of EEC
599 (blue bars) during copulation.

Accepted manuscript